

UNITED STATES DEPARTMENT OF AGRICULTURE
STATISTICAL BULLETIN No. 47

Washington, D.C.

★

January 1935

GRADE, STAPLE LENGTH, AND TENDERABILITY
OF COTTON IN THE UNITED STATES
1928-29 TO 1932-33

Prepared in the Division of Cotton Marketing
Bureau of Agricultural Economics
In Cooperation with State Agricultural Agencies



STATEMENT OF COOPERATION

The data used in the preparation of this report were collected in cooperation with the following:

Alabama Agricultural Experiment Station.
Arizona Agricultural Experiment Station.
Arkansas Agricultural Experiment Station.
Florida Agricultural Experiment Station.
Georgia Experiment Station of the University System.
Louisiana Agricultural Experiment Station.
Mississippi Agricultural Experiment Station.
Missouri Agricultural Experiment Station.
New Mexico Agricultural Experiment Station.
North Carolina Agricultural Experiment Station.
Oklahoma Agricultural Experiment Station.
South Carolina Agricultural Experiment Station.
Tennessee Agricultural Experiment Station.
Texas Agricultural Experiment Station.
California State Department of Agriculture.

UNITED STATES DEPARTMENT OF AGRICULTURE

STATISTICAL BULLETIN No. 47

Washington, D.C.



January 1935

GRADE, STAPLE LENGTH, AND TENDERABILITY OF COTTON IN THE UNITED STATES, 1928-29 TO 1932-33

*Prepared in the Division of Cotton Marketing, Bureau of Agricultural Economics¹
in cooperation with State agricultural agencies*

CONTENTS

	Page
Introduction.....	1
Grade and staple length of American upland cotton ginned, by States, crop of 1932.....	3
Grade and staple length of American-Egyptian cotton ginned, crops of 1928-32.....	15
Staple length of American upland cotton ginned, by States, crops of 1928-32.....	16
Average staple length of American upland cotton ginned, by States and districts thereof, crops of 1928-32.....	19
Grade of American upland cotton ginned, by States and specified periods, crops of 1928-32.....	22
Grade of American-Egyptian cotton ginned, by specified periods, crops of 1928-32.....	70
Staple length of American upland cotton ginned, by States and specified periods, crops of 1928-32.....	71
Staple length of American-Egyptian cotton ginned, by specified periods, crops of 1928-32.....	85
Staple length of American upland cotton ginned, by districts (10 States), crops of 1928-32.....	86
Tenderability of cotton ginned, by States, crops of 1928-32.....	106
Grade and staple length of American upland cotton on hand, August 1, 1932 and 1933.....	108
Tenderability of American upland cotton on hand, August 1, 1928-33.....	108
Grade and staple length of American-Egyptian cotton on hand, August 1, 1928-33.....	110
Staple length of American upland cotton on hand, August 1, 1928-33.....	111
Staple length of foreign cotton on hand August 1, 1928-33.....	112

INTRODUCTION

Information on the grade, staple length, and tenderability of cotton in the United States has been gathered regularly by the United States Department of Agriculture since 1927. It was during that year that Congress, responding to the demand for information on grade and staple of cotton produced in the United States and on hand on August 1, enacted legislation (act of Mar. 3, 1927, 44 Stat. 1372-1374) authorizing and directing the Secretary of Agriculture to make a report on the grade and staple of cotton on hand at the beginning of each cotton season (Aug. 1) and not less than three reports on the grade and staple of the current crop. It was further provided that in any published statistics the cotton recognized as being tenderable on contracts of sale for future delivery under the Cotton Futures Act of August 11, 1916, as amended, should be stated separately from that which may be untenderable under this act as amended. Funds were made available for the fiscal year beginning July 1, 1928, and at that time the Division of Cotton Marketing, Bureau of Agricultural

¹ By W. B. Lanham, senior agricultural economist; J. T. Egan, senior agricultural economist; F. H. Harper, agricultural economist; F. E. Nelson, assistant marketing specialist; and J. L. McCollum, assistant agricultural economist. Credit is due Arthur W. Palmer for general supervision and helpful suggestions; coworkers; the agricultural experiment stations and the one State department of agriculture in the cooperating States; and the cotton ginners, warehousemen, dealers, and manufacturers whose cooperation made this report possible.

Economics, began active work on the project for the entire Cotton Belt.

The information contained in this bulletin supplements in some respects that contained in Statistical Bulletin 40² and presents, in addition, detailed figures on the grade, staple length, and tenderability of cotton ginned from the 1932 crop and of cotton on hand on August 1, 1932 and 1933. Although much of the information herein presented has been released in preliminary reports, this is the first publication issued by the Department of Agriculture in which detailed information on this subject is assembled for the United States as a whole for 5 consecutive years. A statement regarding the procurement of samples, their classification according to official cotton standards,³ and of the method of preparing the reports on grade, staple length, and tenderability is contained in Statistical Bulletin 40.

There are altogether 16 States for which the Bureau of the Census reported separately the ginnings of cotton from each of the crops of 1928 to 1932, inclusive. In addition there is a small group of States, including Illinois, Kansas, and Kentucky, for which the production is not reported separately.

Statistics on the quantity of cotton on hand in the United States at the beginning of the cotton year (Aug. 1) are also prepared by the Bureau of the Census. Statistics herein presented indicate the quality of this cotton for the period to which this bulletin relates. Information on the grade and staple length of cotton on hand at the beginning of the year in relation to the grade and staple length of the supply is contained in Statistical Bulletin 45,⁴ which shows distribution by types of storage.

Changes that have taken place in the grade and staple length of cotton ginned from the crops of 1928 to 1932, inclusive, in individual producing States and in the United States as a whole are reflected by the statistical information herein presented. Although this period is too short to indicate definite long-time trends, certain changes which seem to indicate tendencies have taken place and are important.

Since 1929 there has been a decrease in both the number of bales of cotton shorter than seven-eighths inch and the proportion that cotton of this length group constitutes of total ginnings. This decrease in production of the shorter lengths has been accompanied by marked increases, on the whole, in the production of most of the lengths above seven-eighths inch, and this increase in production of longer cotton has resulted in an appreciable increase in the average length of staple for the United States and for most of the individual producing States (table 4 and fig. 1, pp. 19 and 21).

The most significant decreases in the production of cotton shorter than seven-eighths inch during the 5-year period 1928-32 occurred in western and eastern Texas, southwestern Oklahoma, southern and

² UNITED STATES DEPARTMENT OF AGRICULTURE, BUREAU OF AGRICULTURAL ECONOMICS, GRADE, STAPLE LENGTH, AND TENDERABILITY OF COTTON IN THE UNITED STATES, 1928-29 TO 1931-32. U. S. Dept. Agr. Statis. Bull. 40, 158 pp., illus. 1933.

³ Cotton standardization is reported to have had its early beginnings about 1800 in Liverpool. Permissive standards were established for grade in this country in February 1909, by the Department of Agriculture. The first official standards were promulgated on Dec. 15, 1914, under the original Cotton Futures Act of Aug. 18, 1914. This statute was reenacted with certain changes in 1916, and the official standards of 1914 were repromulgated. Since that time various revisions have been made and additional standards added, both for grade and for staple length, under the Cotton Futures Act and the Cotton Standards Act of 1923. Universal standards for grade and color of American upland cotton were promulgated July 30, 1923, becoming effective Aug. 1, 1924. These are still in effect.

⁴ LANHAM, W. B., and WEAVER, O. T. GRADE AND STAPLE LENGTH OF COTTON CARRIED OVER IN THE UNITED STATES AS RELATED TO THE DOMESTIC SUPPLY, 1928-29 to 1931-32. U. S. Dept. Agr. Statis. Bull. 45, 23 pp., illus. 1934.

north-central Arkansas, southern Tennessee, eastern Mississippi, and in Alabama, Georgia, and the Carolinas. Important increases in the production of some of the staple lengths longer than seven-eighths inch occurred in the Carolinas, Tennessee, Arkansas, Mississippi, Louisiana, Arizona, California, and Missouri. The extent of these changes and the extent of changes in staple length of production in other States are indicated by the distribution of ginnings presented in the tables.

American upland cotton and American-Egyptian cotton are reported separately in the statistics on grade and staple length of ginnings and in the statistics on grade and staple length of cotton on hand on August 1. The foreign growths are also reported separately in the statistics of cotton on hand on August 1.

The grades and staples untenderable in settlement of contracts of sale for future delivery made subject to section 5 of the United States Cotton Futures Act and regulations of the Secretary of Agriculture thereunder are marked by footnotes in each table where they occur.

In some of the tables, percentages, computed to the nearest 0.1 percent, are presented for the convenience of readers.

For economy of space in printing the tables showing statistics on grades of cotton, the color and grade designations for which there are no quantities to report have been omitted.

The grade names have been abbreviated in the following tables as follows: 1-M.F., Middling Fair; 2-S.G.M., Strict Good Middling; 3-G.M., Good Middling; 4-S.M., Strict Middling; 5-M., Middling; 6-S.L.M., Strict Low Middling; 7-L.M., Low Middling; 8-S.G.O., Strict Good Ordinary; 9-G.O., Good Ordinary.

TABLE 1.—*Grade and staple length of American upland cotton: Quantities ginned in the United States, by States, crop of 1932*

[Quantities are given in running bales, except that round bales are counted as half bales. Linters are not included]

UNITED STATES

Grade	All staple lengths		Shorter than $\frac{7}{8}$ inch ¹	$\frac{7}{8}$ and $\frac{2\frac{1}{2}}{8}$ inch	$1\frac{1}{8}$ and $1\frac{1}{2}$ inch	1 and $1\frac{1}{2}$ inches	$1\frac{1}{8}$ and $1\frac{3}{8}$ inches	$1\frac{1}{8}$ and $1\frac{3}{8}$ inches	$1\frac{1}{8}$ and $1\frac{3}{8}$ inches	$1\frac{1}{8}$ inches and longer
All grades.....	1,000 bales 212,701.3	Per cent 100.0	1,000 bales 837.7	1,000 bales 4,786.5	1,000 bales 3,671.0	1,000 bales 1,822.0	1,000 bales 871.8	1,000 bales 622.1	1,000 bales 84.5	1,000 bales 5.7
Extra White.....	460.0	3.6	5.6	63.5	112.0	88.2	110.0	79.7	1.0	-----
Above 3-G.M.....	1.8 106.9	(²) 8	5	2.1	3.7	8.0	.7	1.1	-----	-----
3-G.M.....	132.8	1.1	1.1	9.8	20.6	25.8	45.7	46.6	.3	-----
4-S.M.....	88.2	.7	1.7	15.0	33.5	23.9	47.0	28.3	.2	-----
5-M.....	99.0	.8	1.5	24.4	42.7	25.0	11.5	1.9	.1	-----
6-S.L.M.....	28.9	.2	.8	11.4	10.7	4.7	4.2	1.5	.3	-----
Below 7-L.M. ¹	1.8 (²)	-----	.2	.8	.8	.8	.9	.3	.1	-----
White.....	9,945.3	78.3	564.9	3,731.8	2,857.7	1,477.0	703.8	524.2	80.5	5.4
1-M.F.....	1.2 (²)	-----	3	4	3	1	1	1	-----	-----
2-S.G.M.....	251.3	2.0	24.8	79.3	83.0	39.0	12.9	10.7	.9	.1
3-G.M.....	3,147.0	24.8	107.1	1,140.7	976.4	493.9	198.6	119.1	18.9	2.3
4-S.M.....	4,474.5	35.2	230.0	1,715.2	1,274.1	661.6	313.5	235.9	41.7	2.5
5-M.....	1,569.2	12.4	74.4	595.9	400.3	219.1	133.4	128.8	16.8	.5
6-S.L.M.....	330.3	2.6	23.6	137.6	71.9	38.0	31.7	24.8	2.1	-----
7-L.M.....	116.3	.9	11.2	44.0	31.0	15.0	9.9	4.2	.1	-----
8-S.G.O. ¹	55.5	.4	3.5	18.7	10.2	9.7	3.8	.6	-----	-----

See footnotes at end of table (p. 15).

4 STATISTICAL BULLETIN 47, U. S. DEPT. OF AGRICULTURE

TABLE 1.—*Grade and staple length of American upland cotton: Quantities ginned in the United States, by States, crop of 1932—Continued*

UNITED STATES—Continued

Grade	All staple lengths		Shorter than $\frac{7}{8}$ inch ¹	$\frac{7}{8}$ and $\frac{29}{32}$ inch	$1\frac{1}{16}$ and $\frac{3}{4}$ inch	1 and $1\frac{1}{16}$ inches	$1\frac{1}{16}$ and $1\frac{3}{32}$ inches	$1\frac{1}{8}$ and $1\frac{1}{32}$ inches	$1\frac{3}{16}$ and $1\frac{7}{32}$ inches	$1\frac{1}{4}$ inches and longer
	1,000 bales 2,216.9	Percent 17.4	1,000 bales 247.9	1,000 bales 964.0	1,000 bales 679.7	1,000 bales 248.9	1,000 bales 56.0	1,000 bales 17.3	1,000 bales 2.8	1,000 bales 0.3
Spotted.....										
3-G.M.....	193.6	1.5	9.9	65.0	72.1	39.4	6.2	.8	.2	
4-S.M.....	1,054.0	8.3	57.1	462.7	359.8	132.5	30.5	9.6	1.5	.3
5-M.....	673.0	5.3	58.7	323.6	201.3	66.1	16.4	6.0	.9	
6-S.L.M. ¹	217.5	1.7	83.7	90.3	32.7	7.7	2.1	.8	.2	
7-L.M. ¹	78.8	.6	38.5	22.4	13.8	3.2	.8	.1		
Yellow Tinged.....	33.8	.3	14.6	11.6	5.0	1.7	.5	.3	.1	
2-S.G.M.....										
3-G.M.....	2.7	(3)	.6	1.2	.6	.2		.1		
4-S.M.....	10.7	.1	1.5	5.3	2.9	.6	.2	.1	.1	
5-M ¹	9.0	.1	4.2	3.0	.9	.6	.2	.1		
6-S.L.M. ¹	9.7	.1	7.1	1.9	.4	.2	.1			
7-L.M. ¹	1.7	(3)	1.2	.2	.2	.1				
Light Yellow Stained.....	.3	(3)		.2	.1					
3-G.M.....	.1	(3)			.1					
4-S.M. ¹1	(3)			.1					
5-M ¹1	(3)			.1					
Yellow Stained.....	.1	(3)		.1						
3-G.M.....										
4-S.M.....										
5-M ¹1	(3)		.1						
Gray.....	10.1	.1	.4	3.6	3.0	1.8	.7	.5	.1	
3-G.M.....	1.2	(3)		.6	.4	.2				
4-S.M.....	5.9	(3)	.2	1.7	1.9	1.3	.4	.3	.1	
5-M ¹	3.0	(3)	.2	1.3	.7	.3	.3	.2		
No grade ^{1,4}	34.8	.3	4.3	11.7	13.5	4.4	.8	.1		

ALABAMA

All grades.....	2,933.7	100.0	137.3	657.5	108.7	26.5	3.1	0.5	0.1
Extra White.....	.3	(3)		.2	.1				
Above 3-G.M.....									
3-G.M.....									
4-S.M.....									
5-M.....									
6-S.L.M.....									
7-L.M.....	.2	(3)		.1	.1				
Below 7-L.M. ¹1	(3)		.1					
White.....	783.1	83.9	111.5	552.3	92.7	23.3	2.9	.4	
1-M.F.....									
2-S.G.M.....									
3-G.M.....	9.9	1.1	1.9	6.0	1.6	.4			
4-S.M.....	271.9	29.1	30.8	193.3	37.6	0.2	.9	.1	
5-M.....	440.7	47.2	68.0	310.7	48.0	12.1	1.7	.2	
6-S.L.M.....	58.4	6.2	10.4	40.9	5.2	1.5	.3	.1	
7-L.M.....	1.5	.2	.3	1.0	.2				
8-S.G.O. ¹6	.1	.1	.3	.1		.1		
9-G.O. ¹1	(3)		.1					
Spotted.....	149.0	16.0	25.7	103.9	15.8	3.2	.2	.1	.1
3-G.M.....	6.7	.7	1.0	4.7	.6	.3			
4-S.M.....	82.1	8.8	14.6	56.3	9.4	1.6	.1	.1	
5-M.....	56.5	6.1	9.7	40.4	5.2	1.1	.1		
6-S.L.M. ¹	3.1	.3	.4	2.2	.4				
7-L.M.....	.6	.1		.3	.2		.1		

See footnotes at end of table.

TABLE 1.—*Grade and staple length of American upland cotton: Quantities ginned in the United States, by States, crop of 1932—Continued*

ALABAMA—Continued

Grade	All staple lengths		Shorter than $\frac{7}{8}$ inch ¹	$\frac{7}{8}$ and $2\frac{1}{2}$ inch	$1\frac{1}{8}$ and $2\frac{1}{2}$ inch	1 and $1\frac{1}{2}$ inches	$1\frac{1}{8}$ and $1\frac{1}{2}$ inches	$1\frac{1}{8}$ and $1\frac{1}{2}$ inches	$1\frac{1}{8}$ and $1\frac{1}{2}$ inches	$1\frac{1}{4}$ inches and longer
	1,000 bales	Percent	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales
Yellow Tinged.....	0.9	0.1	0.1	0.7	0.1	-----	-----	-----	-----	-----
2-S.G.M.	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
3-G.M.	.1	(3)	-----	-----	.1	-----	-----	-----	-----	-----
4-S.M.	.6	-----	.1	.1	.4	.1	-----	-----	-----	-----
5-M. ¹	.2	(3)	-----	-----	.2	-----	-----	-----	-----	-----
6-S.L.M. ¹	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
7-L.M. ¹	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
Gray.....	.2	(3)	-----	-----	.2	-----	-----	-----	-----	-----
3-G.M.	.1	(3)	-----	-----	.1	-----	-----	-----	-----	-----
4-S.M.	.1	(3)	-----	-----	.1	-----	-----	-----	-----	-----
5-M. ¹	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
No grade ^{1,4}2	(3)	-----	-----	.2	-----	-----	-----	-----	-----

ARIZONA

All grades.....	² 58.8	100.0	0.1	1.4	6.9	23.5	20.3	6.3	0.3	-----
Extra White.....	45.4	77.2	-----	1.0	4.1	16.1	17.7	6.2	.3	-----
Above 3-G.M.	.2	.3	-----	-----	-----	-----	-----	.1	.1	-----
3-G.M.	11.4	19.4	-----	-----	.4	2.5	5.3	3.0	.2	-----
4-S.M.	19.5	33.2	-----	-----	.7	6.6	9.1	3.0	.1	-----
5-M.	9.4	18.0	-----	-----	1.4	4.9	3.0	.1	-----	-----
6-S.L.M.	3.8	6.6	-----	.3	1.3	2.0	.2	-----	-----	-----
7-L.M.	.9	1.5	-----	.6	.2	.1	-----	-----	-----	-----
Below 7-L.M. ¹	.2	.3	-----	.1	.1	-----	-----	-----	-----	-----
White.....	.1	.2	-----	-----	.1	-----	-----	-----	-----	-----
1-M.F.	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
2-S.G.M.	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
3-G.M.	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
4-S.M.	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
5-M.	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
6-S.L.M.	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
7-L.M.	.1	.2	-----	-----	.1	-----	-----	-----	-----	-----
8-S.G.O. ¹	-----	-----	-----	-----	.1	-----	-----	-----	-----	-----
9-G.O. ¹	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
Spotted.....	12.9	21.9	-----	.1	2.7	7.4	2.6	.1	-----	-----
3-G.M.	.3	.5	-----	-----	-----	-----	1	.2	-----	-----
4-S.M.	8.8	15.0	-----	-----	1.0	5.6	2.1	.1	-----	-----
5-M.	3.5	5.9	-----	.1	1.4	1.7	.3	-----	-----	-----
6-S.L.M. ¹	.3	.5	-----	-----	.3	-----	-----	-----	-----	-----
7-L.M. ¹	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
No grade ^{1,4}4	.7	.1	.3	-----	-----	-----	-----	-----	-----

ARKANSAS

All grades.....	² 1,283.4	100.0	25.8	185.9	375.0	360.4	196.8	120.4	17.3	1.8
Extra White.....	2.9	.2	-----	.1	.2	.4	.3	1.6	.3	-----
Above 3-G.M.	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
3-G.M.	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
4-S.M.	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
5-M.	.6	(3)	-----	-----	-----	-----	-----	-----	-----	-----
6-S.L.M.	1.8	.1	-----	-----	.2	.3	.2	.9	.2	-----
7-L.M.	.5	(3)	-----	-----	.1	.1	.1	.2	-----	-----
Below 7-L.M. ¹	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----

See footnotes at end of table.

TABLE 1.—*Grade and staple length of American upland cotton: Quantities ginned in the United States, by States, crop of 1932—Continued*

ARKANSAS—Continued

Grade	All staple lengths		Shorter than $\frac{7}{8}$ inch ¹	$\frac{7}{8}$ and $2\frac{1}{32}$ inch	$1\frac{1}{16}$ and $2\frac{1}{32}$ inch	1 and $1\frac{1}{32}$ inches	$1\frac{1}{16}$ and $1\frac{1}{32}$ inches	$1\frac{1}{8}$ and $1\frac{1}{32}$ inches	$1\frac{1}{16}$ and $1\frac{1}{32}$ inches	$1\frac{1}{4}$ inches and longer
	1,000 bales	Percent	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales
White.....	1,210.0	94.4	23.8	169.8	346.9	344.2	191.6	117.2	10.6	1.8
1-M. F.....										
2-S.G.M.....										
3-G.M.....	12.7	1.0	.8	2.9	5.2	3.1	.6	.1		
4-S.M.....	337.0	26.3	11.2	60.5	109.5	92.6	37.3	17.5	2.7	.6
5-M.....	549.8	42.9	10.5	77.2	153.5	157.8	90.0	51.3	8.6	.9
6-S.L.M.....	212.6	16.6	.9	13.4	48.2	66.4	42.0	36.8	4.6	.3
7-L.M.....	50.1	3.9	.2	3.7	11.0	12.8	13.0	8.8	.6	
8-S.G.O.I.....	20.8	2.3	.2	3.7	11.6	6.1	5.8	2.3	.1	
9-G.O.I.....	18.0	1.4		2.4	6.9	5.4	2.0	.4		
Spotted.....	60.0	4.7	1.8	14.2	24.3	13.5	4.3	1.5	.4	
3-G.M.....	1.9	.2	.1	.6	.7	.4	.1			
4-S.M.....	29.9	2.3	1.3	8.2	12.1	6.2	1.5	.5	.1	
5-M.....	14.5	1.1	.4	2.8	4.7	4.1	1.7	.6	.2	
6-S.L.M.I.....	5.7	.5		.9	2.4	1.4	.6	.3	.1	
7-L.M.I.....	8.0	.6		1.7	4.4	1.4	.4	.1		
Yellow Tinged.....	.9	.1		.2	.3	.3	.1			
2-S.G.M.....										
3-G.M.....										
4-S.M.....	.1	(3)			.1					
5-M.I.....	.2	(3)				.1	.1			
6-S.L.M.I.....	.2	(3)		.1		.1				
7-L.M.I.....	.4	(3)		.1	.2	.1				
Light Yellow Stained.....	.1	(3)				.1				
3-G.M.....	.1	(3)			.1					
4-S.M.I.....										
5-M.I.....										
Gray.....	.5	(3)				.2	.3			
3-G.M.....										
4-S.M.....	.4	(3)			.2	.2				
5-M.I.....	.1	(3)				.1				
No grade ^{1,4}	8.1	.6	.2	1.0	4.0	1.7	.5	.1		

CALIFORNIA

All grades.....	2 124.4	100.0		2.2	3.9	13.4	57.9	46.8	0.2
Extra White.....	119.2	95.8		.4	1.8	12.7	57.5	46.6	.2
Above 3-G.M.....	1.1	.9					.5	.6	
3-G.M.....	71.6	57.5			.1	3.8	34.7	32.9	.1
4-S.M.....	37.9	30.5			.1	5.9	19.3	12.5	.1
5-M.....	5.7	4.6			.7	1.7		.6	
6-S.L.M.....	1.6	1.3		.2	.6	.5	.3		
7-L.M.....	.5	.4		.2		.3			
Below 7-L.M.I.....	.8	.6			.3	.5			
White.....	.2	.2			.1	.1			
1-M. F.....									
2-S.G.M.....									
3-G.M.....									
4-S.M.....									
5-M.....									
6-S.L.M.....									
7-L.M.....	.1	.1			.1				
8-S.G.O.I.....									
9-G.O.I.....							.1		

See footnotes at end of table.

TABLE 1.—*Grade and staple length of American upland cotton: Quantities ginned in the United States, by States, crop of 1932—Continued*

CALIFORNIA—Continued

Grade	All staple lengths		Shorter than $\frac{7}{8}$ inch ¹	$\frac{7}{8}$ and $2\frac{1}{2}$ inch	$1\frac{1}{8}$ and $3\frac{1}{2}$ inch	1 and $1\frac{1}{2}$ inches	$1\frac{1}{8}$ and $1\frac{3}{4}$ inches	$1\frac{1}{8}$ and $1\frac{1}{2}$ inches	$1\frac{1}{8}$ and $1\frac{1}{2}$ inches	$1\frac{1}{4}$ inches and longer
	1,000 bales	Percent								
Spotted.....	2.5	2.0		0.2	01.3	0.6	0.3	0.1		
3-G.M.....	.1	.1					.1			
4-S.M.....	1.3	1.0				.5	.5	.2	.1	
5-M.....	1.0	.8		.2		.7	.1			
6-S.L.M. ¹1	.1				.1				
7-L.M. ¹										
Gray.....	.2	.2						.1	.1	
3-G.M.....										
4-S.M.....	.2	.2						.1	.1	
No grade ^{1,4}	2.3	1.8		1.6		.7				

FLORIDA

All grades.....	2 15.6	100.0	2.3	12.5	0.7	0.1				
White.....	10.3	66.0	1.6	8.3	.4					
1-M.F.....										
2-S.G.M.....										
3-G.M.....										
4-S.M.....	2.8	17.0	.2	2.4	.2					
5-M.....	0.1	39.1	1.0	5.0	.1					
6-S.L.M.....	1.4	9.0	.4	.9	.1					
7-L.M.....										
8-S.G.O. ¹										
9-G.O. ¹										
Spotted.....	5.3	34.0	.7	4.2	.3	.1				
3-G.M.....	.2	1.3	.1		.1					
4-S.M.....	2.7	17.3	.2	2.3	.2					
5-M.....	2.3	14.8	.4	1.8		.1				
6-S.L.M. ¹1	.6		.1						
7-L.M. ¹										

GEORGIA

All grades.....	2 861.8	100.0	43.1	502.8	103.1	53.9	6.7	1.0	0.3	
Extra White.....	1.6	.2	.1	.7	.3	.4	.1			
Above 3-G.M.....										
3-G.M.....										
4-S.M.....										
5-M.....	.5	.1		.2	.1	.1	.1			
6-S.L.M.....	1.0	.1	.1	.4	.2	.3				
7-L.M.....	.1	(8)		.1						
Below 7-L.M. ¹										
White.....	702.9	81.6	36.1	480.7	128.3	44.3	5.5	1.7	.3	
1-M.F.....	.1	(8)		.1						
2-S.G.M.....										
3-G.M.....	15.3	1.8	.7	0.6	3.6	1.3	.1			
4-S.M.....	228.2	26.5	8.1	146.9	52.0	17.8	2.4	.8	.2	
5-M.....	347.4	40.3	14.9	248.5	59.7	20.7	2.7	.8	.1	
6-S.L.M.....	95.7	11.1	8.9	70.9	11.6	4.1	.2	.1		
7-L.M.....	13.6	1.6	2.6	0.1	1.4	.4	.1			
8-S.G.O.J.....	2.1	.2	.7	1.3	.1					
9-G.O.J.....	.5	.1	.2	.3						

See footnotes at end of table.

TABLE 1.—*Grade and staple length of American upland cotton: Quantities ginned in the United States, by States, crop of 1932—Continued*

GEORGIA—Continued

Grade	All staple lengths		Shorter than $\frac{7}{8}$ inch ¹	$\frac{7}{8}$ and $\frac{21}{32}$ inch	$1\frac{1}{16}$ and $\frac{31}{32}$ inch	1 and $1\frac{1}{32}$ inches	$1\frac{1}{16}$ and $1\frac{3}{32}$ inches	$1\frac{1}{8}$ and $1\frac{9}{32}$ inches	$1\frac{3}{16}$ and $1\frac{15}{32}$ inches	$1\frac{1}{4}$ inches and longer
	1,000 bales	Percent								
Spotted	156.0	18.1								
3-G.M.	12.2	1.4		.2	6.9	3.8	1.1		.1	
4-S.M.	77.3	9.0		2.7	40.8	18.5	5.5	.7	.1	
5-M.	57.9	6.7		2.8	41.4	11.0	2.4	.3		
6-S.L.M. ¹	8.1	.9		1.1	5.8	1.0	.2			
7-L.M. ¹	.5	.1		.1	.4					
Yellow Tinged	.8	.1			.0	.2				
2-S.G.M.										
3-G.M.	.1	(3)			.1					
4-S.M.	.6	.1			.6	.1				
5-M. ¹	.1	(3)				.1				
6-S.L.M. ¹										
7-L.M. ¹										
Gray	.4	(3)			.4					
3-G.M.	.1	(3)			.1					
4-S.M.	.2	(3)			.2					
5-M. ¹	.1	(3)			.1					
No grade ^{1,4}	.1	(3)			.1					

LOUISIANA

All grades	2 599.5	100.0	62.1	131.0	152.6	156.4	88.7	28.2	2.4	0.1
Extra White	15.1	2.5	.4	2.4	4.4	4.1	2.8	.9	.1	
Above 3-G.M.										
3-G.M.	.1	(3)					.1			
4-S.M.	5.8	1.0	.2	1.2	1.5	1.3	1.1	.4	.1	
5-M.	6.1	1.0	.1	.7	1.0	1.7	1.3	.4		
6-S.L.M.	2.8	.5	.1	.5	.9	.8	.4	.1		
7-L.M.	.3	(3)				.1	.2			
Below 7-L.M. ¹										
White	548.7	91.5	58.9	121.7	139.9	140.4	81.4	24.1	2.2	.1
1-M.F.										
2-S.G.M.	.4	.1	.2	.1	.1					
3-G.M.	31.4	5.2	8.9	9.3	6.0	5.5	1.3	.3	.1	
4-S.M.	225.7	37.6	35.6	57.7	50.5	52.3	20.6	8.0	.9	.1
5-M.	221.6	37.0	13.3	45.8	63.5	60.4	25.8	11.8	1.0	
6-S.L.M.	58.5	9.8	.9	7.9	17.6	17.3	11.1	3.5	.2	
7-L.M.	10.3	1.7		.8	1.9	4.6	2.5	.5		
8-S.G.O. ¹	.7	.1		.1	.2	.3	.1			
9-G.O. ¹	.1	(3)			.1					
Spotted	33.7	5.6	2.1	5.9	8.2	11.7	4.5	1.2	.1	
3-G.M.	7.8	1.3	.9	1.4	1.7	2.9	.6	.1		
4-S.M.	18.7	3.1	1.0	3.6	5.1	6.0	2.0	.9	.1	
5-M.	6.7	1.1	.2	.8	1.3	2.6	1.6	.2		
6-S.L.M. ¹	.6	.1		.1	.1	.1	.3			
7-L.M. ¹	.1	(3)				.1				
Yellow Tinged	.9	.2		.9						
2-S.G.M.										
3-G.M.										
4-S.M.	.9	.2		.9						
5-M. ¹										
6-S.L.M. ¹										
7-L.M. ¹										
Gray	.4	.1		.1	.1	.2				
3-G.M.	.1	(3)			.1					
4-S.M.	.2	(3)			.1		.1			
5-M. ¹	.1	(3)					.1			
No grade ^{1,4}	.7	.1	.7							

See footnotes at end of table.

TABLE 1.—*Grade and staple length of American upland cotton: Quantities ginned in the United States, by States, crop of 1932—Continued*

MISSISSIPPI

Grade	All staple lengths		Shorter than $\frac{7}{16}$ inch ¹	$\frac{7}{16}$ and $\frac{29}{32}$ inch	$1\frac{1}{16}$ and $2\frac{1}{2}$ inch	1 and $1\frac{1}{32}$ inches	$1\frac{1}{8}$ and $1\frac{1}{32}$ inches	$1\frac{1}{4}$ and $1\frac{5}{32}$ inches	$1\frac{1}{8}$ and $1\frac{7}{32}$ inches	$1\frac{1}{4}$ inches and longer
	1,000 bales	Percent	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales
All grades.....	2 1,161.2	100.0	32.2	148.6	162.3	170.9	271.5	328.2	45.3	2.2
Extra White.....		.3	(3)	-----	-----	-----	-----	.1	.2	-----
Above 3-G.M.....										
3-G.M.....										
4-S.M.....										
5-M.....										
6-S.L.M.....	.2	(3)						.1	.1	
7-L.M.....	.1	(3)						.1	.1	
Below 7-L.M. ¹										
White.....	1,021.9	88.1	25.2	111.7	124.7	142.0	254.7	317.7	43.8	2.1
1-M.F.....										
2-S.G.M.....	.1	(3)								
3-G.M.....	31.0	2.7	3.0	4.1	2.3	3.3	8.0	9.5	.7	.1
4-S.M.....	348.2	30.0	13.6	55.2	50.9	56.3	84.1	75.9	11.1	1.1
5-M.....	434.9	37.5	7.4	44.1	57.1	60.6	103.1	140.0	21.9	.7
6-S.L.M.....	105.8	14.3	1.1	7.1	12.4	15.9	43.5	76.7	8.9	.2
7-L.M.....	31.8	2.7	.1	.9	1.6	2.4	12.0	13.7	1.1	
8-S.G.O. ¹	7.6	.7		.2	.2	1.9	3.5	1.7	.1	
9-G.O. ¹	2.5	.2		.1	.2	1.6	.5	.1		
Spotted.....	137.6	11.9	6.8	36.7	37.3	28.6	18.6	10.0	1.5	.1
3-G.M.....	6.8	.6	.5	1.9	2.0	1.2	.7	.4	.1	
4-S.M.....	86.1	7.4	4.4	25.1	23.0	17.4	9.4	5.7	1.0	.1
5-M.....	40.3	3.5	1.8	8.8	10.8	9.1	5.9	3.5	.4	
6-S.L.M. ¹	4.1	.4	.1	.8	1.4	.8	.6	.4		
7-L.M. ¹3	(3)		.1	.1	.1				
Yellow Tinged.....	.5	(3)		.1	.1	.2	.1			
2-S.G.M.....										
3-G.M.....	.1	(3)								
4-S.M.....	.3	(3)								
5-M. ¹1	(3)								
6-S.L.M. ¹										
7-L.M. ¹										
Light Yellow Stained.....	.1	(3)			.1					
3-G.M.....										
4-S.M. ¹										
5-M. ¹1	(3)			.1					
Gray.....	.3	(3)						.3		
3-G.M.....										
4-S.M.....	.2	(3)						.2		
5-M. ¹1	(3)						.1		
No grade ¹5	(3)	.2		.2	.1				

MISSOURI

All grades.....	2 300.7	100.0	2.0	52.1	114.9	107.7	23.0	1.0		
Extra White.....	4.1	1.4		.3	1.1	1.6	1.1			
Above 3-G.M.....										
3-G.M.....										
4-S.M.....										
5-M.....										
6-S.L.M.....	.1	(3)			.1					
7-L.M.....	3.7	1.3		.2	1.0	1.5	1.0			
Below 7-L.M.....	.3	.1			.1	.1	.1			

See footnotes at end of table.

10 STATISTICAL BULLETIN 47, U. S. DEPT. OF AGRICULTURE

TABLE 1.—*Grade and staple length of American upland cotton: Quantities ginned in the United States, by States, crop of 1932—Continued*

MISSOURI—Continued

Grade	All staple lengths		Shorter than $\frac{7}{8}$ inch ¹	$\frac{7}{8}$ and $\frac{25}{32}$ inch	$1\frac{1}{16}$ and $8\frac{1}{32}$ inch	1 and $1\frac{1}{32}$ inches	$1\frac{1}{16}$ and $1\frac{3}{32}$ inches	$1\frac{1}{8}$ and $1\frac{1}{32}$ inches	$1\frac{1}{16}$ and $1\frac{1}{32}$ inches	1/4 inches and longer
	1,000 bales 279.5	Percent 92.9								
White.....										
1-M.F.										
2-S.G.M.										
3-G.M.	.4	.1			.1	.1		.2		
4-S.M.	28.0	9.3	.3	8.0	10.8	7.8	1.0	.1		
5-M.	112.2	37.3	1.1	16.8	38.4	44.8	10.7	.4		
6-S.L.M.	77.2	25.7	.2	4.7	27.6	35.7	8.5	.5		
7-L.M.	19.8	6.6	.2	2.1	8.7	7.8	1.0			
8-S.G.O. ¹	21.8	7.2		6.2	10.8	4.6	.2			
9-G.O. ¹	20.1	6.7		8.6	9.3	1.9	.3			
Spotted.....										
3-G.M.	.2	.1	.1	.1						
4-S.M.	1.4	.4	.1	.4	.6	.3				
5-M.	1.8	.6	.2	.2	.6	.9	.1			
6-S.L.M. ¹	.3	.1		.1	.1	.1				
7-L.M. ¹	4.1	1.4		1.7	2.3	.1				
No grade ^{1,4}	9.3	3.1		2.8	4.5	1.9	.1			

NEW MEXICO

All grades.....	2 67.5	100.0	2.1	2.6	3.4	23.6	25.1	10.7		
Extra White.....	38.9	57.6				.6	8.4	19.3	10.6	
Above 3-G.M.	.2	.2							.2	
3-G.M.	8.7	13.0					.7	3.8	4.2	
4-S.M.	23.5	34.9					4.5	13.0	6.0	
5-M.	5.5	8.2				.1	2.8	2.4	.2	
6-S.L.M.	.2	.2					.1	.1		
7-L.M.	.2	.2				.1	.1			
Below 7-L.M. ¹	.6	.9				.4	.2			
White.....	1.1	1.8						.5	.6	
1-M.F.										
2-S.G.M.										
3-G.M.										
4-S.M.	.6	.9						.2	.4	
5-M.	.5	.7						.3	.2	
6-S.L.M.										
7-L.M.										
8-S.G.O. ¹										
9-G.O. ¹										
Spotted.....	21.0	31.2	*		1.2	14.5	5.2	.1		
3-G.M.	4.9	7.3					2.8	2.0	.1	
4-S.M.	13.3	19.8					9.8	3.1		
5-M.	2.3	3.4					.6	1.6	.1	
6-S.L.M. ¹	.5	.7					.2	.3		
7-L.M. ¹										
Yellow Tinged.....	.2	.2						.2		
2-S.G.M.										
3-G.M.	.1	.1						.1		
4-S.M.	.1	.1						.1		
5-M. ¹										
6-S.L.M. ¹										
7-L.M. ¹										
No grade ^{1,4}	6.3	9.4	2.1	2.6	1.6					

See footnotes at end of table.

TABLE 1.—*Grade and staple length of American upland cotton: Quantities ginned in the United States, by States, crop of 1932—Continued*

NORTH CAROLINA

Grade	All staple lengths	Shorter than $\frac{7}{8}$ inch ¹	$\frac{7}{8}$ and $\frac{2\frac{1}{2}}{2\frac{1}{2}}$ inch	$1\frac{1}{16}$ and $2\frac{1}{2}$ inch	1 and $1\frac{1}{2}$ inches	$1\frac{1}{16}$ and $1\frac{3}{4}$ inches	$1\frac{1}{8}$ and $1\frac{5}{8}$ inches	$1\frac{3}{16}$ and $1\frac{7}{8}$ inches	$1\frac{1}{4}$ inches and longer	
All grades	<i>1,000 bales</i> 2 680.3	<i>Percent</i> 100.0	<i>1,000 bales</i> 7.8	<i>1,000 bales</i> 227.8	<i>1,000 bales</i> 223.6	<i>1,000 bales</i> 160.4	<i>1,000 bales</i> 47.5	<i>1,000 bales</i> 11.0	<i>1,000 bales</i> 2.1	
Extra White	2.5	.4	—	.6	.6	.7	.5	.1	—	
Above 3-G.M.	—	—	—	—	—	—	—	—	—	
3-G.M.	—	—	—	—	—	—	—	—	—	
4-S.M.	.1	(²)	—	—	.1	—	—	—	—	
5-M.	.8	—	—	—	.2	.2	.1	—	—	
6-S.L.M.	1.4	.1	—	.3	.2	.4	.4	.1	—	
7-L.M.	.2	(²)	—	—	.1	.1	—	—	—	
Below 7-L.M.	—	—	—	—	—	—	—	—	—	
White	618.9	90.9	7.3	200.6	203.4	150.4	44.4	10.6	2.1	.1
1-M.F.	—	—	—	—	—	—	—	—	—	
2-S.G.M.	—	—	—	—	—	—	—	—	—	
3-G.M.	13.8	2.0	.7	4.2	4.4	3.8	.6	.1	—	
4-S.M.	216.3	31.8	3.9	68.6	73.0	55.3	12.2	2.5	.7	.1
5-M.	294.0	43.2	2.3	95.6	94.8	71.4	23.3	5.6	1.0	—
6-S.L.M.	85.9	12.6	.3	29.2	28.4	18.0	7.6	2.1	.3	—
7-L.M.	8.3	1.2	.1	2.7	2.7	1.8	.6	.3	.1	—
8-S.G.O. ¹	.6	.1	—	.3	.1	.1	.1	—	—	—
9-G.O. ¹	—	—	—	—	—	—	—	—	—	—
Spotted	52.8	7.8	.4	24.5	17.4	8.1	2.2	.2	—	—
3-G.M.	3.7	.5	.1	1.5	1.3	.7	.1	—	—	—
4-S.M.	26.3	3.9	.2	11.1	9.1	4.4	1.4	.1	—	—
5-M.	20.8	3.1	.1	10.5	6.5	2.9	.7	.1	—	—
6-S.L.M. ¹	2.0	.3	—	1.4	.5	.1	—	—	—	—
7-L.M. ¹	—	—	—	—	—	—	—	—	—	—
Yellow Tinged	.4	.1	—	.3	.1	—	—	—	—	—
2-S.G.M.	—	—	—	—	—	—	—	—	—	—
3-G.M.	.1	(²)	—	—	.1	—	—	—	—	—
4-S.M.	.1	(²)	—	—	.1	—	—	—	—	—
5-M. ¹	.1	(²)	—	—	.1	—	—	—	—	—
6-S.L.M. ¹	.1	(²)	—	—	.1	—	—	—	—	—
7-L.M. ¹	—	—	—	—	—	—	—	—	—	—
Gray	5.6	.8	—	1.8	2.1	1.2	.4	.1	—	—
3-G.M.	.7	.1	—	.4	.2	.1	—	—	—	—
4-S.M.	3.5	.5	—	.9	1.3	.9	.3	.1	—	—
5-M. ¹	1.4	.2	—	.5	.6	.2	.1	—	—	—
No grade ^{1,4}	.1	(²)	.1	—	—	—	—	—	—	—

OKLAHOMA

All grades	<i>2 1,072.0</i>	<i>100.0</i>	<i>91.9</i>	<i>535.8</i>	<i>386.1</i>	<i>52.4</i>	<i>5.2</i>	<i>0.5</i>	<i>0.1</i>
Extra White	34.8	3.3	.5	7.2	19.8	6.7	.6	—	—
Above 3-G.M.	—	—	—	—	—	—	—	—	—
3-G.M.	1.6	.2	.1	.3	.8	.4	—	—	—
4-S.M.	9.0	.8	.2	1.7	5.1	1.7	.3	—	—
5-M.	13.5	1.3	.1	2.2	7.8	3.1	.3	—	—
6-S.L.M.	7.9	.7	.1	1.8	4.6	1.4	—	—	—
7-L.M.	2.8	.3	—	1.2	1.5	.1	—	—	—
Below 7-L.M. ¹	—	—	—	—	—	—	—	—	—
White	823.7	76.8	63.3	425.1	292.6	38.0	4.2	.4	.1
1-M.F.	—	—	—	—	—	—	—	—	—
2-S.G.M.	.1	(²)	—	—	.1	—	—	—	—
3-G.M.	24.5	2.3	1.4	8.3	11.7	2.9	.2	—	—
4-S.M.	286.9	20.8	21.6	133.7	113.1	10.5	1.8	.2	—
5-M.	328.2	30.4	22.8	171.4	116.4	13.6	1.7	.2	.1
6-S.L.M.	142.3	13.3	10.5	83.6	43.2	4.5	.5	—	—
7-L.M.	37.6	3.5	4.9	24.8	7.4	.5	—	—	—
8-S.G.O. ¹	5.8	.5	1.9	3.2	.7	—	—	—	—
9-G.O. ¹	.3	(²)	.2	.1	—	—	—	—	—

See footnotes at end of table.

12 STATISTICAL BULLETIN 47, U.S. DEPT. OF AGRICULTURE

TABLE 1.—*Grade and staple length of American upland cotton: Quantities ginned in the United States, by States, crop of 1932—Continued*

OKLAHOMA—Continued

Grade	All staple lengths		Shorter than $\frac{7}{8}$ inch ¹	$\frac{7}{8}$ and $\frac{29}{32}$ inch	$1\frac{1}{16}$ and $\frac{31}{32}$ inch	1 and $1\frac{1}{32}$ inches	$1\frac{1}{16}$ and $1\frac{3}{32}$ inches	$1\frac{1}{8}$ and $1\frac{1}{32}$ inches	$1\frac{1}{16}$ and $1\frac{1}{2}$ inches	$1\frac{1}{4}$ inches and longer
	1,000 bales	Percent								
Spotted.....	210.8	19.7	27.6	102.3	72.7	7.7	0.4	0.1		
3-G.M.....	16.5	1.5	.9	6.9	7.2	1.5				
4-S.M.....	102.4	9.6	4.5	50.6	42.4	4.5	.3	.1		
5-M.....	59.1	5.5	5.3	32.9	19.2	1.6	.1			
6-S.L.M. ¹	22.6	2.1	8.9	10.0	3.6	.1				
7-L.M. ¹	10.2	1.0	8.0	1.9	.3					
Yellow Tinged.....	2.4	.2	.4	1.0	1.0					
2-S.G.M.....										
3-G.M.....	.4	(3)		.1	.3					
4-S.M.....	1.2	.1		.6	.6					
5-M. ¹4	(3)	.1	.2	.1					
6-S.L.M. ¹4	(3)	.3	.1						
7-L.M. ¹										
Light Yellow Stained.....	.1	(3)		.1						
3-G.M.....										
4-S.M. ¹1	(3)		.1						
5-M. ¹										
Gray.....	.1	(3)		.1						
3-G.M.....										
4-S.M.....										
5-M. ¹1	(3)		.1						
No grade ¹1	(3)	.1							

SOUTH CAROLINA

All grades.....	2 722.2	100.0	2.8	181.2	225.6	169.5	81.8	44.0	15.8	1.5
Extra White.....	2.3	.3		.2	.6	.7	.4	.3	.1	
Above 3-G.M.....										
3-G.M.....										
4-S.M.....	.1	(3)			.1					
5-M.....	.4	.1		.1	.2	.1				
6-S.L.M.....	1.7	.2		.1	.3	.6	.3	.3	.1	
7-L.M.....	.1	(3)					.1			
Below 7-L.M. ¹										
White.....	622.9	86.2	2.3	148.4	180.4	149.0	75.7	41.8	15.0	1.3
1-M.F.....										
2-S.G.M.....										
3-G.M.....	10.3	1.4	.2	3.0	2.7	2.2	1.5	.6	.1	
4-S.M.....	210.6	29.2	1.3	57.2	63.7	49.1	23.9	11.9	3.1	.4
5-M.....	327.0	45.2	.7	75.4	102.1	80.4	37.3	21.4	8.0	.8
6-S.L.M.....	69.8	9.7	.1	12.3	19.8	16.0	11.9	7.0	2.6	.1
7-L.M.....	5.0	.7		.5	1.1	1.3	1.0	.8	.3	
8-S.G.O. ¹2	(3)					.1	.1		
9-G.O. ¹										
Spotted.....	95.0	13.2	.5	32.2	35.1	19.3	5.4	1.6	.7	.2
3-G.M.....	3.3	.5	.1	1.1	1.1	.7	.2	.1		
4-S.M.....	48.2	6.7	.2	16.0	18.1	10.0	2.3	.9	.5	.2
5-M.....	39.8	5.5	.2	13.5	14.5	8.2	2.6	.6	.2	
6-S.L.M. ¹	3.5	.5		1.5	1.3	.4	.3			
7-L.M. ¹2	(3)		.1	.1					
Yellow Tinged.....	1.4	.2		.3	.3	.4	.2	.2		
2-S.G.M.....										
3-G.M.....	.1	(3)				.1				
4-S.M.....	.7	.1		.1	.2	.2	.1	.1		
5-M. ¹6	.1		.2	.1	.1	.1	.1		
6-S.L.M. ¹										
7-L.M. ¹										

See footnotes at end of table.

TABLE 1.—*Grade and staple length of American upland cotton: Quantities ginned in the United States, by States, crop of 1932—Continued*

SOUTH CAROLINA—Continued

Grade	All staple lengths		Shorter than $\frac{7}{8}$ inch ¹	$\frac{7}{8}$ and $\frac{29}{32}$ inch	$1\frac{1}{16}$ and $\frac{3}{4}\frac{1}{2}$ inch	1 and $1\frac{1}{2}$ inches	$1\frac{1}{16}$ and $1\frac{1}{2}$ inches	$1\frac{1}{8}$ and $1\frac{1}{2}$ inches	$1\frac{1}{16}$ and $1\frac{1}{2}$ inches	1 $\frac{1}{4}$ inches and longer
	1,000 bales	Percent								
Gray.....	0.6	0.1			0.1					
3-G.M.....	.1	(3)					.1			
4-S.M.....	.4		.1			.1			.1	
5-M. ¹1	(3)				.1				

TENNESSEE

All grades.....	2 467.5	100.0	12.4	115.9	175.3	134.7	24.6	4.3	0.3	
Extra White.....	.6	.1				.2	.4			
Above 3-G.M.....										
3-G.M.....										
4-S.M.....										
5-M.....										
6-S.L.M.....	.3	(3)				.1		.2		
7-L.M. ¹3	(3)				.1		.2		
Below 7-L.M. ¹										
White.....	394.2	84.3	10.0	89.7	144.4	123.6	22.4	3.8	.3	
1-M.F.....										
2-S.G.M.....										
3-G.M.....	2.0	.4	.1	.6	.7	.4		.2		
4-S.M.....	117.9	25.2	2.8	24.9	42.6	38.4	7.7	1.4	.1	
5-M.....	195.9	41.9	5.2	42.1	70.2	65.9	10.8	1.6	.1	
6-S.L.M.....	50.8	10.9	1.6	12.5	18.3	14.7	3.1	.5	.1	
7-L.M.....	13.8	3.0	.2	4.0	6.5	2.4	.4	.3		
8-S.G.O. ¹	8.6	1.8	.1	2.9	4.1	1.4	.1			
9-G.O. ¹	5.2	1.1		2.7	2.0	.4		.1		
Spotted.....	69.1	14.8	2.3	24.7	29.0	10.4	2.2	.5		
3-G.M.....	1.4	.3	.1	.5	.6	.2				
4-S.M.....	37.1	8.0	1.5	13.7	14.2	6.3	1.1	.3		
5-M.....	15.9	3.4	.6	6.1	6.3	2.3	.5	.1		
6-S.L.M. ¹	7.0	1.5	.1	1.8	4.2	.6	.2	.1		
7-L.M. ¹	7.7	1.6		2.6	3.7	1.0	.4	.4		
Yellow Tinged.....	.5	.1		.2	.2	.1				
2-S.G.M.....										
3-G.M.....										
4-S.M.....	.1	(3)					.1			
5-M. ¹2	(3)			.1		.1			
6-S.L.M. ¹1	(3)					.1			
7-L.M. ¹1	(3)					.1			
Gray.....	.1	(3)				.1				
3-G.M.....										
4-S.M.....										
5-M. ¹1	(3)				.1				
No grade ¹	3.0	.7	.1	1.3	1.4	.2				

TEXAS

All grades.....	2 4,307.4	100.0	414.3	1,918.6	1,554.8	362.3	37.5	19.7	0.2	
Extra White.....	191.9	4.5	4.5	50.5	78.3	35.8	9.6	13.2		
Above 3-G.M.....	.3	(3)						.1	.2	
3-G.M.....	13.5	.3	.4	1.8	2.5	.5	1.9	.6	.4	
4-S.M.....	36.9	.9	.6	6.8	13.1	5.8	4.2	6.4		
5-M.....	45.6	1.1	1.4	12.2	21.1	9.1	1.7	.1		
6-S.L.M.....	73.1	1.7	1.3	20.5	33.2	16.9	1.2			
7-L.M.....	22.4	.5	.8	9.2	8.4	3.4	.5	.1		
Below 7-L.M. ¹1	(3)				.1				

See footnotes at end of table.

14 STATISTICAL BULLETIN 47, U. S. DEPT. OF AGRICULTURE

TABLE 1.—*Grade and staple length of American upland cotton: Quantities ginned in the United States, by States, crop of 1932—Continued*

TEXAS—Continued

Grade	All staple lengths		Shorter than $\frac{7}{8}$ inch ¹	$\frac{7}{8}$ and $2\frac{1}{2}$ inch	$1\frac{1}{8}$ and $3\frac{1}{2}$ inch	1 and $1\frac{1}{2}$ inches	$1\frac{1}{8}$ and $1\frac{1}{2}$ inches	$1\frac{1}{4}$ and $1\frac{1}{2}$ inches	$1\frac{1}{8}$ and $1\frac{1}{2}$ inches	$1\frac{1}{4}$ inches and longer
	1,000 bales	Percent	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales
White.....	2,887.4	67.0	221.8	1,352.9	1,077.9	212.9	17.0	4.7	0.2
1-M.F.	.5	(3)	.1	.1	.2	.1
2-S.G.M.	100.0	2.3	7.3	31.1	45.1	16.0	.5
3-G.M.	865.4	20.1	67.7	322.6	371.2	97.4	5.9	.6
4-S.M.	1,196.4	27.7	82.2	573.1	462.7	70.4	5.5	2.4	.1
5-M.	541.6	12.6	38.4	308.1	165.3	24.1	4.1	1.5	.1
6-S.L.M.	137.8	3.2	15.0	87.9	29.4	4.4	.9	.2
7-L.M.	38.1	.9	8.1	25.7	3.7	.5
8-S.G.O. ¹	7.6	.2	3.0	4.3	.3
9-G.O. ¹
Spotted.....	1,198.8	27.8	172.6	505.8	395.1	112.7	10.8	1.8
3-G.M.	127.7	3.0	6.0	39.2	52.8	27.5	2.0	.2
4-S.M.	517.2	11.9	28.2	225.0	204.9	54.2	6.2	.7
5-M.	348.0	8.1	37.0	162.5	117.8	27.4	2.4	.9
6-S.L.M. ¹	160.2	3.7	73.1	65.5	17.0	3.4	.2
7-L.M. ¹	46.7	1.1	30.3	13.6	2.6	.2
Yellow Tinged.....	24.9	.6	14.2	7.3	2.7	.6	.1
2-S.G.M.
3-G.M.	1.7	(3)	.6	.7	.3	.1
4-S.M.	6.0	.1	1.4	2.7	1.8	.1
5-M. ¹	7.1	.2	4.1	2.2	.4	.3	.1
6-S.L.M. ¹	8.9	.2	6.9	1.7	.2	.1
7-L.M. ¹	1.2	(3)	1.2
Yellow Stained.....	1	(3)	1
3-G.M.
4-S.M. ¹
5-M. ¹	.1	(3)1
Gray.....	1.7	(3)	.3	1.0	.3	.1
3-G.M.	.1	(3)1
4-S.M.	.9	(3)	.1	.5	.2	.1
5-M. ¹	.7	(3)	.2	.5
No grade ^{1,4}	2.6	.1	.9	1.1	.5	.1

VIRGINIA

All grades.....	2	31.3	100.0	1.5	19.4	8.7	1.5	0.2
White.....	28.4	90.7	1.2	17.2	8.3	1.5	.2
1-M.F.
2-S.G.M.
3-G.M.
4-S.M.	4.5	14.4	.1	3.5	.5	.3	.1
5-M.	16.8	53.7	.5	9.3	5.9	1.0	.1
6-S.L.M.	7.0	22.3	.6	4.3	1.9	.2
7-L.M.	.1	.31
8-S.G.O. ¹
9-G.O. ¹
Spotted.....	2.9	0.8	.3	2.2	.4
3-G.M.	.7	2.36	.1
4-S.M.	2.1	0.7	.3	1.5	.3
5-M.	.1	.31
6-S.L.M. ¹
7-L.M. ¹

See footnotes at end of table.

TABLE 1.—*Grade and staple length of American upland cotton: Quantities ginned in the United States, by States, crop of 1932—Continued*

ALL OTHER STATES*

Grade	All staple lengths		Shorter than $\frac{7}{8}$ inch ¹	$\frac{7}{8}$ and $2\frac{1}{2}$ inch	$1\frac{1}{8}$ and $3\frac{1}{2}$ inch	1 and $1\frac{1}{2}$ inches	$1\frac{1}{8}$ and longer			
	1,000 bales ²	Percent	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales
All grades.....	2,140	100.0								
Extra White.....	.1	.7					.1			
Above 3-G.M.....										
3-G.M.....										
4-S.M.....										
5-M.....										
6-S.L.M.....	.1	.7					.1			
7-L.M.....										
Below 7-L.M. ³										
White.....	11.1	79.3		.8	4.0	4.0	1.7	.5	.1	
1-M.F.....										
2-S.G.M.....										
3-G.M.....										
4-S.M.....	2.1	15.0		.4	.8	.7	.2			
5-M.....	5.0	35.7		.2	1.6	2.1	.9	.2		
6-S.L.M.....	2.2	15.7		.1	.7	.7	.4	.2	.1	
7-L.M.....	.4	2.9			.1	.1	.1	.1		
8-S.G.O. ⁴4	2.9			.3	.1				
9-G.O. ⁴	1.0	7.1		.1	.6	.3	.1			
Spotted.....	1.7	12.1		.2	.9	.5	.1			
3-G.M.....	.1	.7			.1					
4-S.M.....	.5	3.6			.2	.2	.1			
5-M.....	.5	3.6		.1	.3	.1				
6-S.L.M. ⁵2	1.3			.1	.1				
7-L.M. ⁵4	2.9		.1	.2	.1				
No grade ⁶	1.1	7.9		.2	.5	.2	.1	.1		

* Untenderable in settlement of futures contracts made subject to sec. 5 of the United States Cotton Futures Act and the regulations of the Secretary of Agriculture thereunder.

¹ As reported by the Bureau of the Census.

² Less than 0.05 percent.

³ Includes bales not otherwise classified above.

⁴ Includes Illinois, Kansas, and Kentucky.

TABLE 2.—*Grade and staple length of American-Egyptian cotton: Quantities ginned in the United States, crops of 1932, 1931, 1930, 1929, and 1928*

[Quantities are given in running bales]

Crop and grade	All staple lengths		Shorter than $1\frac{1}{2}$ inches	$1\frac{1}{2}$ and $1\frac{1}{2}\frac{1}{2}$ inches	$1\frac{1}{8}$ and $1\frac{1}{2}\frac{1}{2}$ inches	$1\frac{1}{8}$ to $1\frac{1}{2}\frac{1}{2}$ inches	$1\frac{1}{8}$ inches and longer
1932	1,000 bales ¹	Percent	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales
All grades.....	18.3	100.0					
1 and $1\frac{1}{2}$	2.0	24.1			.3	1.0	.7
2 and $2\frac{1}{2}$	3.7	44.6		.5	1.6	1.3	.3
3 and $3\frac{1}{2}$	2.2	20.5		.9	.9	.3	.1
4 and $4\frac{1}{2}$4	4.8		.3	.1		
5.....							
Below 5.....							
All grades.....	13.7	100.0			2.4	8.4	2.9
1 and $1\frac{1}{2}$	1.2	8.7			.3	.6	.3
2 and $2\frac{1}{2}$	5.9	43.1			1.1	3.5	1.3
3 and $3\frac{1}{2}$	4.5	32.8			.7	3.0	.8
4 and $4\frac{1}{2}$	1.0	13.9			.3	1.2	.4
5.....	.2	1.5				.1	.1
Below 5.....							

¹ As reported by the Bureau of the Census.

TABLE 2.—*Grade and staple length of American-Egyptian cotton: Quantities ginned in the United States, crops of 1932, 1931, 1930, 1929, and 1928—Continued*

[Quantities are given in running bales]

Crop and grade	All staple lengths		Shorter than 1½ inches	1½ and 1½½ inches	1½ and 1½½ inches	1½ and 1½½ inches	1½ and 1½½ inches	1½ and 1½½ inches
	1,000 bales 1 23.3	Percent 100.0	1,000 bales	1,000 bales 2.5	1,000 bales 10.2	1,000 bales 4.6	1,000 bales	1,000 bales
1930								
All grades								
1 and 1½	6.2	26.6		.4	4.1	1.7		
2 and 2½	11.4	48.9		.7	8.1	2.6		
3 and 3½	5.1	21.9		1.1	3.7	.3		
4 and 4½	.6	2.6		.3	.3			
5								
Below 5								
1929								
All grades	1 28.8	100.0		5.3	17.1	6.0		0.4
1 and 1½	5.5	19.1		.6	3.1	1.7		.1
2 and 2½	16.4	56.9		2.3	10.0	3.8		.3
3 and 3½	6.5	22.6		2.1	3.9	.5		
4 and 4½	.4	1.4		.3	.1			
5								
Below 5								
1928								
All grades	1 28.3	100.0	0.7	13.4	12.5	1.6		.1
1 and 1½	5.5	19.4	.2	2.2	2.7	.4		
2 and 2½	13.6	48.1	.3	5.9	6.5	.8		.1
3 and 3½	8.4	29.7	.2	4.8	3.0	.4		
4 and 4½	.7	2.5		.4	.3			
5	.1	.3		.1				
Below 5								

¹ As reported by the Bureau of the Census.TABLE 3.—*Staple length of American upland cotton: Quantities ginned in the United States, by States, crops of 1932, 1931, 1930, 1929, and 1928*

[Quantities are given in running bales, except that round bales are counted as half bales. Linters are not included]

State	Crop year	All staple lengths ¹		Shorter than $\frac{7}{8}$ inch ²		$\frac{7}{8}$ and $\frac{29}{32}$ inch		$\frac{15}{16}$ and $\frac{31}{32}$ inch	
		1,000 bales	Percent	1,000 bales	Percent	1,000 bales	Percent	1,000 bales	Percent
United States	1932	12,701.3	100	837.7	6.6	4,786.5	37.7	3,671.0	28.9
	1931	16,615.2	100	1,019.5	6.1	6,593.3	39.7	4,511.9	27.2
	1930	13,732.2	100	1,829.2	13.3	5,327.7	38.8	3,421.6	24.9
	1929	14,519.0	100	2,921.5	20.1	5,533.7	38.1	2,748.2	18.9
	1928	14,268.2	100	2,072.1	14.5	5,914.8	41.5	3,225.7	22.6
	1932	933.7	100	137.3	14.7	657.5	70.4	108.7	11.7
Alabama	1931	1,385.0	100	136.5	9.9	1,044.7	75.4	108.7	12.2
	1930	1,444.9	100	553.3	38.3	802.2	55.5	77.1	5.3
	1929	1,307.7	100	585.6	44.8	689.0	52.7	28.3	2.2
	1928	1,096.6	100	264.2	24.1	762.6	69.5	49.8	4.5
	1932	58.8	100	.1	.2	1.4	2.4	6.9	11.7
Arizona	1931	97.3	100	3.3	3.4	.6	.6	14.9	15.3
	1930	127.2	100	.3	.2	5.5	4.3	41.2	32.4
	1929	120.7	100	1.5	1.2	4.3	3.6	27.4	22.7
	1928	117.4	100	2.6	2.2	11.3	9.6	17.6	15.0
Arkansas	1932	1,283.4	100	25.8	2.0	185.9	14.5	375.0	29.2
	1931	1,836.1	100	20.9	1.1	307.4	16.7	549.6	29.9
	1930	863.5	100	76.9	8.9	196.1	22.7	252.9	20.3
	1929	1,395.9	100	177.9	12.7	393.4	28.2	370.1	27.2
	1928	1,216.2	100	127.0	10.4	350.4	29.6	200.5	23.9
California	1932	124.4	100			2.2	1.8	3.0	3.1
	1931	171.2	100	8.9	5.2	.1	.1	2.4	1.4
	1930	256.3	100			1.7	.7	13.9	5.4
	1929	254.1	100	.8	.3	5.0	2.0	17.1	6.7
	1928	171.0	100	2.8	1.6	20.5	12.0	29.3	17.1

¹ As reported by the Bureau of the Census.² Untenderable in settlement of futures contracts made subject to sec. 5 of the United States Cotton Futures Act and the regulations of the Secretary of Agriculture thereunder.

TABLE 3.—*Staple length of American upland cotton: Quantities ginned in the United States, by States, crops of 1932, 1931, 1930, 1929, and 1928—Continued*

State	Crop year	All staple lengths ¹		Shorter than $\frac{3}{8}$ inch ²		$\frac{1}{2}$ and $2\frac{1}{2}$ inch		$1\frac{1}{2}$ and $3\frac{1}{2}$ inch	
		1,000 bales	Percent	1,000 bales	Percent	1,000 bales	Percent	1,000 bales	Percent
Florida	1932	15.6	100	2.3	14.8	12.5	80.1	0.7	4.5
	1931	43.4	100	2.9	6.7	33.9	78.1	6.1	14.1
	1930	51.1	100	10.1	19.8	34.9	68.3	5.8	11.3
	1929	29.8	100	11.1	37.3	17.2	57.7	1.2	4.0
	1928	20.1	100	2.3	11.4	14.5	72.2	2.6	12.9
	1932	861.8	100	43.1	5.0	592.8	68.8	163.1	18.9
Georgia	1931	1,393.7	100	69.4	5.0	1,015.6	72.9	242.1	17.4
	1930	1,597.5	100	237.1	14.8	1,096.8	68.7	218.4	13.7
	1929	1,339.8	100	282.2	21.0	918.8	68.6	113.7	8.5
	1928	1,053.2	100	129.5	12.3	743.9	70.6	144.1	13.7
Louisiana	1932	599.5	100	62.1	10.4	131.0	21.8	152.6	25.4
	1931	876.6	100	58.6	6.7	157.3	17.9	203.6	23.2
	1930	704.8	100	91.3	13.0	160.1	22.7	167.4	23.8
	1929	797.7	100	110.6	13.9	205.5	25.8	212.7	26.7
	1928	685.9	100	88.3	12.9	203.1	29.7	178.6	26.0
	1932	1,161.2	100	32.2	2.8	148.6	12.8	162.3	14.0
Mississippi	1931	1,719.4	100	65.4	3.8	285.4	16.6	238.0	13.8
	1930	1,458.5	100	114.9	7.9	299.2	20.5	264.5	18.1
	1929	1,876.0	100	189.7	10.1	352.3	18.8	227.0	12.1
	1928	1,462.0	100	254.3	17.4	258.9	17.7	111.9	7.7
	1932	300.7	100	2.0	.7	52.1	17.3	114.9	38.2
	1931	280.4	100	2.6	.9	85.9	30.6	128.0	45.7
Missouri	1930	153.3	100	11.2	7.3	56.0	36.5	55.6	36.3
	1929	220.9	100	3.1	1.4	57.5	26.0	88.4	40.0
	1928	146.9	100	8.8	6.0	30.2	20.6	52.2	35.5
	1932	67.5	100	2.1	3.1	2.6	3.8	3.4	5.0
	1931	93.8	100	5.1	5.4	1.4	1.5	13.7	14.6
	1930	95.8	100	.5	.5	1.3	1.4	7.8	8.1
New Mexico	1929	86.3	100	4.8	5.6	2.7	3.1	6.3	7.3
	1928	82.2	100	4.9	6.0	6.2	7.5	8.6	10.4
	1932	680.3	100	7.8	1.2	227.8	33.4	223.6	32.9
	1931	771.2	100	7.8	1.0	340.1	44.1	264.6	34.3
	1930	800.6	100	40.4	5.0	410.2	51.2	236.5	29.6
	1929	767.0	100	81.9	10.7	479.4	62.5	151.2	19.7
North Carolina	1928	869.3	100	87.1	10.0	605.5	69.7	129.7	14.9
	1932	1,072.0	100	91.9	8.6	535.8	50.0	386.1	36.0
	1931	1,235.9	100	137.9	11.2	557.4	45.1	414.5	33.5
	1930	856.8	100	126.7	14.8	348.6	40.7	295.1	34.4
	1929	1,125.6	100	321.5	28.6	500.1	44.4	207.9	18.5
	1928	1,187.0	100	162.3	13.7	421.7	35.5	395.5	33.3
Oklahoma	1932	722.2	100	2.8	.4	181.2	25.1	225.6	31.2
	1931	1,010.2	100	12.0	1.2	395.6	39.1	315.2	31.2
	1930	1,015.3	100	26.9	2.6	426.0	42.0	288.6	28.4
	1929	833.1	100	93.1	11.2	440.4	52.9	157.4	18.9
	1928	744.4	100	51.8	7.0	409.9	55.0	144.9	19.5
	1932	467.5	100	12.4	2.7	115.9	24.8	175.3	37.5
Tennessee	1931	578.0	100	39.9	6.9	218.6	37.8	211.3	36.6
	1930	371.4	100	59.7	16.1	128.1	34.5	123.8	33.3
	1929	504.3	100	80.0	15.9	173.4	34.4	163.8	32.5
	1928	423.5	100	88.0	20.8	173.0	40.9	108.0	25.5
	1932	4,307.4	100	414.3	9.6	1,018.6	44.5	1,554.8	36.1
	1931	5,068.8	100	446.5	8.8	2,123.4	41.9	1,719.3	33.9
Texas	1930	3,886.1	100	475.5	12.2	1,330.4	34.2	1,363.2	35.1
	1929	3,803.2	100	974.2	25.6	1,260.5	33.2	955.9	25.1
	1928	4,941.5	100	794.6	16.1	1,856.8	37.6	1,557.1	31.5
	1932	31.3	100	1.5	4.8	19.4	62.0	8.7	27.8
	1931	42.5	100	1.7	4.0	22.9	53.9	16.2	38.1
	1930	42.7	100	3.9	9.1	29.6	69.3	8.4	19.7
Virginia	1929	48.0	100	3.3	6.9	34.1	71.0	9.7	20.2
	1928	44.8	100	2.8	6.3	37.1	82.8	4.7	10.5
	1932	14.0	100	—	—	1.2	8.6	5.4	38.5
	1931	11.7	100	.1	.9	3.0	25.6	3.7	31.6
	1930	6.4	100	.5	7.8	1.0	15.6	1.4	21.9
	1929	8.9	100	.2	2.2	.1	1.1	1.1	12.4
All other ⁴	1928	6.2	100	.8	12.9	.2	3.2	.6	9.7

¹ As reported by the Bureau of the Census.² Untenderable in settlement of futures contracts made subject to sec. 5 of the United States Cotton Futures Act and the regulations of the Secretary of Agriculture thereunder.³ Includes Illinois, Kansas, and Kentucky.

18 STATISTICAL BULLETIN 47, U. S. DEPT. OF AGRICULTURE

TABLE 3.—*Staple length of American upland cotton: Quantities ginned in the United States, by States, crops of 1932, 1931, 1930, 1929, and 1928—Continued*

State	Crop year	1 and 1½ inches	1½ and 1¾ inches	1¾ and 1½ inches	1¾ and 1½ inches	1¼ and longer			
United States		1,000 bales	Per- cent	1,000 bales	Per- cent	1,000 bales	Per- cent	1,000 bales	Per- cent
	1932	1,822.0	14.3	871.8	9.9	622.1	4.9	84.5	0.7
	1931	2,557.1	15.4	1,087.8	6.5	590.0	3.5	224.6	1.4
	1930	1,725.9	12.0	970.9	7.1	393.3	2.9	60.8	.4
	1929	1,693.6	11.7	938.6	6.5	556.1	3.8	119.4	.8
	1928	1,575.8	11.0	794.2	5.6	480.2	3.4	167.9	1.2
		26.5	2.8	3.1	.5	.1	(3)		
	1931	28.0	2.0	5.8	.4	1.3	.1		
	1930	9.2	.6	2.5	.2	.6	(3)		
	1929	3.3	.2	1.4	.1	.1	(3)		
	1928	17.6	1.6	1.8	.2	.5	(3)		
Alabama	1932	23.5	40.0	20.3	34.5	0.3	10.7	.1	(3)
	1931	43.4	44.6	30.8	31.7	4.3	4.4		
	1930	47.8	37.6	29.6	23.3	2.8	2.2		
	1929	65.3	54.1	19.4	16.1	2.8	2.3		
Arizona	1928	40.2	34.2	42.7	36.4	3.0	2.6		
	1932	360.4	28.1	196.8	15.3	120.4	9.4	17.3	1.4
	1931	614.2	33.5	247.0	13.5	81.4	4.4	14.0	.8
	1930	208.8	24.0	106.6	12.3	22.4	2.6	1.7	.2
Arkansas	1929	261.9	18.8	128.0	9.2	45.4	3.2	8.6	.6
	1928	223.6	18.4	130.6	10.7	65.9	5.4	18.9	1.4
	1932	13.4	10.8	57.9	40.5	46.8	37.6	.2	.2
	1931	25.6	15.0	98.0	57.2	33.9	19.8	2.3	1.3
California	1930	41.2	16.1	160.0	62.4	39.2	15.3	.3	.1
	1929	119.6	47.1	104.2	41.0	7.4	2.9		
	1928	47.6	27.9	60.0	35.1	10.8	6.3		
	1932	.1	.6						
Florida	1931	.4	.9	.1	.2				
	1930	.3	.6						
	1929	.2	.7	.1	.3				
	1928	.4	2.0	.2	1.0	.1	.5		
Georgia	1932	53.9	6.3	6.7	.8	1.9	.2	.3	(3)
	1931	54.8	3.9	10.4	.7	3.2	.1		
	1930	36.3	2.3	7.3	.4	1.4	.1	.2	(3)
	1929	21.3	1.6	3.6	.3	.2	(3)		
Louisiana	1928	28.3	2.7	5.4	.5	1.4	.1	.6	.1
	1932	156.4	26.1	68.7	11.5	26.2	4.4	2.4	.4
	1931	233.0	32.3	133.1	15.2	35.4	4.0	5.3	.6
	1930	186.8	26.5	67.2	9.5	28.0	4.0	3.8	.5
Mississippi	1929	170.4	21.3	63.7	8.0	29.6	3.7	5.0	.6
	1928	127.6	18.6	52.1	7.6	27.7	4.0	7.8	1.1
	1932	170.9	14.7	271.5	23.4	328.2	28.2	45.3	3.9
	1931	234.8	13.7	311.0	18.1	368.1	21.4	188.7	11.0
Missouri	1930	209.4	14.4	312.9	21.5	216.6	14.8	39.4	2.7
	1929	202.7	10.8	388.9	20.7	412.4	22.0	97.7	5.2
	1928	127.8	8.7	259.8	17.8	305.9	20.9	121.7	8.3
	1932	107.7	35.8	23.0	7.7	1.0	.3		
New Mexico	1931	53.9	19.2	9.5	3.4	.5	.2		
	1930	22.4	14.6	7.1	4.6	1.0	.7		
	1929	56.2	25.5	13.7	6.2	1.8	.8	.2	.1
	1928	39.3	26.7	14.3	9.7	1.7	1.2	.4	.3
North Carolina	1932	23.6	35.0	25.1	37.2	10.7	15.9		
	1931	36.7	39.1	31.4	33.5	5.5	5.9		
	1930	35.6	37.2	42.5	44.4	8.0	8.3	.1	.1
	1929	40.2	46.8	27.7	32.1	4.6	5.3		
Oklahoma	1928	22.1	26.9	34.0	41.4	6.4	7.8		
	1932	160.4	23.6	47.5	7.0	11.0	1.6	2.1	.3
	1931	126.0	16.3	28.5	3.7	3.8	.5	.4	.1
	1930	81.6	10.2	25.0	3.2	5.6	.7	.7	
South Carolina	1929	45.7	6.0	7.0	.9	1.5	.2	.3	(3)
	1928	30.5	3.5	9.0	1.0	4.9	.6	2.2	.3
	1932	52.4	4.9	5.2	.5	.5	(3)	.1	(3)
	1931	110.8	9.0	14.0	1.1	1.3	.1		
Tennessee	1930	71.6	8.4	12.2	1.4	2.5	.3	.1	(3)
	1929	67.1	6.0	22.6	2.0	6.1	.5	.3	(3)
	1928	172.3	14.5	25.3	2.1	8.2	.7	1.6	.1
	1932	169.5	23.5	81.8	11.3	44.0	6.1	15.8	2.2
Tennessee	1931	169.8	16.8	67.8	6.7	36.0	3.6	12.0	1.2
	1930	140.1	13.8	83.9	8.3	38.4	3.8	10.6	1.0
	1929	81.0	9.7	42.6	5.1	14.4	1.7	3.7	.4
	1928	68.5	9.2	33.9	4.6	23.4	3.1	9.6	1.3
Tennessee	1932	134.7	28.8	24.6	5.3	4.3	.9	.3	(3)
	1931	93.5	16.2	11.7	2.0	2.5	.4	.5	.1
	1930	48.0	12.9	11.0	3.0	.8	.2		
	1929	74.9	14.8	10.3	2.0	1.6	.3	.3	.1
	1928	42.1	9.9	10.4	2.5	1.4	.3	.5	.1

^a Less than 0.05 percent.

TABLE 3.—*Staple length of American upland cotton: Quantities ginned in the United States, by States, crops of 1932, 1931, 1930, 1929, and 1928—Continued*

State	Crop year	1 and 1½ inches		1¾ and 1½ inches		1½ and 1¾ inches		1¾ and 2 inches		1¼ inches and longer	
Texas	(1932)	1,000 bales	Percent	1,000 bales	Percent	1,000 bales	Percent	1,000 bales	Percent	1,000 bales	Percent
	1931	362.3	8.4	37.5	0.9	019.7	0.5	0.2	(*)	0.1	(3)
	1930	586.3	15.1	101.0	2.6	25.7	.7	3.9	.1	.1	(3)
	1929	480.8	12.0	103.2	2.7	26.0	.7	2.4	.1	.2	(3)
	1928	586.8	11.9	113.0	2.3	26.9	.5	5.8	.1	.5	(3)
	1932	1.5	4.8	.2	.6						
	1931	1.7	4.0								
Virginia	1930	.7	1.7	.1	.2						
	1929	.8	1.7	.1	.2						
	1928	.2	.4								
	(1932)	4.8	34.3	1.9	13.6	.6	4.3	.1	.7		
All other *	1931	2.2	18.8	1.8	15.4	.8	6.8	.1	.9		
	1930	1.8	28.1	1.4	21.9	.3	4.7				
	1929	2.2	24.7	2.1	23.7	2.2	24.7	.9	10.1	.1	1.1
	(1928)	.9	14.5	1.7	27.4	1.0	10.1	.7	11.3	.3	4.9

^a Less than 0.05 percent.^b Includes Illinois, Kansas, and Kentucky.TABLE 4.—*Average staple length ¹ of American upland cotton ginned in the United States, by States and districts thereof, crops of 1932, 1931, 1930, 1929, and 1928 (fig. 1)*

State and district ²	1932	1931	1930	1929	1928
	Sixteenth inches 15.45	Sixteenth inches 15.44	Sixteenth inches 15.22	Sixteenth inches 15.11	Sixteenth inches 15.18
United States					
Alabama	14.54	14.58	14.19	14.08	14.34
District 1	14.56	14.66	14.12	14.05	14.35
District 2	14.68	14.55	14.16	13.91	14.30
District 3	14.76	14.47	14.34	14.11	14.48
District 4	14.42	14.58	14.16	14.16	14.32
District 5	14.49	14.48	14.46	14.27	14.37
District 6	14.41	14.65	14.42	14.11	14.39
Arizona	16.90	16.64	16.36	16.37	16.51
Arkansas	16.24	16.08	15.67	15.46	15.62
District 1	15.43	15.35	14.42	14.07	14.47
District 2	15.32	15.64	14.86	14.63	14.89
District 3	15.33	15.86	15.32	14.65	14.81
District 4	16.73	16.33	16.04	16.05	16.29
District 5	15.77	15.86	15.36	15.03	15.05
California	17.66	17.34	17.37	16.85	16.52
Florida	14.41	14.60	14.43	14.19	14.60
Georgia	14.80	14.73	14.55	14.41	14.59
District 1	14.90	14.61	14.42	14.39	14.69
District 2	14.98	14.78	14.70	14.44	14.74
District 3	14.78	14.76	14.43	14.39	14.45
District 4	14.52	14.70	14.46	14.39	14.37
Louisiana	15.71	15.96	15.61	15.48	15.46
District 1	14.43	14.80	14.48	14.49	14.74
District 2	15.79	16.22	15.86	15.64	15.71
District 3	16.26	16.41	15.99	15.90	15.73
District 4	14.63	15.06	14.93	15.24	15.25

¹ Average calculated by multiplying the number of bales in each length group by the midpoint of the group expressed in sixteenths of an inch (that is, 1½ to 15.5), summing, and dividing by the total number of bales. In making calculations, the midpoints of the groups "shorter than 7/8 inch" and "1¼ inches and longer" were considered to be 13.5 and 20.5, respectively.² For location of districts into which States are subdivided, see maps accompanying tables for individual States, by districts, pp. 87-105.

TABLE 4.—Average staple length¹ of American upland cotton ginned in the United States, by States and districts thereof, crops of 1932, 1931, 1930, 1929, and 1928 (fig. 1)—Continued

State and district ²	1932	1931	1930	1929	1928
	Sixteenth inches 16.94	Sixteenth inches 16.92	Sixteenth inches 16.27	Sixteenth inches 16.51	Sixteenth inches 16.45
Mississippi.....					
District 1.....	18.13	18.30	17.80	18.05	18.09
District 2.....	17.00	16.71	16.28	16.01	16.34
District 3.....	15.24	14.99	14.85	14.52	14.18
District 4.....	15.28	15.18	15.14	15.41	14.35
Missouri.....	15.83	15.44	15.25	15.62	15.08
New Mexico.....	16.97	16.61	16.99	16.63	16.64
North Carolina.....	15.58	15.29	15.08	14.75	14.60
District 1.....	15.33	15.19	14.98	14.73	14.70
District 2.....	16.04	15.64	15.45	15.10	14.06
District 3.....	15.73	15.31	15.10	14.73	14.05
District 4.....	15.29	15.35	15.02	14.48	14.55
Oklahoma.....	14.80	14.94	14.92	14.60	15.09
District 1.....	14.84	14.70	14.67	14.37	14.82
District 2.....	14.99	15.25	15.29	15.19	15.64
District 3.....	15.02	15.40	15.29	14.94	15.43
District 4.....	15.63	16.24	16.69	16.44	16.66
District 5.....	14.62	15.07	14.71	14.34	14.73
South Carolina.....	15.98	15.55	15.49	15.02	15.10
District 1.....	15.69	15.34	15.25	14.86	15.02
District 2.....	16.65	16.65	16.35	15.78	15.42
District 3.....	18.33	15.64	15.68	15.17	15.40
District 4.....	15.37	15.13	15.11	14.76	14.52
Tennessee.....	15.62	15.20	15.03	15.04	14.84
District 1.....	15.86	15.48	15.40	15.52	15.30
District 2.....	15.57	15.14	14.94	14.86	14.62
District 3.....	14.55	14.44	14.31	14.10	14.06
District 4.....	15.33	14.61	14.72	14.75	14.71
Texas.....	14.98	15.08	15.14	14.86	14.00
District 1.....	17.56	17.08	17.24	16.84	16.98
District 2.....	14.86	14.36	14.44	13.87	14.43
District 3.....	15.02	14.55	14.44	14.12	14.70
District 4.....	15.12	14.81	14.44	14.42	14.07
District 5.....	14.85	14.68	14.62	14.17	14.89
District 6.....	14.91	14.92	14.66	14.51	14.75
District 7.....	15.06	15.31	15.31	15.19	15.13
District 8.....	14.44	14.98	14.71	14.50	14.78
District 9.....	15.11	15.34	15.25	15.18	15.21
District 10.....	15.03	15.63	15.42	15.60	14.90
District 11.....	15.26	15.76	15.90	16.28	15.47
Virginia.....	14.85	14.92	14.65	14.67	14.55
All other ³	18.19	15.98	16.05	17.37	17.08

¹ Average calculated by multiplying the number of bales in each length group by the midpoint of the group expressed in sixteenths of an inch (that is, $1\frac{1}{16}$ = 15.5), summatting, and dividing by the total number of bales. In making calculations, the midpoints of the groups "shorter than $\frac{3}{8}$ inch" and " $1\frac{1}{4}$ inches and longer" were considered to be 13.5 and 20.5, respectively.

² For location of districts into which States are subdivided, see maps accompanying tables for individual States, by districts, pp. 87-105.

³ Includes Illinois, Kansas, and Kentucky.




FIGURE I.—AVERAGE STAPLE LENGTH OF AMERICAN UPLAND COTTON GINNED IN THE UNITED STATES, BY STATES, CROPS OF 1932, 1931, 1930, 1929, AND 1928.

The cotton-growing States are here arranged according to average staple length of cotton grown in 1932. The average staple length of cotton produced in the United States has increased from year to year since 1929, and the average for each of the individual States is greater for 1932 than for 1929 (table 4).

22 STATISTICAL BULLETIN 47, U. S. DEPT. OF AGRICULTURE

TABLE 5.—Grade of American upland cotton: Quantities ginned in the United States during specified periods,¹ by States, crops of 1932, 1931, 1930, 1929, and 1928

[Quantities are given in running bales, except that round bales are counted as half bales. Linters are not included]

UNITED STATES

Grade	Total, crop of—					Ginned prior to Oct. 1				
	1932	1931	1930	1929	1928	1932	1931	1930	1929	1928
All grades ⁴	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales
	12,701.3	10,615.2	13,732.2	14,510.0	14,268.2	4,834.9	5,407.4	6,300.0	5,869.0	4,055.2
Extra White	460.0	435.8	500.0	468.7	466.7	58.2	54.6	61.1	45.3	63.1
Above 3-G.M.	1.8	.6	1.0	(3)	(3)	.4	.7	(3)	(3)	(3)
3-G.M.	106.9	76.7	123.1	180.4	146.5	20.1	31.6	36.3	27.8	43.1
4-S.M.	132.8	174.2	214.4	167.8	141.9	17.9	12.7	23.4	16.3	14.4
5-M.	88.2	88.1	107.6	80.5	93.6	11.0	2.5	.7	1.0	3.6
6-S.L.M.	99.6	62.2	31.1	32.5	18.0	8.6	5.7	—	.1	.7
7-L.M.	28.9	29.7	4.8	7.5	6.7	2.0	1.7	—	.1	1.3
Below 7-L.M. ⁴	1.8	4.3	18.6	(8)	(8)	—	—	—	(8)	(8)
White	9,945.3	15,039.7	11,042.2	12,237.1	11,940.1	4,449.7	5,130.4	5,882.1	5,463.0	4,060.2
1-M.F.	—	—	—	—	—	—	—	—	.1	.4
2-S.G.M.	1.2	10.9	13.0	38.7	41.8	.8	7.5	11.8	12.4	34.3
3-G.M.	251.3	940.0	892.3	863.9	1,630.0	188.9	570.0	764.7	713.1	1,149.0
4-S.M.	3,147.0	5,873.4	4,364.0	3,877.9	4,845.6	1,891.2	2,711.6	6,027.6	6,235.2	4,230.2
5-M.	4,474.5	5,232.3	4,211.7	4,399.1	3,250.4	1,961.9	1,561.8	1,718.4	1,806.1	940.2
6-S.L.M.	1,569.2	1,759.2	1,749.7	1,881.7	1,387.2	351.2	257.5	323.8	478.1	240.4
7-L.M.	330.3	640.3	576.9	805.4	447.7	46.5	20.6	30.2	83.2	47.0
8-S.G.O. ⁴	116.3	421.9	114.6	290.1	247.5	7.7	1.3	4.6	14.2	7.7
9-G.O. ⁴	55.5	160.8	20.0	80.1	89.2	1.5	.1	1.0	3.4	1.4
Spotted	2,216.9	1,048.2	1,214.3	1,567.3	1,677.9	322.3	220.4	341.5	382.4	217.6
3-G.M.	193.6	115.3	147.2	47.1	142.9	35.5	43.9	70.5	18.6	31.1
4-S.M.	1,054.0	428.5	557.0	648.9	792.7	192.5	142.1	107.1	228.3	114.6
5-M.	673.0	247.9	335.2	564.3	466.1	87.3	31.7	63.3	315.6	62.0
6-S.L.M. ⁴	217.5	185.2	143.7	234.4	196.1	6.4	2.5	4.3	17.4	9.7
7-L.M. ⁴	78.8	71.3	31.2	72.6	80.1	.6	.2	.3	2.5	.9
Yellow Tinged	33.8	23.9	55.6	139.9	41.7	3.1	1.3	13.6	8.3	3.2
2-S.G.M.	—	—	.2	.1	.5	—	—	—	.2	.1
3-G.M.	2.7	1.6	7.4	2.7	4.7	.8	.1	3.2	1.1	.5
4-S.M.	10.7	4.5	20.7	25.1	11.3	1.9	1.0	7.1	4.6	1.9
5-M. ⁴	9.0	7.3	14.9	42.0	6.7	.3	.2	2.6	2.1	.5
6-S.L.M. ⁴	9.7	8.1	10.1	40.0	9.6	.1	—	.4	.3	.1
7-L.M. ⁴	1.7	2.4	2.3	30.0	8.9	—	—	.1	—	—
Light Yellow Stained	.3	.7	1.8	17.2	1.7	.1	.1	.4	.5	.4
3-G.M.	.1	.1	—	.2	.4	.1	—	—	—	.1
4-S.M. ⁴	.1	.2	.6	3.3	.5	—	—	2	.2	.1
5-M. ⁴	.1	.4	1.2	13.7	.8	—	.1	.2	.3	.2
Yellow Stained	.1	.1	.6	10.4	.8	—	—	—	.1	—
3-G.M.	—	—	—	.1	.2	—	—	—	—	.1
4-S.M. ⁴	—	—	.2	1.1	.3	—	—	—	—	—
5-M. ⁴	.1	.1	.4	9.2	.3	—	—	—	—	—
Gray	10.1	12.5	4.6	18.9	21.6	.8	.3	.0	—	.2
3-G.M.	1.2	.5	.4	.2	3.5	—	—	.1	.1	.1
4-S.M.	5.9	6.9	3.2	7.0	12.8	.7	—	.0	—	.1
5-M. ⁴	3.0	5.1	1.0	10.8	5.3	.1	.2	.2	—	—
Blue Stained	—	—	.1	.3	3.0	1.6	—	—	—	—
3-G.M. ⁴	—	—	—	.1	.1	.4	—	—	—	—
4-S.M. ⁴	—	—	.1	—	1.8	.9	—	—	—	—
5-M. ⁴	—	—	.2	1.1	.3	—	—	—	—	—
No grade ^{4,6}	34.8	54.2	12.2	56.5	176.1	.7	.3	.3	.4	1.4

See footnotes at end of table (p. 60).

TABLE 5.—*Grade of American upland cotton: Quantities ginned in the United States during specified periods,¹ by States, crops of 1932, 1931, 1930, 1929, and 1928—Continued*

UNITED STATES—Continued

Grade	Ginned during October					Ginned during November				
	1932	1931	1930	1929	1928	1932	1931	1930	1929	1928
	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales
All grades ²	4,408.3	6,711.5	4,553.4	4,981.3	5,193.9	2,385.7	2,892.7	1,968.2	1,954.0	2,391.3
Extra White	213.5	151.6	154.0	169.4	137.3	150.3	142.4	139.7	158.7	112.8
Above 3-G.M.	1.2	.2	.2	(3)	(3)				.1	(3)
3-G.M.	58.5	43.3	59.5	108.1	84.4	27.1	1.2	25.9	42.2	18.4
4-S.M.	51.6	80.1	83.0	46.4	40.7	53.4	64.1	82.4	64.2	52.4
5-M.	32.9	16.9	10.7	11.6	8.7	30.5	31.0	29.2	37.3	37.4
6-S.L.M.	55.0	0.6	.5	3.0	2.1	30.2	27.7	2.0	11.3	3.8
7-L.M.	14.3	3.3	.1	.3	1.4	8.8	18.4		3.7	.8
Below 7-L.M. ⁴		1.2		(6)	(6)	.3		.1	(6)	(6)
White	3,708.9	6,196.2	3,934.9	4,228.5	4,403.7	1,337.6	2,488.3	1,621.4	1,461.3	1,802.3
1-M.F.				.1	.3					
2-S.G.M.	.3	3.2	.9	25.5	6.1	.1	.2	.2	.8	1.2
3-G.M.	52.1	327.7	97.4	128.5	416.3	9.8	31.1	26.5	16.6	50.5
4-S.M.	979.1	2,601.7	1,075.1	1,261.6	2,018.5	240.6	442.7	221.9	198.9	492.6
5-M.	1,855.1	2,436.6	1,779.7	1,902.1	1,450.1	556.1	1,079.2	592.0	520.9	687.1
6-S.L.M.	699.5	684.4	751.5	728.5	442.0	397.1	650.3	525.5	414.7	445.5
7-L.M.	101.9	122.4	200.2	162.5	97.8	98.0	214.9	214.1	236.0	119.3
8-S.G.O. ⁴	17.0	20.2	25.5	19.1	26.5	31.3	64.9	37.0	68.1	66.3
9-G.O.A.	3.9	1.0	4.6	2.6	6.1	4.6	5.0	4.2	5.3	9.8
Spotted	477.2	355.4	441.1	554.6	580.6	884.6	253.3	199.6	300.0	398.4
3-G.M.	55.8	53.1	61.1	24.3	74.9	95.6	15.6	7.4	3.5	23.9
4-S.M.	281.2	183.0	243.2	283.2	344.2	476.3	86.6	79.1	96.3	206.3
5-M.	122.8	88.2	111.0	207.2	131.2	278.7	88.5	76.8	122.3	119.0
6-S.L.M. ⁴	14.6	28.0	21.4	38.8	28.4	31.5	53.6	30.3	60.4	33.6
7-L.M. ⁴	2.8	3.1	4.4	3.1	1.9	2.5	9.0	6.0	17.6	15.6
Yellow Tinged	1.9	3.2	10.2	24.6	5.4	8.2	2.7	4.4	28.5	4.0
2-S.G.M.					.1					.1
3-G.M.	.6	1.0	3.8	1.1	1.0	1.4	.4	.2	.3	.9
4-S.M.	.9	1.4	10.3	10.1	3.1	5.4	1.2	1.4	5.0	1.6
5-M. ⁴	.4	.6	3.8	10.1	.9	1.2	.8	1.8	8.4	1.3
6-S.L.M. ⁴	.2	.9	2.9	.2	.2	.2	.2	.8	7.5	.6
7-L.M. ⁴	.1	.4	.4	.1			.1	.2	7.3	.4
Light Yellow Stained	.1	.2	.5	1.2	.7	.1		.1	.5	.2
3-G.M.					.2				.1	
4-S.M. ⁴	.1		.2	.5	.2				.1	.1
5-M. ⁴		.2	.3	.7	.3	.1		.1	.3	.1
Yellow Stained										
3-G.M.				.1	.1					
4-S.M. ⁴				.1	.2					
5-M. ⁴				.1			.1	.2	.8	.1
Gray	4.9	4.2	2.7	1.8	1.9	3.1	5.4	.7	2.4	8.1
3-G.M.	.9	.1	.3	.1	.1	.2			.1	.8
4-S.M.	3.2	2.2	2.0	1.5	1.0	1.5	3.7	.5	1.3	4.8
6-M. ⁴	.8	1.9	.4	.2	.8	1.4	1.6	.2	1.0	2.5
Blue Stained					.1			.1		
3-G.M. ⁴					.1					
4-S.M. ⁴								.1		
5-M. ⁴										
No grade ^{4,6}	1.8	.7	.9	.7	4.1	1.8	.4	2.1	1.8	4.5

See footnotes at end of table.

TABLE 5.—*Grade of American upland cotton: Quantities ginned in the United States during specified periods,¹ by States, crops of 1932, 1931, 1930, 1929, and 1928—Continued*

UNITED STATES—Continued

Grade	Ginned Dec. 1-Jan. 15, crop of—					Ginned after Jan. 15, crop of—				
	1932	1931	1930	1929	1928	1932	1931	1930	1929	1928
	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales
All grades ²	778.6	973.9	751.6	1,316.3	1,323.1	293.8	629.7	159.0	367.5	404.7
Extra White.....	26.5	75.5	114.6	72.6	70.0	11.5	11.7	31.2	22.7	23.5
Above 3-G.M.				(3)	(3)				(3)	(3)
3-G.M.	1.2	.6	1.4	2.3	.5					.1
4-S.M.	9.6	16.9	25.3	38.4	26.0	.3	.4	.3	2.5	8.4
5-M.	10.3	35.1	62.8	22.4	35.0	3.5	2.6	4.2	8.2	8.9
6-S.L.M.	3.7	16.4	19.4	7.5	6.8	4.1	5.8	9.2	10.6	4.6
7-L.M.	1.5	5.0	1.7	2.0	1.7	2.3	1.3	3.0	1.4	1.5
Below 7-L.M. ⁴2	1.5	4.0	(6)	(6)	1.3	1.6	14.5	(6)	(6)
White.....	357.8	750.5	431.8	860.8	762.8	91.3	474.3	72.0	223.5	182.1
1-M.F.				1	2					
2-S.G.M.										
3-G.M.4	8.0	3.1	5.3	12.2	.1	3.2	.6	.4	1.4
4-S.M.	26.6	83.7	33.0	54.3	94.5	10.5	33.7	6.4	10.7	9.8
5-M.	83.7	124.4	103.4	146.3	140.0	17.7	32.2	18.2	23.7	18.0
6-S.L.M.	108.6	129.9	131.6	226.3	218.5	12.8	37.1	17.3	36.1	31.8
7-L.M.	68.3	191.7	115.0	260.6	145.2	15.6	90.7	17.4	63.1	38.4
8-S.G.O. ⁴	44.2	174.1	37.4	131.7	104.1	16.1	161.4	10.1	57.0	52.0
9-G.O. ⁴	27.0	38.7	8.2	36.3	42.1	18.5	116.0	2.0	32.5	29.8
Spotted.....	363.1	135.0	184.1	260.2	382.9	169.7	84.1	48.0	70.1	98.4
3-G.M.	5.3	2.4	2.1	.7	12.0	1.4	.3	.1		1.0
4-S.M.	93.1	13.7	33.7	36.4	115.5	10.9	3.1	3.9	4.7	12.1
5-M.	156.4	27.5	69.7	97.5	123.8	27.8	12.0	14.4	21.7	30.1
6-S.L.M.	86.3	62.5	65.3	91.8	83.1	78.7	38.6	22.4	28.0	42.0
7-L.M. ⁴	22.0	28.9	13.3	33.8	48.5	50.9	30.1	7.2	15.7	13.2
Yellow Tinged.....	10.9	6.0	12.4	62.2	18.1	9.7	10.7	6.0	16.3	10.1
2-S.G.M.					1					
3-G.M.1	.2	.2	.2	.2					.1
4-S.M.	2.0	.6	1.7	5.1	4.0	.5	.3	.2	3	.7
5-M. ⁴	4.2	2.2	4.2	19.2	2.7	2.9	3.5	2.5	2.2	1.3
6-S.L.M. ⁴	4.1	2.0	5.3	23.9	4.1	5.3	5.1	2.7	5.4	4.6
7-L.M. ⁴6	.5	1.0	13.8	5.0	1.0	1.8	.6	8.4	3.4
Light Yellow Stained.....	.2	.3	12.6	.2		.2	.5	2.4		.2
3-G.M.1				.1
4-S.M. ⁴2	.1	2.1							.1
5-M. ⁴2	10.5								.1
Yellow Stained.....										.1
3-G.M.1
4-S.M. ⁴1
5-M. ⁴1
Gray.....	1.1	1.5	.3	12.7	10.1	.2	1.1		2.0	1.3
3-G.M.1
4-S.M.5	.8	.1	4.7	6.6					.3
5-M. ⁴6	.6	.2	8.0	1.1					.9
Blue Stained.....										.6
3-G.M. ⁴3
4-S.M. ⁴8
5-M. ⁴3
No grade ^{4,6}	19.2	5.2	7.7	24.2	78.5	11.3	47.6	1.2	20.4	87.6

See footnotes at end of table.

TABLE 5.—*Grade of American upland cotton: Quantities ginned in the United States during specified periods,¹ by States, crops of 1932, 1931, 1930, 1929, and 1928—Continued*

ALABAMA

Grade	Total, crop of—					Ginned prior to Oct. 1				
	1932	1931	1930	1929	1928	1932	1931	1930	1929	1928
	1,000 bales 933.7	1,000 bales 1,385.0	1,000 bales 1,444.9	1,000 bales 1,307.7	1,000 bales 1,096.0	1,000 bales 412.0	1,000 bales 531.0	1,000 bales 582.6	1,000 bales 578.1	1,000 bales 336.0
All grades ²3		.1		.4				.1	.2
Extra White.....										
<i>Above 3-G.M.</i>										
3-G.M.										.1
4-S.M.										.1
5-M.										
6-S.L.M.										
7-L.M.										
Below 7-L.M. ⁴	.2									
.1										
White.....	783.1	1,327.8	1,275.0	1,137.3	931.7	382.8	499.6	539.6	510.9	310.6
<i>1-M.F.</i>										
2-S.G.M.		2.9	.5	1.6	1.5					
3-G.M.	9.9	77.2	49.2	37.8	106.8	5.8	47.3	44.1	31.7	46.1
4-S.M.	271.9	578.8	508.2	430.0	545.8	154.6	253.5	318.7	263.2	197.1
5-M.	440.7	579.8	493.6	484.1	236.4	191.3	178.9	152.7	187.1	60.0
6-S.L.M.	58.4	87.5	138.3	124.1	37.1	10.8	17.4	18.7	26.8	6.5
7-L.M.	1.5	1.6	65.9	42.6	3.5	.3	.3	4.2	.7	.2
8-S.G.O. ⁴	.6		15.8	28.2	.5			.6	.1	
9-G.O. ⁴	.1		3.5	10.9	.1			.1		
Spotted.....	149.0	55.3	161.1	154.9	160.0	48.6	30.7	40.7	64.1	24.7
<i>3-G.M.</i>										
4-S.M.	6.7	11.6	19.6	4.9	20.1	4.0	6.4	12.1	3.0	5.8
5-M.	82.1	36.3	75.6	97.7	110.2	31.6	21.0	21.4	48.5	14.6
6-S.L.M. ⁴	56.6	6.8	47.0	44.2	27.2	12.5	3.2	6.7	12.0	4.1
7-L.M. ⁴	3.1	.5	14.6	7.2	2.2	.5	.1	.4	.6	.2
Yellow Tinged.....	.6	.1	4.3	.9	.3			.1		
<i>2-S.G.M.</i>										
3-G.M.	.9	1.0	6.8	8.6	3.8	.6	.6	2.1	3.0	.5
4-S.M.	.1									
5-M. ⁴	.6	.7	3.0	4.2	2.2	.4	.4	1.2	2.2	.2
6-S.L.M. ⁴	.2	.2	2.1	2.9	.5	.1	.1	.5	.4	.1
7-L.M. ⁴										
Light Yellow Stained.....										
<i>3-G.M.</i>										
4-S.M. ⁴										
5-M. ⁴										
Yellow Stained.....										
<i>3-G.M.</i>										
4-S.M. ⁴										
5-M. ⁴										
Gray.....										
<i>3-G.M.</i>										
4-S.M.	.2	.7		3.4	.4					
5-M. ⁴	.1	.5		1.1	.3					
Blue Stained.....										
<i>3-G.M.⁴</i>										
4-S.M. ⁴										
5-M. ⁴										
No grade ^{4,6}2		1.7	2.4	.1					

See footnotes at end of table.

TABLE 5.—Grade of American upland cotton: Quantities ginned in the United States during specified periods,¹ by States, crops of 1932, 1931, 1930, 1929, and 1928—Continued

ALABAMA—Continued

Grade	Ginned during October					Ginned during November				
	1932	1931	1930	1929	1928	1932	1931	1930	1929	1928
	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales
All grades ²	330.0	647.9	695.4	511.8	482.0	150.6	185.8	202.0	109.9	206.2
Extra White.....					.2	.1				
Above 3-G.M.										
3-G.M.										
4-S.M.					.1					
5-M.					.1					
6-S.L.M.										
7-L.M.						.1				
Below 7-L.M. ⁴										
White.....	280.4	620.1	520.6	434.6	425.0	115.2	180.1	173.4	95.7	160.0
1-M.F.										
2-S.G.M.		.8			.5					
3-G.M.	3.7	27.3	4.9	5.4	51.9	.4	2.0		.7	6.8
4-S.M.	93.7	267.8	165.8	142.7	252.4	21.5	48.9	21.3	16.7	80.4
5-M.	165.8	287.2	237.6	223.4	99.7	72.3	105.6	89.5	40.8	63.1
6-S.L.M.	23.0	45.2	68.9	55.7	17.7	19.8	23.3	41.9	26.6	9.0
7-L.M.	.2	1.0	33.9	6.0	2.4	.8	.2	16.9	8.2	.4
8-S.G.O. ⁴			7.3	.9	.4	.4		3.4	2.3	
9-G.O. ⁴			2.2	.2				.4	.4	
Spotted.....	43.3	18.4	71.3	72.2	56.5	35.2	4.9	27.6	12.7	45.1
3-G.M.	1.7	4.5	6.8	1.7	10.0	.6	.6	.5	.2	2.1
4-S.M.	25.8	11.4	38.5	42.0	38.9	17.3	3.4	12.3	6.3	35.7
5-M.	15.4	2.2	21.4	25.6	6.4	16.4	.8	10.8	4.7	7.1
6-S.L.M. ⁴4	.3	3.8	2.8	.9	.8		3.4	1.4	.2
7-L.M. ⁴8	.1	.3	.1	.1	.6	.1	
Yellow Tinged.....	.2	.2	3.2	4.6	.2		.2	.4	.3	.8
2-S.G.M.										
3-G.M.			.5	.2						
4-S.M.	.2	.1	1.5	1.8	.2		.2	.1	.1	.4
5-M. ⁴1	.9	2.1				.2	.2	.1
6-S.L.M.			.1	.5				.1		
7-L.M. ⁴2							
Light Yellow Stained.....				.1	.2					.1
3-G.M.										
4-S.M. ⁴1	.1					
5-M. ⁴1				.1	
Yellow Stained.....					.1			.1		.1
3-G.M.										
4-S.M. ⁴1					
5-M. ⁴1		.1
Gray.....	.1	.2					.1	.5		1.1
3-G.M.		.1								
4-S.M.			.1				.1	.4		.4
5-M. ⁴1					.1		.7
Blue Stained.....										
3-G.M. ⁴										
4-S.M. ⁴										
5-M. ⁴										
No grade ^{4,6}2	.1	.1			.6	

See footnotes at end of table.

TABLE 5.—*Grade of American upland cotton: Quantities ginned in the United States during specified periods,¹ by States, crops of 1932, 1931, 1930, 1929, and 1928—Continued*

ALABAMA—Continued

Grade	Ginned Dec. 1-Jan. 15, crop of—					Ginned after Jan. 15, crop of—				
	1932	1931	1930	1929	1928	1932	1931	1930	1929	1928
	1,000 bales 35.3	1,000 bales 16.9	1,000 bales 56.9	1,000 bales 95.5	1,000 bales 63.1	1,000 bales 5.8	1,000 bales 3.4	1,000 bales 8.0	1,000 bales 12.4	1,000 bales 9.3
All grades ²										
Extra White.....	.2									
Above 3-G.M.....										
3-G.M.....										
4-S.M.....										
5-M.....										
6-S.L.M.....										
7-L.M.....	.1									
Below 7-L.M. ⁴1									
White.....	16.2	15.9	36.8	84.7	31.8	2.5	3.1	4.6	11.4	4.3
1-M.F.....										
2-S.G.M.....										
3-G.M.....	.6	.2			1.9					.1
4-S.M.....	1.8	6.9	2.1	5.6	14.1	.3	1.7	.3	1.8	1.8
5-M.....	9.6	7.0	12.4	10.6	12.0	1.7	1.1	1.4	2.2	1.6
6-S.L.M.....	4.4	1.4	8.0	13.6	3.4	.4	.2	.8	1.4	.5
7-L.M.....	.2		9.8	24.8	.3		.1	1.1	2.9	.2
8-S.G.O. ⁴2		3.7	20.6				.8	2.3	.1
9-G.O. ⁴6	9.5	.1		.1	.2	.8	
Spotted.....	18.7	1.0	18.2	5.4	28.9	3.2	.3	3.3	.5	4.8
3-G.M.....	.3	.1	.2		2.1	.1				.1
4-S.M.....	6.6	.4	3.3	.9	18.6	.8	.1	.1		2.4
5-M.....	10.6	.5	7.5	1.8	7.8	1.6	.1	.6	.1	1.8
6-S.L.M. ⁴	1.0		6.1	2.2	.4	.4	.1	.9	.2	.5
7-L.M. ⁴2		1.1	.5		.3		1.7	.2	
Yellow Tinged.....	.1		1.1	.5	2.1				.2	.2
2-S.G.M.....										
3-G.M.....										.1
4-S.M.....			.2	.1	1.3					.1
5-M. ⁴1		.5	.1	.3					
6-S.L.M. ⁴4	.2						
7-L.M. ⁴1						
Light Yellow Stained.....										
3-G.M.....										
4-S.M. ⁴										
5-M. ⁴										
Yellow Stained.....										
3-G.M.....										
4-S.M. ⁴										
5-M. ⁴										
Gray.....										
3-G.M.....										
4-S.M.....										
5-M. ⁴										
Blue Stained.....										
3-G.M. ⁴										
4-S.M. ⁴										
5-M. ⁴										
No grade ^{4,6}1		.8	2.0		.1		.1	.3	

See footnotes at end of table.

TABLE 5.—*Grade of American upland cotton: Quantities ginned in the United States during specified periods,¹ by States, crops of 1932, 1931, 1930, 1929, and 1928—Continued*

ARIZONA

Grade	Total, crop of—					Ginned prior to Oct. 1				
	1932	1931	1930	1929	1928	1932	1931	1930	1929	1928
	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales
All grades ²	58.8	97.3	127.2	120.7	117.4	9.8	10.2	19.6	15.1	21.0
Extra White.....	45.4	82.0	109.4	97.7	104.6	9.8	10.1	19.6	15.0	20.0
Above 3-G.M.....	.2					.2				
3-G.M.....	11.4	4.8	23.4	19.5	32.6	6.1	2.4	8.2	5.6	14.0
4-S.M.....	19.5	29.0	51.6	55.3	46.1	3.1	5.4	11.1	8.4	5.4
5-M.....	9.4	32.5	22.6	17.5	23.6	.4	1.6	.3	.8	.8
6-S.L.M.....	3.8	13.3	8.3	5.0	2.1		.6		.1	.1
7-L.M.....	.9	1.5	1.0	.4	.2		.1		.1	
Below 7-L.M. ⁴2	.3	2.5							
White.....	.1	7.3		1.3	.4					.1
1-M.F.....										
2-S.G.M.....				.7						
3-G.M.....						.1				
4-S.M.....						.1				
5-M.....		1.7								
6-S.L.M.....		3.0								
7-L.M.....	.1	2.2		.1						.1
8-S.G.O. ¹4			.2					
9-G.O. ¹5						
Spotted.....	12.9	4.8	17.7	20.6	12.2		.1			.1
3-G.M.....	.3	.1	.1							.1
4-S.M.....	8.8	2.3	6.6	5.9	5.8					
5-M.....	3.5	1.9	6.2	10.4	3.8					
6-S.L.M. ⁴3	.5	3.9	3.8	2.1					
7-L.M. ⁴9	.5						
Yellow Tinged.....		.1	.1		.2					
2-S.G.M.....										
3-G.M.....										
4-S.M.....						.1				
5-M. ⁴1				.1				
6-S.L.M. ⁴1			.1				
7-L.M. ⁴										
Gray.....					.2					
3-G.M.....										
4-S.M.....					.1					
5-M. ⁴1					
No grade ⁴4	3.1		.9						

Grade	Ginned during October					Ginned during November				
	1932	1931	1930	1929	1928	1932	1931	1930	1929	1928
	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales
All grades ²	14.7	20.3	31.4	34.6	27.2	15.8	16.6	28.0	35.5	27.4
Extra White.....	13.4	18.7	31.0	33.2	25.1	8.8	16.2	27.5	28.9	26.0
Above 3-G.M.....										
3-G.M.....	4.5	2.1	11.8	10.5	13.7	.6	.2	3.4	3.0	4.2
4-S.M.....	0.9	9.0	17.5	16.3	8.3	6.5	7.7	18.2	17.8	14.1
5-M.....	1.7	6.2	1.7	4.7	2.6	1.6	7.4	5.5	7.1	7.4
6-S.L.M.....	.3	.5		1.5	.4	.1	.8	.4	.9	.3
7-L.M.....				.2	.1		.1		.1	
Below 7-L.M. ⁴										

See footnotes at end of table.

TABLE 5.—*Grade of American upland cotton: Quantities ginned in the United States during specified periods,¹ by States, crops of 1932, 1931, 1930, 1929, and 1928—Continued*

ARIZONA—Continued

Grade	Ginned during October					Ginned during November				
	1932	1931	1930	1929	1928	1932	1931	1930	1929	1928
	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales
White.....										
1-M.F.										
2-S.G.M.					.7					
3-G.M.						.1				
4-S.M.						.1				
5-M.										
6-S.L.M.										
7-L.M.							.1			
8-S.G.O. ⁴										
9-G.O. ⁴										
Spotted.....	1.3	1.6	0.4	.7	1.9	6.8	0.4	0.5	6.6	1.4
3-G.M.			.1				.3			
4-S.M.	.5	.4	.3	.1	1.3	4.9	.3	.4	1.3	.9
5-M.	.8	.9	.4	.6	.6	1.4	.1	.1	3.8	.4
6-S.L.M. ⁴		.3		.2			.2		1.4	.1
7-L.M. ⁴										.1
Yellow Tinged.....										
2-S.G.M.										
3-G.M.										
4-S.M.										
5-M. ⁴										
6-S.L.M. ⁴										
7-L.M. ⁴										
Gray.....										
3-G.M.										
4-S.M.										
5-M. ⁴										
No grade ^{4,6}1			

Grade	Ginned Dec. 1-Jan. 15, crop of—					Ginned after Jan. 15, crop of—				
	1932	1931	1930	1929	1928	1932	1931	1930	1929	1928
	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales
All grades ²										
All grades ²	12.5	31.9	38.3	29.8	27.7	6.0	18.3	9.9	5.7	14.1
Extra White.....	7.9	30.2	25.6	16.8	23.9	5.5	6.8	5.7	3.8	8.7
Above 3-G.M.....										
3-G.M.	.2	.1		.4	.1					
4-S.M.	2.8	6.3	4.7	12.0	14.2	.2	.3	.1	.8	4.1
5-M.	4.1	15.5	13.9	3.7	8.7	1.6	1.8	1.2	1.2	4.1
6-S.L.M.	.7	7.4	6.2	.7	.8	2.7	4.0	1.7	1.8	.5
7-L.M.			.5		.1	.9	.6	.5		
Below 7-L.M. ⁴1	.2	.3			.1	.1	2.2		
White.....			.2					7.1		.5
1-M.F.										
2-S.G.M.										
3-G.M.										
4-S.M.										
5-M.										
6-S.L.M.			.1							
7-L.M.										
8-S.G.O. ⁴										
9-G.O. ⁴										

See footnotes at end of table.

TABLE 5.—*Grade of American upland cotton: Quantities ginned in the United States during specified periods,¹ by States, crops of 1932, 1931, 1930, 1929, and 1928—Continued*

ARIZONA—Continued

Grade	Ginned Dec. 1-Jan. 15, crop of—					Ginned after Jan. 15, crop of—				
	1932	1931	1930	1929	1928	1932	1931	1930	1929	1928
	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales
Spotted.....	4.6	1.5	12.6	12.0	3.7	0.2	1.2	4.2	1.3	5.1
3-G.M.....										.3
4-S.M.....	3.3	1.1	5.7	4.2	2.5	.1	.5	.2	.3	1.1
5-M.....	1.2	.3	4.6	5.5	.9	.1	.5	1.5	.7	1.9
6-S.L.M. ⁴1	.1	1.7	1.9	.2		.1	2.2	.3	1.8
7-L.M. ⁴6	.4				.3		
Yellow Tinged.....				.1				.1		.1
2-S.G.M.....										
3-G.M.....										
4-S.M.....										
5-M ⁴										
6-S.I.M. ⁴1				.1		
7-L.M. ⁴										
Gray.....					.2					
3-G.M.....										
4-S.M.....					.1					
5-M ⁴1					
No grade ^{4,6}8			.3	3.1	
										.1

ARKANSAS

Grade	Total, crop of—					Ginned prior to Oct. 1				
	1932	1931	1930	1929	1928	1932	1931	1930	1929	1928
	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales
All grades ²	1,283.4	1,836.1	863.5	1,395.9	1,216.2	501.8	268.6	264.2	530.0	361.8
Extra White.....		2.0			1.7	.2	.1			.1
Above 3-G.M.....										
3-G.M.....					.1					.1
4-S.M.....					1.4	.1				1.0
5-M.....	.6				.2					.1
6-S.L.M.....	1.8					.1	.1			
7-L.M.....	.5									
Below 7-L.M. ⁴										
White.....	1,210.9	1,782.9	817.1	1,313.2	1,124.1	488.3	263.8	255.9	530.7	361.0
1-M.F.....						.1				.1
2-S.G.M.....		.2	.3	1.3	1.2		.1	.2	1.1	.8
3-G.M.....	12.7	69.5	63.2	198.4	116.5	10.2	38.9	48.0	161.1	87.4
4-S.M.....	337.9	600.6	256.0	535.6	428.3	208.2	181.0	121.7	277.7	217.9
5-M.....	549.8	592.0	362.1	387.7	338.0	246.9	40.6	81.0	81.9	52.0
6-S.L.M.....	212.6	242.1	111.5	103.2	138.8	21.0	2.4	4.9	8.3	2.0
7-L.M.....	50.1	102.7	26.1	47.5	37.3	1.1	.2	.1	.5	.1
8-S.G.O. ⁴	29.8	112.8	5.6	29.7	34.8					
9-G.O. ⁴	18.0	62.1	1.4	9.8	19.1					
Spotted.....	60.0	87.1	44.0	72.1	83.4	13.1	4.7	8.3	7.1	.6
3-G.M.....		1.9	3.5	5.3	2.4	1.9	.9	.9	2.1	.9
4-S.M.....	20.9	15.6	19.4	22.4	20.2	9.5	3.1	5.1	4.7	.4
5-M.....	14.5	8.9	13.7	25.9	30.8	2.6	.6	1.1	1.3	.1
6-S.L.M. ⁴	5.7	5.2	4.5	13.8	15.4	.1	.1			
7-L.M. ⁴	8.0	3.9	1.1	7.6	9.1					

See footnotes at end of table.

TABLE 5.—*Grade of American upland cotton: Quantities ginned in the United States during specified periods,¹ by States, crops of 1932, 1931, 1930, 1929, and 1928—Continued*

ARKANSAS—Continued

Grade	Total, crop of—					Ginned prior to Oct. 1				
	1932	1931	1930	1929	1928	1932	1931	1930	1929	1928
	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales
Yellow Tinged.....	0.9	0.4	1.3	4.9	1.7	0.1	0.1	0.1	0.1	0.1
2-S.G.M.....										
3-G.M.....										
4-S.M.....	.1	.1	.2	.4	.5	.1	.1	.1	.1	.1
5-M. ⁴2	.1	.4	1.4	.5					
6-S.L.M. ⁴2	.1	.6	1.3	.4					
7-L.M. ⁴4	.1	.1	1.8	.3					
Light Yellow Stained.....	.1			.2	.1	0.1				
3-G.M.....				.1	.1	.1				
4-S.M. ⁴										
5-M. ⁴1						
Yellow Stained.....			.1		.1					
3-G.M.....						.1				
4-S.M. ⁴										
5-M. ⁴1							
Gray.....	.5	.5	.1	.3	2.6					
3-G.M.....			.1							
4-S.M.....	.4	.3	.1		1.8					
5-M. ⁴1	.1		.3	.8					
Blue Stained.....		.1		.1	.1					
3-G.M. ⁴1						
4-S.M. ⁴1								
5-M. ⁴1					
No grade ⁴	8.1	15.1	.9	3.4	3.9	.2				
Grade	Ginned during October					Ginned during November				
	1932	1931	1930	1929	1928	1932	1931	1930	1929	1928
	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales
All grades ⁴	481.0	766.1	368.6	523.4	399.3	181.0	438.3	182.3	189.1	259.1
Extra White.....	2.5			.3	.1	.3			.3	
Above 3-G.M.....										
3-G.M.....										
4-S.M.....				.2					.2	
5-M.....	.6			.1					.1	
6-S.L.M.....	1.5				.1	.2				
7-L.M.....	.4				.1					
Below 7-L.M. ⁴										
White.....	461.7	749.1	353.4	510.5	383.6	170.5	433.1	188.6	166.5	239.4
1-M.F.....										
2-S.G.M.....		.1		.1	.3				.1	.1
3-G.M.....	1.0	27.4	10.4	31.7	24.2	.6	2.4	4.5	4.7	3.8
4-S.M.....	101.0	348.6	96.1	222.0	171.2	25.2	56.5	34.7	31.1	40.6
5-M.....	228.3	314.9	192.1	224.8	160.0	63.9	216.0	70.5	66.8	105.7
6-S.L.M.....	119.8	55.6	51.9	30.9	26.7	51.8	140.2	43.2	42.4	78.7
7-L.M.....	10.2	2.3	2.0	1.0	1.0	20.2	16.4	13.2	10.1	8.6
8-S.G.O. ⁴4	.2			.1	8.2	1.5	2.2	4.8	1.7
9-G.O. ⁴1					.6	.1	.2	.6	.2

See footnotes at end of table.

TABLE 5.—*Grade of American upland cotton: Quantities ginned in the United States during specified periods,¹ by States, crops of 1932, 1931, 1930, 1929, and 1928—Continued*

ARKANSAS—Continued

Grade	Ginned during October					Ginned during November				
	1932	1931	1930	1929	1928	1932	1931	1930	1929	1928
	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales
Spotted.....	16.7	17.0	14.9	12.5	15.6	9.7	4.9	13.2	21.5	17.2
3-G.M.....	.8	2.2	2.3	1.3	1.1	1	.4	.9	.1	.5
4-S.M.....	11.2	10.4	7.6	8.0	9.8	6.1	1.6	5.5	7.6	6.8
5-M.....	4.8	3.8	4.6	3.1	4.2	2.7	2.3	5.2	10.7	8.2
6-S.L.M. ⁴4	.5	.4	.1	.4	.6	.6	1.4	2.5	1.5
7-L.M. ⁴1			.1	.2		.2	.0	.2
Yellow Tinged.....	.1		.2			.1		.3	.4	.5
2-S.G.M.....										
3-G.M.....										
4-S.M.....	.1		.1					.1	.2	.2
5-M ⁴1			.1		.1	.2	.2
6-S.L.M. ⁴1		
7-L.M. ⁴										
Light Yellow Stained.....										.1
3-G.M.....										.1
4-S.M. ⁴										
5-M ⁴										
Yellow Stained.....						.1				.1
3-G.M.....						.1				
4-S.M. ⁴										
5-M ⁴1		
Gray.....			.1			.2	.1		.2	1.9
3-G.M.....										
4-S.M.....			.1			.1	.1			1.5
5-M ⁴1			.2	.4
Blue Stained.....				.1				.1		
3-G.M. ⁴1				.1		
4-S.M. ⁴										
5-M ⁴										
No grade ^{4,6}2	.1	.1	.1	.1
Grade	Ginned Dec. 1-Jan. 15, crop of—					Ginned after Jan. 15, crop of—				
	1932	1931	1930	1929	1928	1932	1931	1930	1929	1928
	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales
All grades ²	89.2	162.9	43.9	99.2	136.0	30.4	200.2	4.5	45.2	60.0
Extra White.....										
Above 3-G.M.....										
3-G.M.....										
4-S.M.....										
5-M.....										
6-S.L.M.....										
7-L.M.....										
Below 7-L.M. ⁴										

See footnotes at end of table.

TABLE 5.—*Grade of American upland cotton: Quantities ginned in the United States during specified periods,¹ by States, crops of 1932, 1931, 1930, 1929, and 1928—Continued*

ARKANSAS—Continued

Grade	Ginned Dec. 1-Jan. 15, crop of—					Ginned after Jan. 15, crop of—				
	1932	1931	1930	1929	1928	1932	1931	1930	1929	1928
	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales
White	72.5	156.8	35.8	75.5	96.1	17.9	180.1	3.4	30.0	44.1
1-M.F.										
2-S.G.M.										
3-G.M.		.6	.3	.8	1.1					.1
4-S.M.		2.9	10.8	3.9	4.1	8.2	.6	3.1	.5	.7
5-M.		9.4	16.8	7.8	12.8	17.8	1.3	4.6	.7	1.9
6-S.L.M.		17.9	32.2	10.6	18.4	24.5	1.2	11.7	.9	3.2
7-L.M.		16.5	45.3	9.1	22.7	19.1	2.1	38.5	.8	7.2
8-S.G.O. ⁴		16.6	39.0	3.1	13.4	18.2	4.6	72.1	.3	11.4
9-G.O. ⁴		9.2	12.1	1.0	3.3	7.2	8.1	49.9	.2	6.0
Spotted		12.4	3.8	6.5	20.4	36.7	8.1	6.7	1.1	10.6
3-G.M.		.1			.1	.2				
4-S.M.		3.0	.5	1.0	1.8	8.5	.1	2	.3	.7
5-M.		4.6	1.1	2.2	8.7	14.0	.3	1.1	.6	2.1
6-S.L.M. ⁴		2.8	1.4	2.4	6.8	7.8	1.8	2.6	.3	4.2
7-L.M. ⁴		1.9	.8	.9	3.0	6.2	5.9	3.0		4.0
Yellow Tinged		.2	.1	.8	2.3	.9	.5	.2		2.1
2-S.G.M.										
3-G.M.										
4-S.M.										.1
5-M. ⁴										3
6-S.L.M. ⁴										.6
7-L.M. ⁴										.1
Light Yellow Stained										.1
3-G.M.										
4-S.M. ⁴										
5-M. ⁴										.1
Yellow Stained										
3-G.M.										
4-S.M. ⁴										
5-M. ⁴										
Gray		.3	.4		.1	.3				.4
3-G.M.										
4-S.M. ⁴										
5-M. ⁴										
Blue Stained										.1
3-G.M. ⁴										
4-S.M. ⁴										
5-M. ⁴										
No grade ^{4,6}		3.8	1.8	.8	.0	1.9	3.9	13.2		2.4
										1.9

CALIFORNIA

Grade	Total, crop of—					Ginned prior to Oct. 1				
	1932	1931	1930	1929	1928	1932	1931	1930	1929	1928
	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales
All grades ²	124.4	171.2	256.3	254.1	171.0	7.4	27.8	15.4	13.5	20.3
Extra White	119.2	151.1	254.9	219.1	159.1	7.4	27.8	15.4	8.3	20.0
Above 3-G.M.	1.1	.4	.2							
3-G.M.	71.6	52.4	59.9	112.9	77.5	6.3	23.6	8.9	7.2	18.0
4-S.M.	37.9	73.0	107.7	66.0	42.6	1.0	3.7	6.4	1.1	1.4
5-M.	5.7	20.8	56.8	24.8	28.5	.1	.1	.1		.5
6-S.L.M.	1.6	2.2	16.9	13.5	8.5					.1
7-L.M.	.5	.6	2.9	1.9	2.0					
Below 7-L.M. ⁴	.8	.8	10.5							

See footnotes at end of table.

34 STATISTICAL BULLETIN 47, U. S. DEPT. OF AGRICULTURE

TABLE 5.—*Grade of American upland cotton: Quantities ginned in the United States during specified periods,¹ by States, crops of 1932, 1931, 1930, 1929, and 1928—Continued*

CALIFORNIA—Continued

Grade	Total, crop of—					Ginned prior to Oct. 1				
	1932	1931	1930	1929	1928	1932	1931	1930	1929	1928
	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales
White.....	.2	9.6	23.7	8.3						
1-M.F.										
2-S.G.M.			23.4							5.1
3-G.M.			.2							.1
4-S.M.		.8								.3
5-M.		2.7								
6-S.L.M.		3.4								
7-L.M.		.1	2.1							
8-S.G.O. ⁴			.5							
9-G.O. ⁴			.1							
Spotted.....		2.5	2.0	1.4	5.1	3.3				
3-G.M.		.1								
4-S.M.		1.3	1.5	.8	1.5	1.6				
5-M.		1.0	.4	.4	2.7	.7				
6-S.L.M.		.1	.1	.2	.8	.3				
7-L.M. ⁴					.1					
Yellow Tinged.....										.1
2-S.G.M.										
3-G.M.										
4-S.M.										
5-M. ⁴										
6-S.L.M.										
7-L.M. ⁴										
Gray.....		.2	.5		.1					
3-G.M.										
4-S.M.			.3							
5-M. ⁴			.2		.1					
No grade ^{4,6}		2.3	8.0		6.1	.2				
Grade	Ginned during October					Ginned during November				
	1932	1931	1930	1929	1928	1932	1931	1930	1929	1928
	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales
All grades ²	53.6	66.6	75.3	87.4	61.3	45.4	38.9	72.3	78.0	43.2
Extra White.....	53.0	66.6	75.3	69.4	60.2	44.3	38.5	72.3	76.8	43.0
Above 3-G.M.										
3-G.M.	1.1		.1							
4-S.M.	42.1	28.1	32.7	64.9	49.2	22.2	.5	17.0	38.9	10.0
5-M.	9.2	34.2	41.3	4.0	10.4	21.2	29.1	43.3	35.0	18.8
6-S.L.M.	.3	3.9	1.2	.2	.5	.8	8.9	11.7	2.4	12.3
7-L.M.	.3	.2		.3		.1		.2	.5	1.7
Below 7-L.M. ⁴										.2
White.....		.1								.5
1-M.F.										
2-S.G.M.				17.8						.5
3-G.M.				.1						
4-S.M.										
5-M.										
6-S.L.M.										
7-L.M.										
8-S.G.O. ⁴										
9-G.O. ⁴										

See footnotes at end of table.

TABLE 5.—*Grade of American upland cotton: Quantities ginned in the United States during specified periods,¹ by States, crops of 1932, 1931, 1930, 1929, and 1928—Continued*

CALIFORNIA—Continued

Grade	Ginned during October					Ginned during November				
	1932	1931	1930	1929	1928	1932	1931	1930	1929	1928
Spotted.....										
	<i>1,000 bales</i>	<i>1,000 bales</i>	<i>1,000 bales</i>	<i>1,000 bales</i>	<i>1,000 bales</i>	<i>1,000 bales</i>	<i>1,000 bales</i>	<i>1,000 bales</i>	<i>1,000 bales</i>	<i>1,000 bales</i>
Spotted.....	.5					1.1	0.3			
3-G.M.....						.1	.1			
4-S.M.....						.1	.6	.2		.3
5-M.....		.5					.3	.1		.3
6-S.L.M. ⁴1			.1			.1
7-L.M. ⁴										
Yellow Tinged.....										
2-S.G.M.....										
3-G.M.....										
4-S.M.....										
5-M. ⁴										
6-S.L.M. ⁴										
7-L.M. ⁴										
Gray.....									.1	
3-G.M.....										
4-S.M.....										
5-M. ⁴1		
No grade ^{4,6}1				
 Grade										
Ginned Dec. 1-Jan. 15, crop of—										
Grade	1932	1931	1930	1929	1928	1932	1931	1930	1929	1928
All grades ²										
	<i>1,000 bales</i>	<i>1,000 bales</i>	<i>1,000 bales</i>	<i>1,000 bales</i>	<i>1,000 bales</i>	<i>1,000 bales</i>	<i>1,000 bales</i>	<i>1,000 bales</i>	<i>1,000 bales</i>	<i>1,000 bales</i>
All grades ²	13.2	23.5	70.2	49.3	24.8	4.8	14.4	23.1	25.9	21.4
Extra White.....	11.7	17.0	69.3	46.9	22.9	2.8	1.2	22.6	17.7	13.0
Above 3-G.M.....										
3-G.M.....	1.0	.2	1.3	1.9	.2					.1
4-S.M.....	6.4	6.8	16.5	24.5	8.0	.1	.1	.2	1.4	4.0
5-M.....	3.7	7.4	40.9	15.5	10.9	.8	.5	2.9	6.7	4.3
6-S.L.M.....	.5	1.7	9.4	4.1	3.3	.7	.3	7.3	8.6	3.4
7-L.M.....	.1	.4	.6	.9	.5	.4	.2	2.3	1.0	1.2
Below 7-L.M. ⁴5	.6			.8	.1	9.9		
White.....		5.1				.8	.1	4.5		.1
1-M.F.....										6.4
2-S.G.M.....										
3-G.M.....										
4-S.M.....										
5-M.....			2.2					.5		
6-S.L.M.....			1.6					1.8		.1
7-L.M.....			.4					1.7		.6
8-S.G.O. ⁴1				.5		.4		5.5
9-G.O. ⁴3	.1			.2
Spotted.....	.8	1.0	.9	2.1	1.1	.1	.7	.5	2.2	1.8
3-G.M.....										.3
4-S.M.....	.7	.8	.6	.9	.6		.5	.2	.3	.8
5-M.....	.1	.2	.2	1.0	.2	.1	.1	.2	1.3	.4
6-S.L.M. ⁴1				.1	.1		.6
7-L.M. ⁴1					.3
Yellow Tinged.....										.1
2-S.G.M.....										
3-G.M.....										
4-S.M.....										
5-M. ⁴										
6-S.L.M. ⁴										
7-L.M. ⁴										

See footnotes at end of table.

TABLE 5.—*Grade of American upland cotton: Quantities ginned in the United States during specified periods,¹ by States, crops of 1932, 1931, 1930, 1929, and 1928—Continued*

CALIFORNIA—Continued

Grade	Ginned Dec. 1-Jan. 15, crop of—					Ginned after Jan. 15, crop of—				
	1932	1931	1930	1929	1928	1932	1931	1930	1929	1928
	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales
Gray.....	.2	.3	—	—	—	—	.1	—	—	—
3-G.M.....	—	—	—	—	—	—	—	—	—	—
4-S.M.....	—	.3	—	—	—	—	—	—	—	—
5-M. ⁴2	—	—	—	—	—	.1	—	—	.1
No grade ^{4,6}5	.1	—	0.3	—	1.8	7.9	—	5.8	0.1

FLORIDA

Grade	Total, crop of—					Ginned prior to Oct. 1				
	1932	1931	1930	1929	1928	1932	1931	1930	1929	1928
	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales
All grades ²	15.6	43.4	51.1	29.8	20.1	10.9	32.9	40.3	24.9	11.2
White.....	10.3	31.0	48.6	22.2	12.5	7.5	22.0	39.5	18.9	8.7
1-M.F.....	—	—	—	—	—	—	—	—	—	—
2-S.G.M.....	—	—	—	—	—	—	—	—	—	—
3-G.M.....	—	.1	1.7	.1	.1	—	—	—	1.6	.1
4-S.M.....	2.8	6.2	23.9	7.1	2.9	1.7	3.5	22.1	6.6	2.6
5-M.....	6.1	20.0	18.3	12.1	5.7	4.7	14.1	13.8	10.1	4.3
6-S.L.M.....	1.4	4.6	3.3	2.6	3.2	1.1	4.3	1.7	1.9	1.6
7-L.M.....	—	.1	.9	.3	.8	—	.1	.2	.2	.1
8-S.G.O. ⁴	—	—	.4	—	—	—	—	.1	—	—
9-G.O. ⁴	—	—	.1	—	—	—	—	—	—	—
Spotted.....	5.3	12.4	2.5	7.6	7.5	3.4	10.9	.8	6.0	2.5
3-G.M.....	.2	.4	.1	.1	.2	.1	.2	.1	.1	.1
4-S.M.....	2.7	6.3	1.3	5.3	3.3	1.7	5.3	.4	4.4	.9
5-M.....	2.3	5.1	1.0	2.1	3.1	1.5	4.8	.3	1.4	1.2
6-S.L.M. ⁴1	.6	.1	.1	.8	.1	.6	—	.1	.3
7-L.M. ⁴	—	—	—	—	.1	—	—	—	—	—
No grade ^{4,6}	—	—	—	—	.1	—	—	—	—	—

Grade	Ginned during October					Ginned during November				
	1932	1931	1930	1929	1928	1932	1931	1930	1929	1928
	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales
All grades ²	3.4	8.6	8.7	4.0	6.6	0.9	1.5	1.5	0.8	1.9
White.....	2.1	7.3	8.0	2.7	3.6	.5	1.4	.7	.5	.2
1-M.F.....	—	—	—	—	—	—	—	—	—	—
2-S.G.M.....	—	—	—	—	—	—	—	—	—	—
3-G.M.....	—	.1	.1	—	—	—	—	—	—	—
4-S.M.....	.7	2.1	1.6	.4	.3	.2	.4	.2	.1	—
5-M.....	1.1	4.9	3.8	1.6	1.3	.3	.9	.4	.3	.1
6-S.L.M.....	.3	.2	1.4	.6	1.5	—	.1	.1	.1	.1
7-L.M.....	—	—	.7	.1	.5	—	—	—	—	—
8-S.G.O. ⁴	—	—	.3	—	—	—	—	—	—	—
9-G.O. ⁴	—	—	.1	—	—	—	—	—	—	—
Spotted.....	1.3	1.3	.7	1.3	2.8	.4	.1	.8	.3	1.7
3-G.M.....	.1	.1	—	—	—	.1	—	.5	.2	.0
4-S.M.....	.8	.9	.3	.7	1.4	.1	.1	.5	.2	.0
5-M.....	.4	.3	.3	.6	1.0	.3	—	.3	.1	.0
6-S.L.M. ⁴	—	—	.1	—	.2	—	—	—	—	.2
7-L.M. ⁴	—	—	—	—	.1	—	—	—	—	—
No grade ^{4,6}	—	—	—	—	.1	—	—	—	—	—

See footnotes at end of table.

TABLE 5.—*Grade of American upland cotton: Quantities ginned in the United States during specified periods,¹ by States, crops of 1932, 1931, 1930, 1929, and 1928—Continued*

FLORIDA—Continued

Grade	Ginned Dec. 1-Jan. 15, crop of—					Ginned after Jan. 15, crop of—				
	1932	1931	1930	1929	1928	1932	1931	1930	1929	1928
	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales
All grades ²2	.2	.5	.1	.3	.2	.2	.2	.1	.2
White.....										
1-M.F.										
2-S.G.M.										
3-G.M.										
4-S.M.										
5-M.										
6-S.L.M.										
7-L.M.										
8-S.G.O. ⁴										
9-G.O. ⁴										
Spotted.....	.2	.2		.3			.1			.2
3-G.M.										
4-S.M.										
5-M.										.2
6-S.L.M. ⁴										
7-L.M. ⁴										
No grade ⁴										

GEORGIA

Grade	Total, crop of—					Ginned prior to Oct. 1				
	1932	1931	1930	1929	1928	1932	1931	1930	1929	1928
	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales
All grades ²	861.8	1,393.7	1,507.5	1,339.8	1,053.2	426.2	649.4	842.2	678.2	308.9
Extra White.....	1.6	.1	.2		.4	1.0	.1	.2		.8
Above 3-G.M.										
3-G.M.										
4-S.M.										.1
5-M.		.5					.1			.1
6-S.L.M.		1.0	.1			.2	.7	.1		.1
7-L.M.		.1								
Below 7-L.M. ⁴										
White.....	702.9	1,325.7	1,276.6	1,048.1	836.2	382.2	619.0	764.6	497.6	248.7
1-M.F.										
2-S.G.M.										.4
3-G.M.	15.3	108.7	47.0	26.1	63.5	9.3	47.8	42.0	16.8	11.3
4-S.M.	228.2	512.1	471.8	280.4	308.3	128.7	219.7	365.0	193.1	67.1
5-M.	347.4	547.0	470.1	377.9	236.0	186.8	289.9	281.1	225.3	104.5
6-S.L.M.	95.7	143.1	189.2	189.3	153.9	49.6	57.1	66.0	57.1	57.1
7-L.M.	13.6	10.2	78.7	122.3	53.0	6.4	1.4	9.0	4.4	7.5
8-S.G.O. ⁴	2.1	.4	16.1	41.0	15.8	1.2	.1	.7	.5	.8
9-G.O. ⁴	.5		2.6	10.1	3.8	.2				.1
Spotted.....	166.0	66.1	295.1	255.5	208.9	42.8	30.0	70.1	78.8	58.9
3-G.M.	12.2	13.4	35.8	5.0	20.0	2.7	4.0	11.3	3.1	4.5
4-S.M.	77.3	39.5	151.8	111.5	121.4	21.6	20.7	37.7	55.9	32.3
5-M.	57.9	12.0	81.3	97.7	58.6	15.7	5.2	19.2	18.6	19.8
6-S.L.M. ⁴	8.1	1.0	20.9	36.1	8.4	2.6	.1	1.8	1.1	2.2
7-L.M. ⁴	.5	.2	5.3	5.2	.5	.2		.1	.1	.1
Yellow Tinged.....	.8	1.1	23.1	20.1	4.8	.1	.3	6.9	1.7	1.0
2-S.G.M.										
3-G.M.										
4-S.M.		.6	10.9	5.7	2.4	.1	.3	1.7	.3	.8
5-M.		.1	.1	5.2	7.9	.9			1.3	.3
6-S.L.M. ⁴				2.5	4.3	.4			.2	.1
7-L.M. ⁴				.5	1.4				.1	

See footnotes at end of table.

TABLE 5.—*Grade of American upland cotton: Quantities ginned in the United States during specified periods,¹ by States, crops of 1932, 1931, 1930, 1929, and 1928—Continued*

GEORGIA—Continued

Grade	Total, crop of—					Ginned prior to Oct. 1				
	1932	1931	1930	1929	1928	1932	1931	1930	1929	1928
	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales
Light Yellow Stained.....										
3-G.M.....										
4-S.M. ⁴1	.3	.2						
5-M. ⁴4	.6	.1				.1	
Yellow Stained.....				.2	.4					.1
3-G.M.....										
4-S.M.1	.1						.1
5-M. ⁴1	.3						
Gray.....	0.4	.6		7.7	.9	0.1				
3-G.M.....	.1			.1	.4					
4-S.M.2	.2		3.9	.5	.1				
5-M. ⁴1	.4		3.7						
Blue Stained.....				1.0	.1					
3-G.M. ⁴										
4-S.M. ⁴7	.1					
5-M. ⁴3						
No grade ^{4,6}1		1.6	6.2	1.8					.1
 Ginned during October										
Grade	1932	1931	1930	1929	1928	1932	1931	1930	1929	1928
	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales
	275.0	529.0	531.3	452.8	475.5	114.4	166.2	139.2	146.0	186.4
All grades ²										
Extra White.....	.3					.1	.2			
Above 3-G.M.....										
3-G.M.....										
4-S.M.										
5-M.....	.2									
6-S.L.M.1					.1	.1			
7-L.M.1			
Below 7-L.M. ⁴										
White.....	219.4	507.7	366.6	318.5	400.6	82.5	157.5	102.9	107.6	140.6
1-M. F.....						.1				
2-S.G.M.1	1.2	.1	.1	1.0					.3
3-G.M.	4.9	50.3	4.1	6.9	39.8	1.0	7.5	.5	2.1	0.7
4-S.M.	71.6	209.1	87.8	66.7	157.9	23.5	67.7	15.4	14.0	65.0
5-M.....	105.5	173.4	135.4	107.0	80.2	45.2	67.7	41.1	32.2	38.7
6-S.L.M.	30.5	66.6	87.0	80.5	69.9	11.7	13.7	26.8	28.9	19.1
7-L.M.	5.8	6.9	45.0	46.6	35.4	.9	.9	14.3	22.2	6.1
8-S.G.O.8	.2	6.6	9.5	13.1	.1		4.0	7.1	1.2
9-G.O.2		.6	1.3	3.2	.1		.8	1.1	.5
Spotted.....	54.6	20.7	153.3	120.7	72.3	31.3	8.3	35.0	32.8	44.6
3-G.M.....	6.0	6.5	21.7	1.7	8.9	1.6	1.9	1.9	.1	3.7
4-S.M.	31.2	10.8	86.9	44.1	41.1	15.5	4.8	19.4	9.1	28.4
5-M.....	15.7	3.1	36.8	56.3	18.4	13.0	1.5	11.0	14.6	11.8
6-S.L.M. ⁴	1.6	.1	5.7	17.6	3.7	1.2	.1	2.0	8.0	.7
7-L.M. ⁴1	2	2.2	1.0	.2			.7	1.0	
Yellow Tinged.....	.4	.2	10.9	11.2	1.0	.3	.3	.8	4.1	.6
2-S.G.M.1	.1	2.0	.5	.1					.1
3-G.M.2		6.1	3.5	.6	.3	.2	.4	.9	.2
4-S.M.1	.1	2.0	5.0	.2			.1	1.9	
5-M. ⁴7	1.9	.1				1.2	
6-S.L.M. ⁴1	.3				.2	.1	
7-L.M. ⁴										

See footnotes at end of table.

TABLE 5.—*Grade of American upland cotton: Quantities ginned in the United States during specified periods,¹ by States, crops of 1932, 1931, 1930, 1929, and 1928—Continued*

GEORGIA—Continued

Grade	Ginned during October					Ginned during November				
	1932	1931	1930	1929	1928	1932	1931	1930	1929	1928
	1,000 bales	1,000 bales	1,000 bales 0.3	1,000 bales 0.4	1,000 bales 0.1	1,000 bales	1,000 bales	1,000 bales	1,000 bales 0.2	1,000 bales
Light Yellow Stained.....										
3-G.M.....										
4-S.M. ⁴1	.2					
5-M. ⁴2	.2					.2
Yellow Stained.....									0.1	.1
3-G.M.....										
4-S.M. ⁴										
5-M. ⁴1	.1
Gray.....	0.3	0.4			1.5			0.1		.7
3-G.M. ⁴1
4-S.M. ⁴6
5-M. ⁴2
Blue Stained.....										
3-G.M. ⁴										
4-S.M. ⁴										
5-M. ⁴										
No grade ^{4,6}2	.3	1.4	0.1		.4	.5
									.3	

Grade	Ginned Dec. 1-Jan. 15, crop of—					Ginned after Jan. 15, crop of—				
	1932	1931	1930	1929	1928	1932	1931	1930	1929	1928
	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales
All grades ²	37.2	35.0	66.9	129.6	67.2	9.0	13.2	17.9	33.2	15.2
Extra White.....	.1									
Above 3-G.M.....										
3-G.M.....										
4-S.M.....										
5-M.....										
6-S.L.M.....	.1									
7-L.M.....										
Below 7-L.M. ⁴										
White.....	15.3	30.7	35.3	99.8	37.9	3.5	10.8	7.0	24.6	8.4
1-M.P.....										
2-S.G.M.....										
3-G.M.....										
4-S.M.....	3.2	12.0	3.4	5.7	16.9	1.2	3.6	.2	.9	1.4
5-M.....	8.8	12.1	11.3	11.3	10.3	1.1	3.9	1.2	2.1	2.3
6-S.L.M.....	3.1	3.5	7.9	18.9	5.8	.8	2.2	1.5	4.0	2.0
7-L.M.....	.2	.5	8.8	37.4	2.2	.3	.5	1.6	11.7	1.8
8-S.G.O. ⁴										
9-G.O. ⁴										
Spotted.....	21.8	4.9	28.2	16.8	27.3	5.5	2.2	8.5	6.4	5.8
3-G.M.....	1.4	.9	.9	.1	2.7	.5	.1			.2
4-S.M.....	7.7	2.7	6.9	1.9	17.7	1.3	.5	.9	.5	1.9
5-M.....	11.4	1.1	11.6	5.8	6.4	2.1	1.1	2.7	2.4	2.2
6-S.L.M. ⁴	1.2	.2	7.7	6.7	.5	1.5	.5	3.7	2.7	1.3
7-L.M. ⁴1		1.1	2.3		.1		1.2	.8	.2
Yellow Tinged.....		.1	2.5	2.2	1.4		.2	2.0	.9	.8
2-S.G.M.....										
3-G.M.....		.1	.1							
4-S.M.....										
5-M. ⁴										
6-S.L.M.....										
7-L.M. ⁴										

See footnotes at end of table.

TABLE 5.—Grade of American upland cotton: Quantities ginned in the United States during specified periods,¹ by States, crops of 1932, 1931, 1930, 1929, and 1928—Continued

GEORGIA—Continued

Grade	Ginned Dec. 1-Jan. 15, crop of—					Ginned after Jan. 15, crop of—				
	1932	1931	1930	1929	1928	1932	1931	1930	1929	1928
	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales
Light Yellow Stained...										
3-G.M.										
4-S.M. ⁴		.1	.1						.1	
5-M. ⁴			.1							.2
Yellow Stained...					0.1					
3-G.M.										
4-S.M. ⁴										
5-M. ⁴				.1						
Gray...						5.1	0.5			
3-G.M.							.3			
4-S.M.						1.9	.2			
5-M. ⁴						3.2				.4
Blue Stained...						.9				.1
3-G.M. ⁴										.1
4-S.M. ⁴						.6				.1
5-M. ⁴						.3				
No grade ^{4,6}7	4.7	.1		.3 .0

LOUISIANA

Grade	Total, crop of—					Ginned prior to Oct. 1				
	1932	1931	1930	1929	1928	1932	1931	1930	1929	1928
	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales
All grades ² ...	599.5	876.6	704.8	797.7	685.9	393.9	300.1	399.6	542.4	370.3
Extra White...	15.1	1.6				.1	8.7			.1
Above 3-G.M.										
3-G.M.	.1						.1			
4-S.M.	5.8						4.1			.1
5-M.	6.1	.3					3.7			
6-S.L.M.	2.8	.9					.8			
7-L.M.	.8	.4								
Below 7-L.M. ⁴										
White...	548.7	840.4	672.7	769.6	658.7	362.2	287.3	384.7	524.0	362.0
1-M.F.										
2-S.G.M.	.4	.3	.6	.4	.2		.2	.6	.1	.1
3-G.M.	31.4	28.3	52.0	41.6	102.1	24.8	20.0	48.4	38.2	76.7
4-S.M.	225.7	263.6	236.0	263.1	279.5	180.9	140.4	197.6	227.9	173.9
5-M.	221.6	385.0	222.9	314.3	180.6	143.7	114.4	113.0	201.9	76.8
6-S.L.M.	68.5	124.9	120.2	129.1	75.8	12.2	11.4	23.2	53.9	24.4
7-L.M.	10.3	21.6	29.5	12.1	15.8	.3	.8	1.7	1.6	7.2
8-S.G.O. ⁴	.7	8.9	8.9	6.6	2.5		.1	.2	.1	.0
9-G.O. ⁴	.1	7.8	2.6	2.3	.1					.1
Spotted...	33.7	28.9	27.9	25.7	25.8	21.0	12.6	14.3	18.2	7.4
3-G.M.	7.6	4.1	3.2	1.4	3.7	6.4	1.9	2.6	.8	.6
4-S.M.	18.7	16.7	12.1	12.2	10.5	13.0	7.9	8.4	9.2	3.5
5-M.	6.7	5.7	9.4	10.1	8.9	2.4	2.2	3.1	7.0	2.5
6-S.L.M. ⁴	.6	1.1	2.7	1.0	2.1	.1	.5	.2	1.1	.8
7-L.M. ⁴	.1	1.3	.6	.1	.6		.1		.1	.1

See footnotes at end of table.

TABLE 5.—*Grade of American upland cotton: Quantities ginned in the United States during specified periods,¹ by States, crops of 1932, 1931, 1930, 1929, and 1928—Continued*

LOUISIANA—Continued

Grade	Total, crop of—					Ginned prior to Oct. 1				
	1932	1931	1930	1929	1928	1932	1931	1930	1929	1928
	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales
Yellow Tinged.....										
2-S.G.M.	.9									
3-G.M.										
4-S.M.										
5-M. ⁴										
6-S.L.M. ⁴										
7-L.M. ⁴										
Light Yellow Stained.....										
3-G.M.										
4-S.M. ⁴										
5-M. ⁴										
Yellow Stained.....										
3-G.M.										
4-S.M. ⁴										
5-M. ⁴										
Gray.....										
3-G.M.	.4	2.3	1.1	1.0	.3					
4-S.M.										
5-M. ⁴										
Blue Stained.....										
3-G.M. ⁴										
4-S.M. ⁴										
5-M. ⁴										
No grade ^{4,0}										
	.7	3.4	2.2	.6	.9					
Grade	Ginned during October					Ginned during November				
	1932	1931	1930	1929	1928	1932	1931	1930	1929	1928
	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales
All grades ²										
156.0	367.5	207.8	202.8	215.9	40.2	146.1	74.7	34.3	77.3	
Extra White.....										
Above 3-G.M.	6.4	1.1								
3-G.M.										
4-S.M.										
5-M.	1.7									
6-S.L.M.	2.4	.1								
7-L.M.	2.0	.7								
Below 7-L.M. ⁴	.3	.3								
White.....										
1-M.F.	143.5	366.3	197.1	197.5	205.1	34.4	139.6	71.6	32.0	71.2
2-S.G.M.										
3-G.M.	.1	.1								
4-S.M.	5.9	4.9	3.0	3.1	23.2	.7	2.9	.6	.3	1.9
5-M.	38.5	87.6	30.8	30.5	84.7	5.2	28.2	6.5	4.6	19.2
6-S.L.M.	66.0	198.8	83.2	100.8	73.2	9.6	57.0	22.2	10.8	25.9
7-L.M.	29.4	58.6	64.1	60.9	22.3	13.3	41.8	28.4	11.0	20.9
8-S.G.O. ⁴	2.7	5.6	13.1	2.1	1.3	5.5	8.4	10.6	3.5	3.1
9-G.O. ⁴	.2	.7	2.3		.3	.1	1.0	3.0	.0	
	.1		.6				.3	.4		
Spotted.....										
6.0	9.0	10.0	5.1	10.5	5.0	4.7	2.7	2.3	5.0	
3-G.M.	.8	2.0	.5	.3	2.6	.3	.2	.1	.3	.6
4-S.M.	3.2	6.0	3.0	1.9	4.4	2.1	2.5	.5	1.1	2.3
5-M.	1.8	1.5	4.9	2.3	3.2	2.2	1.6	.9	.8	2.6
6-S.L.M. ⁴	.1	.1	1.3	.6	.3	.4	.3	1.0	.1	.4
7-L.M. ⁴	.1		.3				.1	.2		

See footnotes at end of table.

TABLE 5.—*Grade of American upland cotton: Quantities ginned in the United States during specified periods,¹ by States, crops of 1932, 1931, 1930, 1929, and 1928—Continued*

LOUISIANA—Continued

Grade	Ginned during October					Ginned during November				
	1932	1931	1930	1929	1928	1932	1931	1930	1929	1928
	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales
Yellow Tinged										
2-S.G.M.										
3-G.M.										
4-S.M.										
5-M. ⁴			.1							
6-S.L.M. ⁴										
7-L.M. ⁴										
Light Yellow Stained					.1					
3-G.M.										
4-S.M. ⁴										
5-M. ⁴				.1						
Yellow Stained										
3-G.M.										
4-S.M. ⁴										
5-M. ⁴										
Gray	0.1	0.2	.5	.1	.1	0.2	1.3	0.2		0.2
3-G.M.										
4-S.M.			.1							
5-M. ⁴										
Blue Stained										
3-G.M. ⁴										
4-S.M. ⁴										
5-M. ⁴										
No grade ^{4,6}			.3	.1		.1	.6	.1	.1	
Grade	Ginned Dec. 1-Jan. 15, crop of—					Ginned after Jan. 15, crop of—				
	1932	1931	1930	1929	1928	1932	1931	1930	1929	1928
	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales
All grades ²	7.7	37.0	19.5	13.9	19.0	1.7	25.9	3.2	4.3	3.4
Extra White			.1							
Above 3-G.M.										
3-G.M.										
4-S.M.										
5-M.			.1							
6-S.L.M.										
7-L.M.										
Below 7-L.M. ⁴										
White	7.1	34.9	16.3	11.8	17.7	1.5	22.3	3.0	4.3	2.7
1-M.F.										
2-S.G.M.										
3-G.M.			.4			.3				
4-S.M.		.7	4.7	1.0	.1	1.6	.4	2.7	.1	.1
5-M.		1.3	9.4	4.2	.6	4.4	.4	5.4	.3	.3
6-S.L.M.		3.3	10.1	3.8	2.1	7.7	.3	3.0	.7	.3
7-L.M.		1.7	5.1	3.5	4.3	2.8	.1	1.7	.6	.6
8-S.G.O. ⁴		.1	3.2	2.6	4.1	.0	.3	3.9	.8	1.5
9-G.O. ⁴			2.0	1.2	.6			5.5	.4	1.7
Spotted		.6	.9	.8	.1	1.3	.2	1.1	.1	
3-G.M.			.1							
4-S.M.		.3	.2	.2		.3	.1	.1		
5-M.		.2	.4	.5		.6	.1			
6-S.L.M. ⁴			.2	.1	.1	.4			.1	
7-L.M. ⁴			.1					1.0		.5

See footnotes at end of table.

TABLE 5.—*Grade of American upland cotton: Quantities ginned in the United States during specified periods,¹ by States, crops of 1932, 1931, 1930, 1929, and 1928—Continued*

LOUISIANA—Continued

Grade	Ginned Dec. 1-Jan. 15, crop of—					Ginned after Jan. 15, crop of—										
	1932		1931		1930	1929		1928	1932		1931		1930	1929		
	<i>1,000 bales</i>		<i>1,000 bales</i>		<i>1,000 bales</i>	<i>1,000 bales</i>		<i>1,000 bales</i>	<i>1,000 bales</i>		<i>1,000 bales</i>		<i>1,000 bales</i>	<i>1,000 bales</i>		
Yellow Tinged.....																
2-S.G.M.....																
3-G.M.....																
4-S.M.....																
5-M. ⁴																
6-S.L.M. ⁴																
7-L.M. ⁴																
Light Yellow Stained.....																
3-G.M.....																
4-S.M. ⁴																
5-M. ⁴																
Yellow Stained.....						0.1										
3-G.M.....																
4-S.M. ⁴1										
5-M. ⁴																
Gray.....		0.3	0.3		.9							0.5				
3-G.M.....																
4-S.M.....					.1											
5-M. ⁴3	.2		.9							.5				
Blue Stained.....					.2	.4										
3-G.M. ⁴																
4-S.M. ⁴1										
5-M. ⁴2	.3										
No grade ⁶8	1.9		.6							2.0	0.1			

MISSISSIPPI

Grade	Total, crop of—					Ginned prior to Oct. 1										
	1932		1931		1930	1929		1928	1932		1931		1930	1929		
	<i>1,000 bales</i>		<i>1,000 bales</i>		<i>1,000 bales</i>	<i>1,000 bales</i>		<i>1,000 bales</i>	<i>1,000 bales</i>		<i>1,000 bales</i>		<i>1,000 bales</i>	<i>1,000 bales</i>		
All grades ²																
1,161.2	1,710.4		1,458.5		1,876.0		1,462.0		511.7	358.5	532.1	908.4	560.3			
Extra White.....		.3				.2		.1						.1		.1
Above 3-G.M.....																
3-G.M.....																.1
4-S.M.....																
5-M.....																
6-S.L.M.....		.2														
7-L.M.....		.1														
Below 7-L.M. ⁴																
White.....	1,021.9	1,677.2	1,416.9	1,700.7	1,400.6	491.5	356.0	518.4	863.4	555.7						
1-M.F.....			.3	.7	.8	2.4					.1			.2	.7	.8
2-S.G.M.....		31.0	103.8	171.5	252.0	231.8	28.0	58.3	121.1	224.3						126.3
3-G.M.....		348.2	803.8	536.2	600.8	646.7	253.8	253.3	274.1	416.2						295.8
4-S.M.....		434.9	503.6	472.2	522.4	364.2	192.6	42.3	111.5	194.0						119.9
5-M.....		165.8	133.9	188.2	199.2	100.0	16.1	1.8	10.2	27.1						11.0
6-S.L.M.....		31.8	47.2	50.7	86.5	33.1	-.7	.2	.8	1.0						1.7
7-L.M.....		7.6	50.9	13.5	37.6	17.0	.1									.2
8-S.G.O. ⁴		2.6	33.7	3.6	10.4	8.4	.1									

See footnotes at end of table.

TABLE 5.—*Grade of American upland cotton: Quantities ginned in the United States during specified periods,¹ by States, crops of 1932, 1931, 1930, 1929, and 1928—Continued*

MISSISSIPPI—Continued

Grade	Total, crop of—					Ginned prior to Oct. 1				
	1932	1931	1930	1929	1928	1932	1931	1930	1929	1928
Spotted	<i>1,000 bales</i> 137.0	<i>1,000 bales</i> 36.1	<i>1,000 bales</i> 39.9	<i>1,000 bales</i> 162.9	<i>1,000 bales</i> 46.7	<i>1,000 bales</i> 19.9	<i>1,000 bales</i> 2.5	<i>1,000 bales</i> 13.7	<i>1,000 bales</i> 44.4	<i>1,000 bales</i> 4.2
3-G.M.	6.8	6.6	3.7	13.4	3.2	2.4	.5	2.3	4.4	.4
4-S.M.	86.1	22.5	17.5	86.4	27.0	14.0	1.2	8.8	26.0	2.3
5-M.	40.3	5.2	13.8	51.0	13.6	3.4	.7	2.3	12.9	1.2
6-S.L.M. ⁴	4.1	1.1	4.3	10.3	2.6	.1	.1	.3	1.1	.2
7-L.M. ⁴	.3	.7	.6	1.8	.3					.1
Yellow Tinged	.5	.1	.5	.4	.3	.1			.2	.1
2-S.G.M.				.1	.1				.1	.1
3-G.M.	.1					.1				
4-S.M.	.3	.1	.2	.2	.1				.1	
5-M. ⁴	.1		.2		.1				.1	
6-S.L.M. ⁴			.1							
7-L.M. ⁴				.1						
Light Yellow Stained	.1		.1	.2	.2				.2	.2
3-G.M.						.1				.1
4-S.M. ⁴				.1	.1				.1	.1
5-M. ⁴	.1		.1	.1					.1	
Gray	.3	.1	.1	.5	3.8					
3-G.M.				.1						
4-S.M.	.2	.1		.1	1.9					
5-M. ⁴	.1		.1	.4	1.9					
Blue Stained				.3	.1					
3-G.M. ⁴										
4-S.M. ⁴					.1					
5-M. ⁴				.3						
No grade ^{4,5}	.5	5.9	1.0	1.8	1.2	.2			.1	

Grade	Ginned during October					Ginned during November				
	1932	1931	1930	1929	1928	1932	1931	1930	1929	1928
All grades ²	<i>1,000 bales</i> 385.2	<i>1,000 bales</i> 781.2	<i>1,000 bales</i> 594.0	<i>1,000 bales</i> 632.8	<i>1,000 bales</i> 539.7	<i>1,000 bales</i> 198.3	<i>1,000 bales</i> 378.2	<i>1,000 bales</i> 251.0	<i>1,000 bales</i> 151.3	<i>1,000 bales</i> 247.0
Extra White	.2			.1		.1				
Above 3-G.M.										
3-G.M.				.1						
4-S.M.										
5-M.										
6-S.L.M.	.1					.1				
7-L.M.	.1									
Below 7-L.M. ⁴										
White	333.3	754.4	580.2	544.7	518.6	145.5	373.7	243.0	136.1	235.2
1-M.F.										
2-S.G.M.		.2	.4	.1	1.4					
3-G.M.	2.0	39.9	33.5	22.3	90.6	.8	4.2	15.7	3.1	12.8
4-S.M.	69.7	447.0	189.1	153.0	268.7	20.1	82.5	62.2	21.2	70.3
5-M.	178.4	239.9	255.4	259.3	139.3	51.1	190.4	88.8	48.2	91.0
6-S.L.M.	75.8	26.6	86.6	99.0	16.8	59.5	81.6	52.4	40.2	50.7
7-L.M.	6.8	12.3	10.6	1.7		12.6	8.6	20.3	19.7	9.2
8-S.G.O.	.5	.8	2.3	.4	.1	1.3	.4	3.2	3.4	1.0
9-G.O. ⁴	.1		.6			.1		.4	.3	

See footnotes at end of table.

TABLE 5.—*Grade of American upland cotton: Quantities ginned in the United States during specified periods,¹ by States, crops of 1932, 1931, 1930, 1929, and 1928—Continued*

MISSISSIPPI—Continued

Grade	Ginned during October					Ginned during November				
	1932	1931	1930	1929	1928	1932	1931	1930	1929	1928
	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales
Spotted	51.3	26.7	13.4	87.9	20.8	52.4	4.5	7.7	15.1	8.7
3-G.M.	3.2	5.6	.9	8.8	2.4	1.1	.5	.5	.2	.3
4-S.M.	33.8	18.4	5.5	53.7	14.5	31.9	2.7	2.4	5.4	4.5
5-M.	13.2	2.6	5.2	24.1	3.6	17.4	1.1	3.8	7.6	3.3
6-S.L.M. ⁴	1.0	.1	1.6	1.2	.3	1.9	.1	.9	1.8	.6
7-L.M. ⁴	.1		.2	.1		.1	.1	.1	.1	
Yellow Tinged		.1	.1			.1	.2		.1	.1
2-S.G.M.										
3-G.M.										
4-S.M.			.1	.1		.1	.2			.1
5-M. ⁴									.1	.1
6-S.L.M. ⁴										
7-L.M. ⁴										
Light Yellow Stained							.1		.1	
3-G.M.										
4-S.M. ⁴										
5-M. ⁴							.1		.1	
Gray		.8		.1	.1		.1			3.0
3-G.M.							.1			
4-S.M.		.2					.1			1.5
5-M. ⁴		.1		.1						1.5
Blue Stained										
3-G.M. ⁴										
4-S.M. ⁴										
5-M. ⁴										
No grade ^{4,6}				.2			.1			.1

Grade	Ginned Dec. 1-Jan. 15, crop of—					Ginned after Jan. 15, crop of—				
	1932	1931	1930	1929	1928	1932	1931	1930	1929	1928
	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales
All grades ²	53.6	80.0	72.4	121.2	81.3	12.4	112.5	9.0	62.3	33.7
Extra White										
Above 3-G.M.										
3-G.M.										
4-S.M.										
5-M.										
6-S.L.M.										
7-L.M.										
Below 7-L.M. ⁴										
White	41.8	87.3	67.4	111.3	69.2	9.8	105.8	7.9	54.2	30.9
1-M.F.										
2-S.G.M.										
3-G.M.	.2	.5	1.4	2.1	2.0	.9	1	.2		.1
4-S.M.	3.4	15.0	8.8	8.2	10.1	1.2	6.0	2.0	2.2	1.8
5-M.	10.4	22.3	14.0	17.3	12.4	2.4	2.7	2.5	3.6	1.6
6-S.L.M.	13.6	20.5	17.4	25.1	23.7	.8	3.4	1.6	7.8	3.8
7-L.M.	10.9	19.0	16.2	40.1	14.9	.8	18.6	1.1	15.1	5.6
8-S.G.O. ⁴	3.0	8.8	7.2	16.3	4.7	2.7	41.7	.5	17.4	11.0
9-G.O. ⁴	.3	1.2	2.3	2.2	1.4	1.9	32.5	.1	7.9	7.0

See footnotes at end of table.

TABLE 5.—*Grade of American upland cotton: Quantities ginned in the United States during specified periods,¹ by States, crops of 1932, 1931, 1930, 1929, and 1928—Continued*

MISSISSIPPI—Continued

Grade	Ginned Dec. 1-Jan. 15, crop of—					Ginned after Jan. 15, crop of—										
	1932		1931		1930	1929		1928	1932		1931		1930	1929		1928
	<i>1,000 bales</i>		<i>1,000 bales</i>		<i>1,000 bales</i>	<i>1,000 bales</i>		<i>1,000 bales</i>	<i>1,000 bales</i>		<i>1,000 bales</i>		1930	<i>1,000 bales</i>		
Spotted.....	11.6		1.4		4.0	0.2		11.3	2.4		1.0		1.1	0.3		1.7
3-G. M.....	.1															
4-S. M.....	5.3		.2		.5		.9		5.3		1.1			.3		.4
5-M.....	5.2		.5		2.0		4.0		4.7		1.1		.3	.5	2.4	.8
6-S. L. M. ⁴9		.5		1.3		3.6		1.1		.2		.3	.2	2.6	.4
7-L. M. ⁴1		.2				.7						.4	.1	.9	.1
Yellow Tinged.....		.1														.1
2-S. G. M.....																
3-G. M.....																
4-S. M.....		.1														
5-M. ⁴																
Light Yellow Stained.....																
3-G. M.....																
4-S. M. ⁴																
5-M. ⁴																
Gray.....		.1							.3		.6					.1
3-G. M.....																
4-S. M.....			.1								.3					
5-M. ⁴3		.3					.1
Blue Stained.....											.1					.2
3-G. M. ⁴																
4-S. M. ⁴1
5-M. ⁴2
No grade ^{4, 6}1		.2		.7		.3		.2		.2		5.7		1.4
																.9

MISSOURI

Grade	Total, crop of—					Ginned prior to Oct. 1										
	1932		1931		1930	1929		1928	1932		1931		1930	1929		1928
	<i>1,000 bales</i>		<i>1,000 bales</i>		<i>1,000 bales</i>	<i>1,000 bales</i>		<i>1,000 bales</i>	<i>1,000 bales</i>		<i>1,000 bales</i>		1930	<i>1,000 bales</i>		
All grades ²	<i>1,000 bales</i> 300.7	<i>1,000 bales</i> 280.4	<i>1,000 bales</i> 153.3	<i>1,000 bales</i> 220.9	<i>1,000 bales</i> 146.9	<i>1,000 bales</i> 80.7	<i>1,000 bales</i> 32.5	<i>1,000 bales</i> 52.9	<i>1,000 bales</i> 23.1	<i>1,000 bales</i> 13.6						
Extra White.....	4.1										.2					
Above 3-G. M.....																
3-G. M.....																
4-S. M.....																
5-M.....		.1														
6-S. L. M.....		3.7														
7-L. M.....		.3														
Below 7-L. M. ⁴																
White.....	279.5	266.0	148.2	191.9	126.5	79.2	32.2	52.5	22.7							13.5
1-M. F.....																
2-S. G. M.....																
3-G. M.....	.4	.8	5.5	1.3	1.6	.4	.7	4.7	1.2							1.4
4-S. M.....	28.0	27.5	37.3	33.1	20.9	24.8	16.5	24.2	11.2							10.2
5-M.....	112.2	76.7	64.1	70.6	29.1	50.1	13.7	22.8	9.1							1.8
6-S. L. M.....	77.2	72.2	20.8	24.6	27.8	3.9	1.2	.8	1.1							1.
7-L. M.....	19.8	23.3	13.2	19.0	7.8											
8-S. G. O. ⁴	21.8	40.2	6.3	30.7	17.2											
9-G. O. ⁴	20.1	25.3	1.0	12.6	22.1											

See footnotes at end of table.

TABLE 5.—*Grade of American upland cotton: Quantities ginned in the United States during specified periods,¹ by States, crops of 1932, 1931, 1930, 1929, and 1928—Continued*

MISSOURI—Continued

Grade	Total, crop of—					Ginned prior to Oct. 1				
	1932	1931	1930	1929	1928	1932	1931	1930	1929	1928
	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales
Spotted.....	7.8	9.2	3.0	18.3	12.8	1.3	0.3	0.4	0.4	0.1
3-G.M.....	.2	.5	.1	.1	.1	.2				.1
4-S.M.....	1.4	3.2	.9	2.6	2.6	.8	.3	.3	.3	
5-M.....	1.8	2.2	1.0	3.3	5.7	.3		.1	.1	
6-S.L.M. ⁴3	.9	.9	5.0	2.2					
7-L.M. ⁴	4.1	2.4	.1	7.3	2.2					
Yellow Tinged.....			.1		.9		.1			
2-S.G.M.....										
3-G.M.....										
4-S.M.....										
5-M. ⁴										
6-S.L.M. ⁴1		.2	.1					
7-L.M. ⁴7						
Gray.....				.1						
3-G.M.....										
4-S.M.....				.1						
5-M. ⁴										
No grade ^{1,6}	9.3	5.1	2.0	9.8	7.5					
Grade	Ginned during October					Ginned during November				
	1932	1931	1930	1929	1928	1932	1931	1930	1929	1928
	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales
All grades ²	118.3	102.7	64.4	79.2	35.6	55.6	65.9	28.8	58.3	45.4
Extra White.....	3.2						.7			
Above 3-G.M.....										
3-G.M.....										
4-S.M.....										
5-M.....	.1									
6-S.L.M.....	2.8						.7			
7-L.M.....	.3									
Below 7-L.M. ⁴										
White.....	114.4	98.1	62.9	77.1	33.4	53.4	64.8	27.3	51.6	40.0
1-M.F.....										
2-S.G.M.....										
3-G.M.....	.1	.7	.1	.2				.1		
4-S.M.....	2.6	10.1	12.3	19.7	9.6	.6	.8	.8	2.2	1.0
5-M.....	53.3	50.1	37.2	46.5	16.4	8.5	12.8	4.1	14.8	10.1
6-S.L.M.....	53.4	34.8	11.4	10.6	6.8	18.7	35.4	8.3	11.7	19.8
7-L.M.....	4.8	2.9	1.3	.2	.4	11.4	13.9	9.5	11.5	5.5
8-S.G.O. ⁴3	1				11.3	1.8	4.0	11.0	2.4
9-G.O. ⁴						2.9	.1	.6	.4	1.2
Spotted.....	.7	4.6	1.4	2.1	2.2	1.2	1.1	1.1	6.7	5.2
3-G.M.....		.4	.1	.1				.1		
4-S.M.....	.2	2.8	.6	1.5	1.0	.3	.1		.8	1.1
5-M.....	.5	1.3	.4	.5	1.0	.7	.7	.5	2.3	2.7
6-S.L.M. ⁴1	.3		.2	.1	.2	.5	2.2	.9
7-L.M. ⁴1		.1	1.4	.5

See footnotes at end of table.

TABLE 5.—*Grade of American upland cotton: Quantities ginned in the United States during specified periods,¹ by States, crops of 1932, 1931, 1930, 1929, and 1928—Continued*

MISSOURI—Continued

Grade	Ginned during October					Ginned during November				
	1932	1931	1930	1929	1928	1932	1931	1930	1929	1928
	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales
Yellow Tinged										
2-S.G.M.										
3-G.M.										
4-S.M.										
5-M. ⁴										
6-S.L.M. ⁴										
7-L.M. ⁴										
Gray			0.1							
2-G.M.										
4-S.M.			.1							
5-M. ⁴										
No grade ^{4,6}						0.3		0.4		0.1
Grade	Ginned Dec. 1-Jan. 15, crop of—					Ginned after Jan. 15, crop of—				
	1932	1931	1930	1929	1928	1932	1931	1930	1929	1928
	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales
All grades ²	35.2	48.4	5.9	37.7	40.4	10.9	30.9	1.3	22.6	11.0
Extra White										
Above 3-G.M.										
3-G.M.										
4-S.M.										
5-M.										
6-S.L.M.										
7-L.M.										
Below 7-L.M.										
White	28.0	47.3	4.6	27.0	32.1	6.5	23.6	.9	13.5	7.5
1-M.F.										
2-S.G.M.										
3-G.M.										
4-S.M. ⁴							.1			
5-M.	.1	.1		.2	.3	.2				
6-S.L.M.	1.2	.8	.3	.6	.9				.6	.2
7-L.M.	3.5	5.7	2.0	5.0	1.5	.1	.7	.4	2.2	.4
8-S.G.O. ⁴	10.2	30.3	1.9	16.0	12.2		8.0	.4	3.7	2.6
9-G.O. ⁴	11.0	10.4	.4	5.2	17.2	6.2	14.8	.1	7.0	3.7
Spotted	1.3	.9	8.3	3.6	3.3	2.3	.1	.8	1.7	
3-G.M.										
4-S.M.	.1				.3					
5-M.	.3	.2		.4	1.3					
6-S.L.M. ⁴	.2	.4		2.6	.6		.2	.1	.2	.5
7-L.M. ⁴	.7	.3		5.3	1.4	3.3	2.1		.6	.3
Yellow Tinged		.1		.6						
2-S.G.M.										
3-G.M.										
4-S.M.										
5-M. ⁴										
6-S.L.M. ⁴			.1		.2					
7-L.M. ⁴					.4					
Gray										
2-G.M.										
4-S.M.										
5-M. ⁴										
No grade ^{4,6}	7.9	.1	1.3	1.8	4.7	1.1	5.0	.3	8.0	2.7

See footnotes at end of table.

TABLE 5.—*Grade of American upland cotton: Quantities ginned in the United States during specified periods,¹ by States, crops of 1932, 1931, 1930, 1929, and 1928—Continued*

NEW MEXICO

Grade	Total, crop of—					Ginned prior to Oct. 1				
	1932	1931	1930	1929	1928	1932	1931	1930	1929	1928
	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales
All grades ²	67.5	93.8	95.8	86.3	82.2	4.6	5.6	17.0	8.4	6.1
xtra White.....	38.9	81.5	82.2	58.3	68.4	4.6	5.6	17.0	8.3	6.1
Above 3-G.M.....	.2	.2	.5			.1		.4		
3-G.M.....	8.7	14.0	30.7	10.4	17.5	3.4	3.8	15.1	6.2	3.9
4-S.M.....	23.5	42.1	30.1	19.9	25.4	1.0	1.8	1.6	2.1	1.9
5-M.....	5.5	18.1	14.5	15.0	22.0	.1				.3
6-S.L.M.....	.2	4.2	2.1	3.4	2.4					
7-L.M.....	.2	1.0	.3	.6	1.1					
Below 7-L.M. ⁴6	1.9	4.0							
White.....		1.1	1.6		1.6	7.9				.1
1-M.F.....										
2-S.G.M.....					1.6					.1
3-G.M.....						1.2				
4-S.M.....						2.5				
5-M.....						.3				
6-S.L.M.....			.8							
7-L.M.....			.4							
8-S.G.O. ⁴2		3.9					
9-G.O. ⁴2							
Spotted.....		21.0	7.5	13.6	15.2	4.6				
3-G.M.....		4.9		.6	.1	.2				
4-S.M.....		13.3	5.3	7.7	4.9	2.5				
5-M.....		2.3	2.0	3.5	9.0	1.3				
6-S.L.M. ⁴5	.2	1.0	1.1	.6				
7-L.M. ⁴7	.1					
Yellow Tinged.....		.2		.1	4.2	.1				
2-S.G.M.....										
3-G.M.....		.1			.1					
4-S.M.....		.1			1.5					
5-M. ⁴					2.0	.1				
6-S.L.M. ⁴1	.4					
7-L.M. ⁴2					
Light Yellow Stained.....					2.6					
3-G.M.....										
4-S.M. ⁴8					
5-M. ⁴					1.8					
Yellow Stained.....					1.9					
3-G.M.....										
4-S.M. ⁴2					
5-M. ⁴					1.7					
No grade ⁴		6.3	3.2		2.5	1.2				

Grade	Ginned during October					Ginned during November				
	1932	1931	1930	1929	1928	1932	1931	1930	1929	1928
	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales
All grades ²	23.1	34.0	32.2	28.6	27.1	25.7	29.2	28.6	27.6	24.1
Extra White.....	22.0	33.0	28.6	27.1	23.5	10.5	25.0	25.2	19.1	22.8
Above 3-G.M.....	.1	.2	.1							
3-G.M.....	4.4	9.6	10.4	13.0	11.9	.9	.3	5.1	.2	1.7
4-S.M.....	15.1	19.5	12.5	11.8	9.5	7.1	17.6	13.5	5.3	11.7
5-M.....	2.4	3.7	5.3	2.3	2.1	2.1	6.6	5.8	11.0	9.1
6-S.L.M.....				.2			.1	.4	.7	2.4
7-L.M.....				.1				.1		.2
Below 7-L.M. ⁴3		.1		

See footnotes at end of table.

TABLE 5.—Grade of American upland cotton: Quantities ginned in the United States during specified periods,¹ by States, crops of 1932, 1931, 1930, 1929, and 1928—Continued

NEW MEXICO—Continued

Grade	Ginned during October					Ginned during November				
	1932	1931	1930	1929	1928	1932	1931	1930	1929	1928
	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales
White.....	1.1			1.5	3.6					.4
1-M.F.										
2-S.G.M.				1.5						
3-G.M.							1.0			.2
4-S.M.	.6						2.3			
5-M.	.5						.3			
6-S.L.M.										
7-L.M.										
8-S.G.O.										
9-G.O. ⁴										
Spotted.....		1.0	3.6			15.0	4.2	3.4	7.5	.0
3-G.M.			.2			4.5		.3		.1
4-S.M.	.5	2.6				10.0	3.1	1.6	2.4	.7
5-M.	.4	.7				.5	1.0	1.2	4.9	.1
6-S.L.M. ⁴	.1	.1					.1	.2	.2	
7-L.M. ⁴									.1	
Yellow, Tinged.....							.2			1.0
2-S.G.M.										
3-G.M.							.1			
4-S.M.							.1			.4
5-M. ⁴										.5
6-S.L.M.										
7-L.M. ⁴										.1
Light Yellow Stained.....										
3-G.M.										
4-S.M. ⁴										
5-M. ⁴										
Yellow Stained.....										
3-G.M.										
4-S.M. ⁴										
5-M. ⁴										
No grade ⁴										

Grade	Ginned Dec. 1-Jan. 15, crop of—					Ginned after Jan. 15, crop of—				
	1932	1931	1930	1929	1928	1932	1931	1930	1929	1928
	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales
All grades ²	10.7	17.6	16.2	19.7	19.7	3.4	7.4	1.8	2.0	5.2
Extra White.....	1.4	15.9	9.7	3.3	15.0	.4	2.0	1.7	.6	1.0
Above 3-G.M.										
3-G.M.		.3	.1							
4-S.M.	.3	3.2	2.6	.7	2.2					.1
5-M.	.8	7.7	3.3	1.6	10.2	.1	.1	.1	.1	.3
6-S.L.M.	.1	3.2	1.2	1.0	1.6					.6
7-L.M.	.2	.8	.2		1.0					.4
Below 7-L.M. ⁴		.7	2.3			.3	1.2	1.6		
White.....					1.8		1.6			2.1
1-M.F.										
2-S.G.M.										
3-G.M.										
4-S.M.										
5-M.										
6-S.L.M.										
7-L.M.										
8-S.G.O.					1.8					
9-G.O. ⁴										2.1

See footnotes at end of table.

TABLE 5.—*Grade of American upland cotton: Quantities ginned in the United States during specified periods,¹ by States, crops of 1932, 1931, 1930, 1929, and 1928—Continued*

NEW MEXICO—Continued

Grade	Ginned Dec. 1-Jan. 15, crop of—					Ginned after Jan. 15, crop of—									
	1932		1931		1930	1929		1928	1932		1931		1930	1929	
	<i>1,000 bales</i>		<i>1,000 bales</i>		<i>1,000 bales</i>	<i>1,000 bales</i>		<i>1,000 bales</i>	<i>1,000 bales</i>		<i>1,000 bales</i>		1928	<i>1,000 bales</i>	
Spotted.....	.5	.2	1.7	6.4	7.6	2.1			.8	0.6	0.6	0.1			1.6
3-G.M.	.4				.1	.1									
4-S.M.	3.1		1.5		3.5	2.5		1.3		.2	.2				.5
5-M.	1.4		.2		1.5	4.1		.5		.4	.4				.7
6-S.L.M. ⁴	.3				.7	.8		.2		.2					.1
7-L.M. ⁴					.6	.1									.4
Yellow Tinged.....					.1	3.0		.1							.2
2-S.G.M.															
3-G.M.															
4-S.M.															.1
5-M. ⁴															
6-S.L.M. ⁴															.1
7-L.M. ⁴															
Light Yellow Stained.....															.2
3-G.M.															
4-S.M. ⁴															.1
5-M. ⁴															.1
Yellow Stained.....															.2
3-G.M.															
4-S.M. ⁴															
5-M. ⁴															.2
No grade ⁴			4.1						1.7	.7	2.2	3.2			.8
															.5

NORTH CAROLINA

Grade	Total, crop of—					Ginned prior to Oct. 1									
	1932		1931		1930	1929		1928	1932		1931		1930	1929	
	<i>1,000 bales</i>		<i>1,000 bales</i>		<i>1,000 bales</i>	<i>1,000 bales</i>		<i>1,000 bales</i>	<i>1,000 bales</i>		<i>1,000 bales</i>		1928	<i>1,000 bales</i>	
All grades ²	680.3	680.3	771.2	771.2	800.6	767.0	767.0	869.3	226.1	215.6	198.5	50.2	50.2	60.1	60.1
Extra White.....		2.5							.3	1.8					.1
Above 3-G.M.....															
3-G.M.															
4-S.M.	.1														
5-M.	.8														.1
6-S.L.M.	1.4														
7-L.M.	.2														
Below 7-L.M. ⁴															
White.....	618.9	618.9	718.6	718.6	702.6	641.5	641.5	628.7	219.5	203.7	176.5	44.8	44.8	50.3	
1-M.F.															
2-S.G.M.			.6	.6	.1	.1	.1	.3							.1
3-G.M.	13.8	13.8	25.1	25.1	13.9	8.7	8.7	13.7	6.2	16.5	7.1	1.4	1.4		.3
4-S.M.	216.3	216.3	230.0	230.0	232.9	167.2	167.2	213.5	106.0	104.2	85.4	17.3	17.3		12.0
5-M.	294.0	294.0	351.0	351.0	324.3	292.6	292.6	272.0	102.1	80.0	75.6	21.5	21.5		25.4
6-S.L.M.	85.9	85.9	108.8	108.8	98.0	141.3	141.3	104.4	6.1	2.6	7.0	4.2	4.2		11.3
7-L.M.	8.3	8.3	3.1	3.1	29.3	26.2	26.2	20.6				.7	.7		1.2
8-S.G.O. ⁴	.0	.0			3.8	3.3	3.3	2.9				.1	.1		.1
9-O.O. ⁴					.3	2.1	1.3								
Spotted.....	52.8	52.8	60.3	60.3	92.3	115.4	115.4	222.8	4.7	11.9	20.8	5.3	5.3		9.4
3-G.M.	3.7	3.7	10.5	10.5	18.2	3.7	3.7	20.9		.7	2.5	5.9	.1		.7
4-S.M.	28.3	28.3	31.7	31.7	51.5	61.8	61.8	145.0	3.2	8.5	13.3	4.2	4.2		4.9
5-M.	20.8	20.8	7.4	7.4	16.8	43.3	43.3	53.0	.8	.9	1.6	1.0	1.0		3.5
6-S.L.M. ⁴	2.0	2.0	.7	.7	5.8	5.4	5.4	3.7							.3
7-L.M. ⁴					1.0	1.2	1.2	.2							

See footnotes at end of table.

TABLE 5.—*Grade of American upland cotton: Quantities ginned in the United States during specified periods,¹ by States, crops of 1932, 1931, 1930, 1929, and 1928—Continued*

NORTH CAROLINA—Continued

Grade	Total, crop of—					Ginned prior to Oct. 1				
	1932	1931	1930	1929	1928	1932	1931	1930	1929	1928
Yellow Tinged.....	1,000 bales 0.4	1,000 bales 0.4	1,000 bales 4.9	1,000 bales 3.7	1,000 bales 5.0	1,000 bales -----	1,000 bales -----	1,000 bales 1.1	1,000 bales 0.1	1,000 bales 0.1
2-S.G.M.....										
3-G.M.....	.1	.1	1.3	.2	1.5				.6	
4-S.M.....	.1	.2	1.4	.9	2.4				.5	
5-M. ⁴1	.1	1.1	1.0	.8				.1	
6-S.L.M. ⁴1		.8	.5	.2					
7-L.M. ⁴3	1.1	.1					
Light Yellow Stained.....		.1	.3	.2	.4					.1
3-G.M.....							.1			
4-S.M. ⁴1			
5-M. ⁴1	.3	.2	.2					.1
Yellow Stained.....			.2		.4					
3-G.M.....										
4-S.M. ⁴1		.2					
5-M. ⁴1		.2					
Gray.....	5.6	1.8		3.4	8.9	0.1				
3-G.M.....	.7				2.0					
4-S.M.....	3.5	1.0		1.9	0.2	.1				
5-M. ⁴	1.4	.8		1.5	.7					
Blue Stained.....			.1		1.1					
3-G.M. ⁴1		.2					
4-S.M. ⁴7					
5-M. ⁴2					
No grade ^{4,6}1		.2	2.8	1.7			.1		.1

Grade	Ginned during October					Ginned during November				
	1932	1931	1930	1929	1928	1932	1931	1930	1929	1928
All grades ²	1,000 bales 264.1	1,000 bales 382.2	1,000 bales 391.0	1,000 bales 340.4	1,000 bales 437.0	1,000 bales 125.2	1,000 bales 134.7	1,000 bales 133.8	1,000 bales 207.7	1,000 bales 243.2
Extra White.....	.6					.1	.1			.1
Above 3-G.M.....										
3-G.M.....										
4-S.M.....										.1
5-M.....	.1									
6-S.L.M.....	.4					.1				
7-L.M.....	.1					.1				
Below 7-L.M. ⁴										
White.....	247.7	357.8	363.1	296.3	326.8	108.5	121.0	114.5	172.0	177.2
1-M.F.....										
2-S.G.M.....		2	.1		.3					
3-G.M.....	6.7	7.4	6.4	5.9	9.0	.7	.6	.3	1.2	3.4
4-S.M.....	80.1	95.5	131.3	100.5	119.5	23.4	18.6	12.1	35.7	60.9
5-M.....	124.5	186.2	183.5	138.9	149.2	51.4	68.2	46.8	84.7	75.2
6-S.L.M.....	34.5	67.4	29.3	44.1	42.5	29.0	32.9	43.5	42.8	29.0
7-L.M.....	1.9	1.1	2.4	6.2	5.7	3.7	1.6	11.5	7.1	2.5
8-S.G.O. ⁴1	.6	.6	.3		.3	.5	.1
9-G.O. ⁴1						
Spotted.....	12.2	23.6	36.8	43.7	107.1	14.6	11.6	19.1	34.9	63.8
3-G.M.....	1.6	5.9	10.5	2.5	12.0	.8	1.5	1.2	1.0	4.7
4-S.M.....	7.6	14.1	23.7	28.4	69.5	8.2	7.5	11.0	20.0	46.6
5-M.....	2.8	3.4	2.6	12.2	23.8	5.4	2.5	5.5	12.2	12.2
6-S.L.M. ⁴2	.2		.5	1.1	.2	.1	1.2	1.7	.3
7-L.M. ⁴1	.1				.2	

See footnotes at end of table.

TABLE 5.—*Grade of American upland cotton: Quantities ginned in the United States during specified periods,¹ by States, crops of 1932, 1931, 1930, 1929, and 1928—Continued*

NORTH CAROLINA—Continued

Grade	Ginned during October					Ginned during November				
	1932	1931	1930	1929	1928	1932	1931	1930	1929	1928
Yellow Tinged.....	1,000 bales 0.2	1,000 bales 0.2	1,000 bales 1.0	1,000 bales 0.2	1,000 bales 2.0	1,000 bales 0.1	1,000 bales 0.2	1,000 bales 0.8	1,000 bales 0.7	
2-S.G.M.										
3-G.M.	.1	.1	.5		.6			.1	.2	.2
4-S.M.		.1	.4	.1	1.0			.1	.3	.3
5-M. ⁴	.1		.1	.1	.3		.1		.2	.2
6-S.L.M. ⁴					.1				.1	
7-L.M. ⁴										
Light Yellow Stained.....		.1		.1	.2					.1
3-G.M.										
4-S.M. ⁴										.1
5-M. ⁴		.1		.1	.1					
Yellow Stained.....			.1							
3-G.M.										
4-S.M. ⁴			.1							
5-M. ⁴										
Gray.....	3.3	.5			.7	2.0	1.1			1.3
3-G.M.	.5				.1	.2				.3
4-S.M.	2.3	.3			.4	1.0	.6			.9
5-M. ⁴	.5	.2			.2	.8	.5			.1
Blue Stained.....										
3-G.M. ⁴										
4-S.M. ⁴										
5-M. ⁴										
No grade ^{4,6}	.1			.1	.1					

Grade	Ginned Dec. 1-Jan. 15, crop of—					Ginned after Jan. 15, crop of—				
	1932	1931	1930	1929	1928	1932	1931	1930	1929	1928
All grades ²	1,000 bales 51.9	1,000 bales 31.2	1,000 bales 56.7	1,000 bales 139.4	1,000 bales 104.9	1,000 bales 13.0	1,000 bales 7.5	1,000 bales 20.6	1,000 bales 29.3	1,000 bales 24.1
Extra White.....										
Above 3-G.M.										
3-G.M.										
4-S.M.										
5-M.										
6-S.L.M.										
7-L.M.										
Below 7-L.M. ⁴										
White.....	35.8	28.8	44.0	109.7	58.0	7.4	6.4	14.5	18.7	16.4
1-M.F.										
2-S.G.M.										
3-G.M.	.2	.4	.1	.1	.9		.2		.1	.1
4-S.M.	5.2	9.2	3.6	11.7	13.7	2.6	2.5	.5	2.0	1.4
5-M.	13.8	14.3	14.3	42.1	18.9	2.2	2.3	4.1	5.4	3.3
6-S.L.M.	14.0	4.8	14.2	44.3	18.2	1.7	1.1	3.4	5.9	3.4
7-L.M.	1.0	.1	10.8	9.5	5.9	.7	.3	3.0	3.2	5.3
8-S.G.O. ⁴	.1		.9	1.0	.3	.2		2.4	1.1	1.8
9-G.O. ⁴			.1	1.0	.1		.2	1.0		1.1
Spotted.....	15.9	2.2	11.6	24.3	38.3	5.4	1.0	4.0	7.2	4.2
3-G.M.	.2	.6	.5	.1	2.8	.4		.1		.1
4-S.M.	5.8	1.1	2.8	8.2	22.7	1.5	.5	.7	1.0	1.3
5-M.	9.5	.4	4.7	14.4	12.0	2.3	.2	1.4	3.6	1.5
6-S.L.M. ⁴	.4	.1	3.1	1.4	.7	1.2	.3	1.6	1.8	1.3
7-L.M. ⁴			.5	.2	.1			.3	.9	

See footnotes at end of table.

TABLE 5.—*Grade of American upland cotton: Quantities ginned in the United States during specified periods,¹ by States, crops of 1932, 1931, 1930, 1929, and 1928—Continued*

NORTH CAROLINA—Continued

Grade	Ginned Dec. 1-Jan. 15, crop of—					Ginned after Jan. 15, crop of—				
	1932	1931	1930	1929	1928	1932	1931	1930	1929	1928
	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales
Yellow Tinged	0.1	0.1	0.9	1.3	1.7	0.1	0.1	1.7	1.3	0.6
2-S.G.M.										
3-G.M.				.1						
4-S.M.		.1	.3	.4	.9					
5-M. ⁴			.4	.4	.1				.6	.3
6-S.L.M. ⁴				.1					.7	.3
7-L.M. ⁴				.4					.3	.7
Light Yellow Stained									.3	.1
3-G.M.										
4-S.M. ⁴										
5-M. ⁴									.3	.1
Yellow Stained						.3			.1	
3-G.M.										
4-S.M. ⁴										
5-M. ⁴										
Gray	.1	.1		2.5	6.3	.1	0.1		.9	.6
3-G.M.										
4-S.M.		.1		1.6	4.7		1		.3	.2
5-M. ⁴				.9	.1				.6	.3
Blue Stained				.1						
3-G.M. ⁴				.1						
4-S.M. ⁴				.1						
5-M. ⁴										
No grade ^{4,5}				.1	1.6	.1				
									1.1	1.4

OKLAHOMA

Grade	Total, crop of—					Ginned prior to Oct. 1				
	1932	1931	1930	1929	1928	1932	1931	1930	1929	1928
	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales
All grades ²	1,072.0	1,235.9	856.8	1,125.6	1,187.0	290.2	281.1	276.6	255.1	208.1
Extra White	34.8	.3			.1					
Above 3-G.M.										
3-G.M.	1.6									
4-S.M.	9.0	.1								
5-M.	13.5									
6-S.L.M.	7.9	.1								
7-L.M.	2.8	.1								
Below 7-L.M. ⁴										
White	823.7	1,047.5	760.3	984.1	1,026.0	286.7	275.7	256.2	240.1	259.8
1-M.F.					.1					
2-S.G.M.	.1	.1	.1	.1	.7					
3-G.M.	24.5	69.6	17.1	9.9	125.7	20.9	38.0	13.8	4.5	61.9
4-S.M.	286.9	370.0	209.7	125.6	453.7	145.4	184.8	144.8	46.4	168.4
5-M.	326.2	186.7	258.3	307.2	175.4	95.5	49.1	81.4	93.6	20.6
6-S.L.M.	142.3	147.7	195.6	292.8	117.1	21.9	3.6	14.8	64.9	2.4
7-L.M.	37.6	174.6	75.2	209.6	80.3	2.8	.2	1.4	28.0	.2
8-S.G.O.	5.8	91.8	4.1	36.6	60.6	.1				1.8
9-G.O. ⁴	.3	7.0	.2	2.3	12.4	.1				

See footnotes at end of table.

TABLE 5.—*Grade of American upland cotton: Quantities ginned in the United States during specified periods,¹ by States, crops of 1932, 1931, 1930, 1929, and 1928—Continued*

OKLAHOMA—Continued

Grade	Total, crop of—					Ginned prior to Oct. 1				
	1932	1931	1930	1929	1928	1932	1931	1930	1929	1928
	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales
Spotted	210.8	177.2	92.5	92.6	108.2	3.5	5.3	20.4	14.6	8.1
3-G.M.	16.5	5.2	4.3	.1	3.9	.2	1.6	3.5	—	1.6
4-S.M.	102.4	41.7	21.3	9.5	28.7	2.6	3.2	12.0	3.5	6.0
5-M.	59.1	58.6	33.4	24.8	19.9	.5	.4	4.3	8.1	.5
6-S.L.M. ⁴	22.6	50.0	27.5	37.1	22.5	.1	.1	.5	2.9	—
7-L.M. ⁴	10.2	21.7	6.0	21.1	33.2	.1	—	.1	.1	—
Yellow Tinged	2.4	9.6	3.8	37.3	5.8	—	.1	—	.4	.1
2-S.G.M.	—	—	—	—	—	—	—	—	—	—
3-G.M.	.4	.4	—	—	.1	—	—	—	—	—
4-S.M. ⁴	1.2	.7	.2	.7	.2	—	.1	—	.2	.1
5-M. ⁴	.4	2.3	1.2	5.3	.3	—	—	—	—	—
6-S.L.M. ⁴	.4	5.0	2.1	14.7	1.5	—	—	—	—	—
7-L.M. ⁴	—	1.2	.3	16.6	3.7	—	—	—	—	—
Light Yellow Stained	.1	.2	—	2.1	.1	—	—	—	—	—
3-G.M.	—	.1	—	—	—	—	—	—	—	—
4-S.M. ⁴	.1	.1	—	.1	.1	—	—	—	—	—
5-M. ⁴	—	—	—	2.0	—	—	—	—	—	—
Yellow Stained	—	—	—	2.5	—	—	—	—	—	—
3-G.M.	—	—	—	—	—	—	—	—	—	—
4-S.M. ⁴	—	—	—	—	—	—	—	—	—	—
5-M. ⁴	—	—	—	2.5	—	—	—	—	—	—
Gray	.1	.1	.1	.1	.2	—	—	—	—	—
3-G.M.	—	—	—	—	—	—	—	—	—	—
4-S.M.	—	.1	—	—	—	—	—	—	—	—
5-M. ⁴	.1	—	.1	.1	.2	—	—	—	—	—
No grade ^{4,6}	—	.1	1.0	.1	6.9	46.6	—	—	—	.1
Grade	Ginned during October					Ginned during November				
	1932	1931	1930	1929	1928	1932	1931	1930	1929	1928
	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales
All grades ²	464.5	465.0	313.0	460.6	484.8	241.5	307.5	203.9	254.4	196.3
Extra White	—	11.9	.1	—	—	.1	22.8	.1	—	—
Above 3-G.M.	—	.7	—	—	—	—	.9	—	—	—
3-G.M.	.7	—	—	—	—	—	—	—	—	—
4-S.M.	3.8	.1	—	—	—	.1	5.2	—	—	—
5-M.	4.4	—	—	—	—	—	9.0	—	—	—
6-S.L.M.	2.3	—	—	—	—	—	5.6	.1	—	—
7-L.M.	.7	—	—	—	—	—	2.1	—	—	—
Below 7-L.M. ⁴	—	—	—	—	—	—	—	—	—	—
White	401.8	410.1	203.6	436.9	460.3	108.0	246.6	185.2	217.4	162.5
1-M.F.	—	—	—	—	—	.1	—	—	—	—
2-S.G.M.	.1	.1	—	.1	.2	—	—	.1	—	.1
3-O.M.	3.0	29.0	2.7	5.0	61.1	.6	1.7	.6	.3	1.8
4-S.M.	130.1	174.8	49.7	71.9	268.7	10.5	8.2	14.6	6.1	14.4
5-M.	196.2	96.2	119.9	168.7	102.3	31.8	38.4	54.3	39.9	36.0
6-S.L.M.	62.0	57.9	85.0	135.0	22.6	48.2	72.4	83.5	67.6	51.4
7-L.M.	10.0	42.1	33.8	52.9	4.4	14.9	88.5	31.4	79.9	28.8
8-S.G.O. ⁴	.4	9.4	2.4	2.9	.7	2.0	35.4	.6	22.4	25.9
9-G.O. ⁴	—	.6	.1	.4	.2	—	2.0	.1	1.2	3.2

See footnotes at end of table.

TABLE 5.—*Grade of American upland cotton: Quantities ginned in the United States during specified periods,¹ by States, crops of 1932, 1931, 1930, 1929, and 1928—Continued*

OKLAHOMA—Continued

Grade	Ginned during October					Ginned during November									
	1932		1931		1930	1929		1928	1932		1931		1930	1929	
	<i>1,000 bales</i>		<i>1,000 bales</i>		<i>1,000 bales</i>	<i>1,000 bales</i>		<i>1,000 bales</i>	<i>1,000 bales</i>		<i>1,000 bales</i>		<i>1,000 bales</i>	<i>1,000 bales</i>	
Spotted	50.5	53.7			19.1	13.3	23.8		109.0	59.9			22.7	26.7	31.4
3-G.M.	7.4	2.5	.7					1.4		8.8	1.0	.1		1	.4
4-S.M.	36.2	22.0	5.6	4.1		16.6			58.6	15.5	3.3	1.5		5.3	
5-M.	6.3	21.7	8.3	6.9		5.2			36.9	28.5	12.6	5.6		0.4	
6-S.L.M. ⁴	.5	6.8	3.9	1.9		.5			4.6	12.8	5.8	10.8		7.2	
7-L.M. ⁴	.1	.7	.6	.4		.1			.1	2.1	.9	8.7		9.1	
Yellow Tinged	.2	.9	.1	.4		.1			1.7	.9	1.0	9.0		.9	
2-S.G.M.															
3-G.M.	.1	.2								.3	.2				.1
4-S.M.	.1	.3		.1		.1			1.1	.2	.2	.2		.2	
5-M. ⁴	.2	.1	.2						.2	.3	.6	1.3		.2	
6-S.L.M. ⁴	.2			.1					.1	.1	.2	2.1		.3	
7-L.M. ⁴										.1		5.4		.3	
Light Yellow Stained	.1							.1							.1
3-G.M.															
4-S.M. ⁴	.1							.1							.1
5-M. ⁴															
Yellow Stained															.5
3-G.M.															
4-S.M. ⁴															
5-M. ⁴															.5
Gray															
3-G.M.															
4-S.M.															
5-M. ⁴															
No grade ^{4,6}								.4							.7
															1.5

Grade	Ginned Dec. 1-Jan. 15, crop of—					Ginned after Jan. 15, crop of—										
	1932		1931		1930	1929		1928	1932		1931		1930	1929		
	<i>1,000 bales</i>		<i>1,000 bales</i>		<i>1,000 bales</i>	<i>1,000 bales</i>		<i>1,000 bales</i>	<i>1,000 bales</i>		<i>1,000 bales</i>		<i>1,000 bales</i>	<i>1,000 bales</i>		
All grades ²	55.6	145.3			53.1	144.7		202.2			20.2	37.0		5.2	20.8	35.6
Extra White	.1	.1														
Above 3-G.M.																
3-G.M.																
4-S.M.																
5-M.	.1															
6-S.L.M.																
7-L.M.		.1														
Below 7-L.M. ⁴																
White	22.4	102.1			23.4	82.7		134.0			4.8	13.0		1.9	7.0	9.4
1-M.F.															.2	
2-S.G.M.															.1	
3-G.M.		.7			.5	.1		.7			.2				.1	
4-S.M.	.7	2.0	.5		1.0	2.1			.2		.2				.1	
5-M.	2.3	2.8	2.5		4.9	9.4			.4		.2				.3	
6-S.L.M.	9.1	12.2	11.5		24.1	37.9			1.1		1.6				2.8	
7-L.M.	8.4	39.0	7.9		44.6	44.0			1.5		4.8				2.9	
8-S.G.O. ⁴	1.7	41.3	1.0		7.7	31.9			1.6		5.7				2.1	
9-G.O. ⁴	.2	4.1			.3	8.0					.3				1.0	

See footnotes at end of table.

TABLE 5.—*Grade of American upland cotton: Quantities ginned in the United States during specified periods,¹ by States, crops of 1932, 1931, 1930, 1929, and 1928—Continued*

OKLAHOMA—Continued

Grade	Ginned Dec. 1-Jan. 15, crop of—					Ginned after Jan. 15, crop of—				
	1932	1931	1930	1929	1928	1932	1931	1930	1929	1928
	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales
Spotted.....	32.8	39.0	27.2	33.2	42.8	15.0	19.3	3.1	4.8	2.1
3-G.M.....	1	1			.5					
4-S.M.....	4.9	.9	.4	.4	.7	.1	.1			.1
5-M.....	14.2	6.0	7.9	3.5	4.6	1.2	2.0	.3	.7	.2
6-S.L.M. ⁴	11.1	20.0	15.5	19.8	14.1	6.3	10.3	1.8	1.7	.7
7-L.M. ⁴	2.5	12.0	3.4	9.5	22.9	7.4	6.9	1.0	2.4	1.1
Yellow Tinged.....	.2	3.3	2.5	21.7	4.2	.8	4.4	.2	5.8	.5
2-S.G.M.....										
3-G.M.....										
4-S.M.....										
5-M. ⁴2	.8	.5	3.6	.1	1.0		.1		
6-S.L.M. ⁴		2.1	1.7	10.4	1.1	.3	2.6	.2	2.0	.1
7-L.M. ⁴3	.3	7.5	3.0		.8		3.7	.4
Light Yellow Stained.....								.1		.3
3-G.M.....										.1
4-S.M. ⁴										
5-M. ⁴3
Yellow Stained.....					2.0					
3-G.M.....										
4-S.M. ⁴										
5-M. ⁴					2.0					
Gray.....	.1				.1	.2				
3-G.M.....										
4-S.M.....										
5-M. ⁴1				.1	.2				
No grade ^{4,6}										
	.7			3.3	21.0	.1	.2		2.9	23.6

SOUTH CAROLINA

Grade	Total, crop of—					Ginned prior to Oct. 1				
	1932	1931	1930	1929	1928	1932	1931	1930	1929	1928
	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales
All grades ²	722.2	1,010.2	1,015.3	833.1	744.4	295.0	370.6	377.4	162.6	120.7
Extra White.....	2.3				.2	1.1				.1
Above 3-G.M.....										
3-G.M.....										
4-S.M.....										
5-M.....	.1									
6-S.L.M.....	.4				.1	.2				.1
7-L.M.....	.1				.1					
Below 7-L.M. ⁴										
White.....	622.9	959.4	830.6	606.7	486.4	284.7	360.1	334.3	137.5	86.1
1-M.F.....										
2-S.G.M.....			.7	.3		.3		.7	.3	.1
3-G.M.....	10.3	39.6	21.1	3.3	19.9	7.6	29.9	19.1	2.2	2.0
4-S.M.....	210.6	373.4	271.5	134.3	187.9	132.6	181.9	161.9	58.5	18.4
5-M.....	327.0	450.4	383.2	277.9	166.3	133.9	139.2	131.1	62.7	30.3
6-S.L.M.....	69.8	93.6	134.3	130.5	87.9	10.5	8.3	21.4	12.4	25.6
7-L.M.....	5.0	1.7	25.9	46.1	34.3	.2	.1	.5	1.5	8.3
8-S.G.O. ⁴2	.1	3.0	12.9	7.2				.2	1.2
9-G.O. ⁴3	1.7	2.6			.2

See footnotes at end of table.

TABLE 5.—*Grade of American upland cotton: Quantities ginned in the United States during specified periods,¹ by States, crops of 1932, 1931, 1930, 1929, and 1928—Continued*

SOUTH CAROLINA—Continued

Grade	Total, crop of—					Ginned prior to Oct. 1				
	1932	1931	1930	1929	1928	1932	1931	1930	1929	1928
	1,000 bales 95.0	1,000 bales 49.8	1,000 bales 168.8	1,000 bales 213.1	1,000 bales 249.9	1,000 bales 9.1	1,000 bales 10.4	1,000 bales 41.2	1,000 bales 24.6	1,000 bales 33.3
Spotted										
3-G.M.	3.3	12.0	23.1	3.8	17.7	1.0	3.0	6.7	.9	3.2
4-S.M.	48.2	28.6	103.4	109.2	142.8	6.1	6.9	26.5	10.9	14.8
5-M.	39.8	8.4	34.6	80.1	68.4	2.0	.5	7.7	3.7	12.0
6-S.L.M. ⁴	3.5	.8	6.7	18.3	20.1			.3	.1	2.9
7-L.M. ⁴	.2		1.0	1.7	.9					.4
Yellow Tinged	1.4	.7	6.5	10.5	4.0			.1	1.7	.6
2-S.G.M.										
3-G.M.	.1	.2	1.1	.7	.6				.4	.1
4-S.M.	.7	.4	3.2	5.6	2.1			.1	1.0	.4
5-M. ⁴	.6	.1	1.4	3.2	.9			.3		.2
6-S.L.M. ⁴			.6	.9	.3					.1
7-L.M. ⁴			.2	.1	.1					
Light Yellow Stained				.2	.3				.1	
3-G.M.										
4-S.M. ⁴				.1	.2			.1		
5-M. ⁴			.1	.1	.2					.1
Yellow Stained						.1	.1			
3-G.M.										
4-S.M. ⁴						.1	.1			
5-M. ⁴										
Gray	.6	.3		1.6	1.8	.1				
3-G.M.	.1					.6				
4-S.M.	.4	.2		.7	1.0	.1				
5-M. ⁴	.1	.1		.9	.2					
Blue Stained						.4				
3-G.M. ⁴										
4-S.M. ⁴						.3				
5-M. ⁴						.1				
No grade ^{4,6}				.2	.4	1.7			.1	
										.3

Grade	Ginned during October					Ginned during November				
	1932	1931	1930	1929	1928	1932	1931	1930	1929	1928
	1,000 bales 258.8	1,000 bales 459.4	1,000 bales 436.7	1,000 bales 371.2	1,000 bales 392.7	1,000 bales 112.1	1,000 bales 144.7	1,000 bales 128.2	1,000 bales 168.8	1,000 bales 168.5
All grades ²										
Extra White	.9					.1	.2			
Above 3-G.M.										
3-G.M.										
4-S.M.										
5-M.							.1			
6-S.L.M.								.1		
7-L.M.							.1			
Below 7-L.M. ⁴										
White	226.3	432.0	349.6	261.0	258.5	85.2	134.9	103.1	111.6	107.4
1-M.F.										
2-S.G.M.										
3-G.M.	2.4	8.3	1.8	.9	14.8	.3	.9	.1	.1	2.2
4-S.M.	57.9	140.1	94.7	58.9	96.3	15.9	37.0	11.8	12.1	43.7
5-M.	126.7	228.8	186.0	143.5	84.4	49.9	69.7	46.6	51.7	39.8
6-S.L.M.	36.5	54.2	63.9	50.7	39.6	17.4	26.8	35.1	36.9	14.9
7-L.M.	2.7	.6	3.2	6.0	17.6	1.0	.5	8.9	9.6	5.2
8-S.G.O. ⁴	.1			.9	4.5	.1		.5	1.1	.0
9-G.O. ⁴				.1	1.1			.1	.1	.7

See footnotes at end of table.

TABLE 5.—*Grade of American upland cotton: Quantities ginned in the United States during specified periods,¹ by States, crops of 1932, 1931, 1930, 1929, and 1928—Continued*

SOUTH CAROLINA—Continued

Grade	Ginned during October					Ginned during November				
	1932	1931	1930	1929	1928	1932	1931	1930	1929	1928
	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales
Spotted.....	30.9	28.9	84.2	103.5	131.9	26.6	9.4	24.7	54.5	59.8
3-G.M.....	1.5	7.4	14.7	2.1	9.5	.5	1.3	1.5	.7	3.8
4-S.M.....	17.6	15.4	56.1	57.8	76.8	12.6	5.0	15.3	25.9	37.7
5-M.....	11.1	3.9	13.0	40.2	31.2	12.7	2.6	6.7	22.9	16.6
6-S.L.M. ⁴7	.2	.4	3.3	14.0	.8	.5	1.0	4.7	1.6
7-L.M. ⁴1	.4			.2	.3	.1
Yellow Tinged.....	.3	.4	2.9	6.3	1.4		.2	.4	2.3	.4
2-S.G.M.....										
3-G.M.....	.1	.1	.7	.4	.3		.1			
4-S.M.....	.1	.2	2.0	4.1	.9		.1	.2	1.1	.2
5-M. ⁴1	.1	.2	1.7	.2			.1	.6	.1
6-S.L.M. ⁴1				.1	.3	
7-L.M. ⁴1
Light Yellow Stained.....					.3	.1				
3-G.M.....										
4-S.M. ⁴2						
5-M. ⁴1	.1					
Yellow Stained.....					.1	.1				
3-G.M.....										
4-S.M. ⁴1	.1					
5-M. ⁴										
Gray.....	.4	.1				.1	.1	.2	.3	.5
3-G.M.....	.1									.2
4-S.M.....	.3	.1				.1		.1		.3
5-M. ⁴1	.1		
Blue Stained.....										
3-G.M. ⁴										
4-S.M. ⁴										
5-M. ⁴										
No grade ^{4,6}5				.1	.4
Grade	Ginned Dec. 1-Jan. 15, crop of—					Ginned after Jan. 15, crop of—				
	1932	1931	1930	1929	1928	1932	1931	1930	1929	1928
	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales
All grades ²	42.0	25.1	56.0	106.7	49.7	14.3	10.4	17.0	23.8	12.8
Extra White.....						.1				
Above 3-G.M.....										
3-G.M.....										
4-S.M.....										
5-M.....										
6-S.L.M.....										
7-L.M.....										
Below 7-L.M. ⁴										
White.....	20.2	22.9	41.4	79.1	26.7	6.5	9.5	11.2	17.5	7.7
1-M.F.....										
2-S.G.M.....										
3-G.M.....										
4-S.M.....	2.8	8.1	2.2	3.7	8.1	1.5	0.3	.9	1.1	1.4
5-M.....	12.3	10.3	14.3	16.5	9.3	4.2	2.4	5.2	3.5	2.5
6-S.L.M.....	4.7	3.5	11.2	26.0	6.1	.7	.7	2.7	3.6	1.7
7-L.M.....	.4	.4	11.9	22.4	2.2	.1	.1	1.4	6.6	1.0
8-S.G.O. ⁴1	1.7	8.3	.1			.8	2.4	.5
9-O.O. ⁴1	1.2	.2			.1	.3	.4

See footnotes at end of table.

TABLE 5.—*Grade of American upland cotton: Quantities ginned in the United States during specified periods,¹ by States, crops of 1932, 1931, 1930, 1929, and 1928—Continued*

SOUTH CAROLINA—Continued

Grade	Ginned Dec. 1-Jan. 15, crop of—					Ginned after Jan. 15, crop of—				
	1932	1931	1930	1929	1928	1932	1931	1930	1929	1928
	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales
Spotted.....	21.6	2.2	14.1	25.1	20.6	6.8	0.9	4.6	5.4	4.3
3-G.M.....	.1	.3	.2	.1	1.2	.2				
4-S.M.....	8.9	1.1	4.6	5.1	12.1	3.0	.2	.9	.5	1.4
5-M.....	11.2	.7	5.5	11.4	6.6	2.8	.7	1.7	1.9	2.0
6-S.L.M. ⁴	1.3	.1	3.3	7.7	.7	.7		1.7	2.5	.9
7-L.M. ⁴1		.5	.8		.1		.3	.5	
Yellow Tinged.....	.2		.4	1.0	.8	.9		1.1	.4	.6
2-S.G.M.....										
3-G.M.....						.2				
4-S.M.....	.2			.2		.4				.2
5-M. ⁴3	.4		.1	.5		.2	
6-S.L.M. ⁴1	.3		.1			.4	.2
7-L.M. ⁴1					.2	
Light Yellow Stained.....									.1	.1
3-G.M.....										
4-S.M. ⁴1
5-M. ⁴1		
Yellow Stained.....										
3-G.M.....										
4-S.M. ⁴										
5-M. ⁴										
Gray.....				1.0	1.2					.3
3-G.M.....										
4-S.M.....				.3	.6					.1
5-M. ⁴7	.2					.2
Blue Stained.....				.2						.2
3-G.M. ⁴										
4-S.M. ⁴2						.1
5-M. ⁴1
No grade ^{4,6}1	.3	.4					.1

TENNESSEE

Grade	Total, crop of—					Ginned prior to Oct. 1				
	1932	1931	1930	1929	1928	1932	1931	1930	1929	1928
	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales
All grades ²	467.5	578.0	371.4	504.3	423.5	61.3	48.4	87.4	71.5	64.7
Extra White.....	.6			.1		.1				
Above 3-G.M.....										
3-G.M.....										
4-S.M.....				.1						
5-M.....										
6-S.L.M.....	.5									
7-L.M.....	.3									
Below 7-L.M. ⁴										
White.....	394.2	568.0	349.0	431.6	362.0	58.3	48.0	84.8	67.0	64.6
1-M.F.....										
2-S.G.M.....		.1	.2	.2	1.0					
3-G.M.....	2.0	20.6	41.9	30.0	41.5	.6	6.5	19.6	14.8	23.4
4-S.M.....	117.9	218.1	162.5	215.2	135.6	20.4	34.7	48.8	37.4	33.7
5-M.....	195.9	210.0	108.0	119.5	112.2	25.9	6.7	16.2	13.2	6.6
6-S.L.M.....	50.8	73.0	22.3	27.7	43.8	2.1	.1	.2	1.4	.1
7-L.M.....	13.8	17.1	8.9	23.9	10.6	.3				.2
8-S.G.O. ⁴	8.6	12.3	4.2	13.2	11.7					
9-G.O. ⁴	5.2	6.8	1.0	1.9	4.7					

See footnotes at end of table.

TABLE 5.—*Grade of American upland cotton: Quantities ginned in the United States during specified periods,¹ by States, crops of 1932, 1931, 1930, 1929, and 1928—Continued*

TENNESSEE—Continued

Grade	Total, crop of—					Ginned prior to Oct. 1				
	1932	1931	1930	1929	1928	1932	1931	1930	1929	1928
	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales
Spotted	69.1	16.3	20.1	67.5	59.0	2.7	0.4	2.6	4.4	0.1
3-G.M.	1.4	.9	2.1	1.8	.7	.3	.2	1.0	.9	
4-S.M.	37.1	3.8	6.5	16.4	17.4	1.9	.2	1.5	2.4	.1
5-M.	15.9	3.7	7.1	22.4	20.5	.4		.1	.9	
6-S.L.M. ⁴	7.0	3.9	3.6	19.9	9.1	.1			.2	
7-L.M. ⁴	7.7	4.0	.8	7.0	5.3					
Yellow Tinged		.6	.7	1.5	2.6	.7				.1
2-S.G.M.										
3-G.M.										
4-S.M.		.1	.1	.2		.3				
5-M. ⁴		.2	.1	.6	.4	.3				.1
6-S.L.M. ⁴		.1	.2	.5	1.0	.1				
7-L.M. ⁴		.1	.3	.2	1.2					
Gray		.1	.1		.1	.2				
3-G.M.										
4-S.M.										
5-M. ⁴		.1			.1	.1				
Blue Stained					.2					
3-G.M. ⁴										
4-S.M. ⁴										
5-M. ⁴										
No grade ^{4,6}	3.0	2.9	.8	2.2	1.5					

Grade	Ginned during October					Ginned during November				
	1932	1931	1930	1929	1928	1932	1931	1930	1929	1928
	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales
All grades ⁴	201.1	288.9	169.0	220.0	154.0	123.0	167.6	85.2	98.5	110.8
Extra White	.2			0.1	0.1	0.1				
Above 3-G.M.										
3-G.M.										
4-S.M.										
5-M.				.1		.1				
6-S.L.M.		.1								
7-L.M.		.1								
Below 7-L.M. ⁴										
White	186.5	286.8	165.8	216.5	146.8	100.8	163.4	77.3	86.1	101.3
1-M.F.										
2-S.G.M.			.1	.2	.2	1.1				
3-G.M.	1.0	12.5	19.3	11.7	16.2	.4	1.1	2.7	2.7	1.3
4-S.M.	58.7	139.7	89.8	144.0	73.4	25.2	33.6	20.2	28.5	22.7
5-M.	108.4	115.0	52.7	56.1	47.8	50.8	76.1	33.5	38.6	48.1
6-S.L.M.	17.4	18.6	3.7	4.2	7.9	18.0	46.6	14.0	10.7	28.5
7-L.M.	1.0	.9	.1	.3	.4	4.1	5.5	4.8	6.0	2.4
8-S.G.O.							2.0	.4	1.8	1.8
9-G.O.							.3	.1	.3	
Spotted	14.3	2.1	3.2	9.4	7.0	21.9	4.0	7.5	12.4	9.1
3-G.M.	.5	.4	.8	.9	.4	.4	.3	.2		.1
4-S.M.	10.7	1.3	2.0	6.9	3.0	16.3	2.0	2.3	4.1	3.1
5-M.	2.9	.4	.4	1.6	2.7	4.0	1.4	3.9	5.9	4.9
6-S.L.M. ⁴	.2					.8	.9	.3	2.0	.9
7-L.M. ⁴						.1	.3		.4	.1
Yellow Tinged						.1	.1	.2	.3	
2-S.G.M.										
3-G.M.										
4-S.M.										
5-M. ⁴										.2
6-S.L.M. ⁴										
7-L.M. ⁴										

See footnotes at end of table.

TABLE 5.—Grade of American upland cotton: Quantities ginned in the United States during specified periods,¹ by States, crops of 1932, 1931, 1930, 1929, and 1928—Continued

TENNESSEE—Continued

Grade	Ginned during October					Ginned during November				
	1932	1931	1930	1929	1928	1932	1931	1930	1929	1928
	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales
Gray.....										
3-G.M.....										
4-S.M.....										
5-M. ⁴										
Blue Stained.....										
3-G.M. ⁴										
4-S.M. ⁴										
5-M. ⁴										
No grade ^{4,6}	0.1					0.1		0.1		0.1
 Grade										
Ginned Dec. 1-Jan. 15, crop of—										
1932	1931	1930	1929	1928	1932	1931	1930	1929	1928	
1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	
All grades ²	66.0	48.1	25.6	77.0	75.2	16.1	25.0	4.2	31.3	18.8
Extra White.....										
Above 3-G.M.....										
3-G.M.....										
4-S.M.....										
5-M.....										
6-S.L.M.....										
7-L.M.....										
Below 7-L.M. ⁴										
White.....	42.3	43.8	18.4	45.3	38.2	6.3	18.0	2.7	16.7	11.1
1-M.F.....										
2-S.G.M.....										
3-G.M.....	.2	.1	.8	.6	.3	.2				
4-S.M.....	3.1	7.1	3.3	6.4	5.4	1.5	3.0	.4	.9	.4
5-M.....	8.5	10.5	4.4	10.2	8.0	2.3	1.7	1.2	1.4	1.7
6-S.L.M.....	12.3	6.7	4.1	9.7	7.2	1.0	1.0	.3	1.7	2.1
7-L.M.....	7.8	9.0	3.5	12.4	5.6	.6	1.7	.5	5.0	2.2
8-S.G.O. ⁴	5.8	8.3	2.3	5.4	8.6	.8	3.6	.1	6.2	.28
9-G.O. ⁴	4.8	2.0	.7	.4	2.8	.1	4.7		1.5	.9
Spotted.....	21.3	4.0	5.5	29.3	36.2	8.9	5.8	1.3	12.0	6.6
3-G.M.....	.2		.1		.2					
4-S.M.....	7.3	.3	.7	2.3	10.4	.9			.7	.8
5-M.....	7.4	1.3	2.3	11.7	16.1	1.2	.6	.4	2.3	2.8
6-S.L.M. ⁴	3.4	1.7	1.9	12.2	5.3	2.4	1.9	.8	5.5	2.1
7-L.M. ⁴	3.0	.7	.5	3.1	4.2	4.4	3.3	.1	3.5	.9
Yellow Tinged.....	.4		1.0	1.7	.3		.5	.2		.8
2-S.G.M.....										
3-G.M.....										
4-S.M.....	.1		.1		.1					
5-M. ⁴1		.4	.2	.1					
6-S.L.M. ⁴1		.3	.8	.1					
7-L.M. ⁴1		.2	.7						
Gray.....	.1	.1			.1					
3-G.M.....										
4-S.M.....		.1			.1					
5-M. ⁴1									
Blue Stained.....					.1					
3-G.M. ⁴										
4-S.M. ⁴1					
5-M. ⁴										
No grade ^{4,6}	1.9	.2	.7	.6	.4	.9	2.7		1.0	1.0

See footnotes at end of table.

TABLE 5.—*Grade of American upland cotton: Quantities ginned in the United States during specified periods,¹ by States, crops of 1932, 1931, 1930, 1929, and 1928—Continued*

TEXAS

Grade	Total, crop of—					Ginned prior to Oct. 1				
	1932	1931	1930	1929	1928	1932	1931	1930	1929	1928
All grades ²	<i>1,000 bales</i> 4,307.4	<i>1,000 bales</i> 5,068.8	<i>1,000 bales</i> 3,886.1	<i>1,000 bales</i> 3,803.2	<i>1,000 bales</i> 4,941.5	<i>1,000 bales</i> 1,595.2	<i>1,000 bales</i> 2,268.3	<i>1,000 bales</i> 2,584.7	<i>1,000 bales</i> 2,128.6	<i>1,000 bales</i> 2,430.9
Extra White	191.9	119.2	53.8	91.6	72.7	23.2	11.0	8.8	12.5	15.1
Above 3-G.M.	.3		.3			.3		.3		
3-G.M.	13.5	5.5	9.1	28.4	18.7	4.2	1.8	4.1	8.7	6.4
4-S.M.	36.9	28.5	24.9	25.0	27.1	8.6	1.8	4.3	3.6	5.3
5-M.	45.6	16.4	13.5	23.0	19.0	5.5	.8	.1	.2	1.7
6-S.L.M.	73.1	41.4	3.8	10.6	4.6	2.8	5.0			.4
7-L.M.	22.4	26.1	.6	4.6	3.3	1.8	1.6			1.3
Below 7-L.M. ⁴	.1	1.3	1.6							
White	2,887.4	4,436.5	3,588.2	3,303.5	4,278.8	1,418.9	2,150.6	2,465.6	1,999.2	2,340.9
1-M.F.				.1	.3					.2
2-S.G.M.	.5	1.5	9.1	8.1	29.4	.4	8	9.0	3.4	28.4
3-G.M.	100.0	396.7	407.0	254.1	803.6	75.1	206.1	395.1	216.7	712.4
4-S.M.	865.4	1,885.0	1,406.4	1,074.8	1,615.9	523.2	1,136.6	1,259.1	796.6	1,032.7
5-M.	1,196.4	1,295.5	1,021.1	1,224.2	1,115.9	584.0	587.9	633.6	705.3	439.9
6-S.L.M.	541.6	505.8	540.1	508.7	485.4	194.5	146.8	154.0	218.9	107.3
7-L.M.	137.8	231.6	168.1	168.1	148.8	34.3	17.2	11.5	43.8	20.5
8-S.G.O. ⁴	88.1	103.2	32.5	50.7	66.4	6.3	1.1	2.6	11.2	4.4
9-G.O. ⁴	7.6	16.6	3.3	14.7	13.1	1.1	.1	.7	3.3	1.1
Spotted	1,198.8	493.3	233.3	336.7	466.2	151.1	100.2	108.0	114.5	68.0
3-G.M.	127.7	46.5	31.0	10.3	48.7	16.5	22.7	28.9	4.4	14.0
4-S.M.	517.2	172.0	80.0	99.8	144.2	86.4	63.4	61.7	49.3	34.7
5-M.	348.0	119.3	66.6	135.7	142.4	45.2	13.1	16.7	48.6	17.0
6-S.L.M. ⁴	159.2	118.5	46.8	73.2	103.6	2.7	.9	.7	10.0	2.1
7-L.M. ⁴	46.7	37.0	8.9	17.7	27.3	.3	.1		2.2	.2
Yellow Tinged	24.9	9.7	6.3	46.2	14.6	1.4	.1	1.3	2.0	.5
2-S.G.M.						.1				.1
3-(I.M.)	1.7	.5	.3	.3	.2	.6		.2	.1	.1
4-S.M.	6.0	1.6	.9	5.7	.8	.5		.5	.6	.2
5-M. ⁴	7.1	4.2	2.6	17.8	2.2	.2	.1	.5	1.1	.1
6-S.L.M. ⁴	8.9	2.6	2.0	15.9	6.6	.1		.1		
7-L.M. ⁴	1.2	.8	.5	6.5	4.7					.1
Light Yellow Stained		.2	.3	10.3	.4			.1	.2	
3-G.M.					.1					
4-S.M. ⁴					.1			.1	.1	
5-M. ⁴		.2	.2	8.4	.2			.1		
Yellow Stained	.1		.1	5.3	.1					.1
3-G.M.						.1				
4-S.M. ⁴						.0				
5-M. ⁴	.1		.1	4.7						
Gray	1.7	5.4	3.1	.2	2.2	.4	.3	.8		.2
3-G.M.	.1	.2	.3		.2			.1		.1
4-S.M.	.9	2.9	2.3		.7	.3		.6		.1
5-M. ⁴	.7	2.3	.5	.2	1.3	.1	.2	.1		
No grade ^{4,6}	2.6	4.5	1.0	9.4	106.5	.2	.1	.1	.2	.1

See footnotes at end of table.

TABLE 5.—*Grade of American upland cotton: Quantities ginned in the United States during specified periods,¹ by States, crops of 1932, 1931, 1930, 1929, and 1928—Continued*

TEXAS—Continued

Grade	Ginned during October					Ginned during November				
	1932	1931	1930	1929	1928	1932	1931	1930	1929	1928
	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales
All grades ²	1,362.8	1,766.1	709.7	1,016.0	1,434.1	944.3	656.9	393.4	372.9	537.2
Extra White.....	98.9	32.1	19.1	39.2	27.7	62.0	62.2	14.7	33.6	20.9
Above 3-G.M.										
3-G.M.	6.8	3.5	4.6	19.6	9.6	2.5	.2	.4	.1	2.5
4-S.M.	14.9	16.4	11.7	14.0	12.3	13.3	9.7	7.4	5.9	7.7
5-M.	20.6	3.0	2.5	4.3	3.3	17.0	8.0	6.2	16.7	8.6
6-S.L.M.	44.4	5.2	.3	1.2	1.4	22.9	26.2	.7	7.5	1.5
7-L.M.	12.2	3.0		.1	1.1	6.3	18.1		3.4	.6
Below 7-L.M. ⁴		1.0								
White.....	1,068.4	1,582.4	659.8	893.0	1,277.9	322.3	457.0	343.6	204.1	411.3
1-M.F.				.1	.1					
2-S.G.M.		.6	.1	4.6	.9	.1	.1	.1	.2	
3-G.M.	20.6	120.5	10.3	35.2	83.3	4.3	7.8	1.5	1.2	5.9
4-S.M.	271.0	677.5	120.0	245.9	505.0	68.6	60.2	21.7	24.4	63.0
5-M.	488.5	524.7	278.1	420.0	487.3	115.8	163.5	90.8	79.4	145.2
6-S.L.M.	215.1	191.3	195.3	152.6	166.3	105.5	128.7	144.5	93.0	122.6
7-L.M.	55.5	57.8	51.4	30.3	28.8	22.1	70.0	70.2	52.0	47.3
8-S.G.O. ⁴	14.3	9.6	4.2	3.9	6.6	5.4	24.3	13.9	12.7	22.8
9-G.O. ⁴	.3	.4	.4	.5	1.6	.5	2.4	1.0	1.2	3.9
Spotted.....	193.1	147.3	28.1	81.6	125.8	553.5	134.8	33.5	64.0	101.8
3-G.M.	32.2	15.6	1.8	4.9	25.7	76.5	7.8	.2	.8	7.4
4-S.M.	102.1	67.8	9.9	33.7	64.9	291.5	37.7	4.6	9.5	31.0
5-M.	46.9	42.6	12.3	33.0	28.8	164.2	44.2	14.2	25.6	38.9
6-S.L.M. ⁴	9.5	19.2	3.8	8.6	5.9	19.7	38.5	12.0	23.3	18.9
7-L.M. ⁴	2.4	2.1	.3	1.3	.5	1.6	6.6	2.5	4.8	5.6
Yellow Tinged.....	.4	1.2	.7	1.0	.4	5.6	.8	.8	10.5	.4
2-S.G.M.										
3-G.M.	.1	.5	.1			1.0			.1	
4-S.M.	.2	.6	.1	.5	.1	3.7	.4	.1	1.7	.1
5-M. ⁴		.1	.3	1.0	.2	.8	.3	.5	3.2	.2
6-S.L.M. ⁴			.1	.3		.1	.1	.2	3.8	.1
7-L.M. ⁴	.1		.1	.1	.1				1.7	
Light Yellow Stained.....		.1	.1	.1	.2					.1
3-G.M.						.1				
4-S.M. ⁴						.1				
5-M. ⁴		.1	.1	.1						.1
Yellow Stained.....										.2
3-G.M.										
4-S.M. ⁴										
5-M. ⁴										.2
Gray.....	.4	2.7	1.8	.1	.9	.5	1.9	.6	.1	.7
3-G.M.										.1
4-S.M. ⁴										.1
5-M. ⁴										.5
No grade ^{4,6}	1.6	.3	.1	.2	1.2	.4	.2	.3	.4	2.0

See footnotes at end of table.

TABLE 5.—*Grade of American upland cotton: Quantities ginned in the United States during specified periods,¹ by States, crops of 1932, 1931, 1930, 1929, and 1928—Continued*

TEXAS—Continued

Grade	Ginned Dec. 1-Jan. 15, crop of—					Ginned after Jan. 15, crop of—										
	1932		1931		1930	1929		1928	1932		1931		1930	1929		
	<i>1,000 bales</i>		<i>1,000 bales</i>		<i>1,000 bales</i>	<i>1,000 bales</i>		<i>1,000 bales</i>	<i>1,000 bales</i>		<i>1,000 bales</i>		<i>1,000 bales</i>	<i>1,000 bales</i>		
All grades ²	262.0		256.7		166.5	240.9		403.8	143.1		120.8		31.8	44.8		135.5
Extra White.....		5.1		12.2		10.0		5.6	8.2		2.7		1.7	1.2		.7
Above 3-G.M.																
3-G.M.																
4-S.M.																
5-M.																
6-S.L.M.																
7-L.M.																
Below 7-L.M. ⁴																
White.....		54.4		171.1		105.6		124.8	214.2		23.4		69.4	13.6		22.4
1-M.F.																
2-S.G.M.																
3-G.M.																
4-S.M.																
5-M.																
6-S.L.M.																
7-L.M.																
Spotted.....		192.4		70.2		47.7		64.7	126.3		108.7		40.8	16.0		12.0
3-G.M.																
4-S.M.																
5-M.																
6-S.L.M. ⁴																
7-L.M. ⁴																
Yellow Tinged.....																
2-S.G.M.																
3-G.M.																
4-S.M.																
5-M.																
6-S.L.M. ⁴																
7-L.M. ⁴																
Light Yellow Stained.....																
3-G.M.																
4-S.M. ⁴																
5-M. ⁴																
Yellow Stained.....																
3-G.M.																
4-S.M. ⁴																
5-M. ⁴																
Gray.....																
3-G.M.																
4-S.M.																
5-M. ⁴																
No grade ^{4,6}																

See footnotes at end of table.

TABLE 5.—Grade of American upland cotton: Quantities ginned in the United States during specified periods,¹ by States, crops of 1932, 1931, 1930, 1929, and 1928—Continued

VIRGINIA

Grade	Total, crop of—					Ginned prior to Oct. 1				
	1932	1931	1930	1929	1928	1932	1931	1930	1929	1928
All grades ²	<i>1,000 bales</i> 31.3	<i>1,000 bales</i> 42.5	<i>1,000 bales</i> 42.7	<i>1,000 bales</i> 48.0	<i>1,000 bales</i> 44.8	<i>1,000 bales</i> 6.1	<i>1,000 bales</i> 6.2	<i>1,000 bales</i> 7.9	<i>1,000 bales</i> 0.3	<i>1,000 bales</i> 1.0
White	28.4	40.9	41.6	44.3	37.8	6.1	5.8	7.7	.3	.8
1-M.F.										
2-S.G.M.										
3-G.M.			.1	.4	.1					
4-S.M.	4.5	1.1	8.7	10.2	13.1	2.4	.5	3.4	.1	.3
5-M.	10.8	27.1	21.0	26.1	17.4	3.4	4.8	4.0	.2	.5
6-S.L.M.	7.0	12.4	7.3	6.9	4.7	.3	.5	.2		
7-L.M.	.1	.3	4.3	.7	.9			.1		
8-S.G.O. ⁴			.2		.3					
9-G.O. ⁴					.1					
Spotted	2.0	1.5	1.1	3.3	6.1		.4	.2		.2
3-G.M.						.4				
4-S.M.	.7	1.4	.6	1.8	3.5		.4			.1
5-M.	2.1	.1	.4	1.4	2.0			.1		.1
6-S.L.M. ⁴	.1		.1	.1	.2			.1		
7-L.M. ⁴										
Yellow Tinged					.1	.4				
2-S.G.M.						.1				
3-G.M.						.2				
4-S.M.					.1	.1				
5-M. ⁴										
6-S.L.M. ⁴										
7-L.M. ⁴										
Gray		.1		.3	.3					
3-G.M.						.1				
4-S.M.		.1		.1	.2					
5-M.				.2						
Blue Stained						.2				
3-G.M. ⁴						.1				
4-S.M. ⁴						.1				
5-M. ⁴										
Grade	Ginned during October					Ginned during November				
	1932	1931	1930	1929	1928	1932	1931	1930	1929	1928
All grades ²	<i>1,000 bales</i> 11.2	<i>1,000 bales</i> 22.1	<i>1,000 bales</i> 22.2	<i>1,000 bales</i> 16.8	<i>1,000 bales</i> 10.7	<i>1,000 bales</i> 9.0	<i>1,000 bales</i> 11.3	<i>1,000 bales</i> 9.1	<i>1,000 bales</i> 18.0	<i>1,000 bales</i> 15.9
White	11.1	21.3	21.5	16.3	17.6	8.2	11.1	9.0	17.6	14.3
1-M.F.										
2-S.G.M.						.1				
3-G.M.			.1	.2	.5				.2	.7
4-S.M.	2.0	.5	5.0	5.1	7.7	.1	.1	.3	4.2	4.5
5-M.	8.3	13.7	13.4	9.9	8.2	4.5	6.1	3.0	12.0	2.0
6-S.L.M.	.8	6.8	2.9	1.0	1.1	3.6	4.0	3.4	1.2	.1
7-L.M.	.3	.1	.1	.1				2.3		
8-S.G.O. ⁴										
9-G.O. ⁴										
Spotted	.1	.8	.7	.5	2.1	.8	.1	.1	1.0	1.5
3-G.M.						.1				.2
4-S.M.	.1	.7	.6	.3	.9	.2	.1		.8	1.2
5-M.		.1	.1	.2	1.0	.0		.1	.2	.1
6-S.L.M. ⁴										
7-L.M. ⁴										

See footnotes at end of table.

TABLE 5.—*Grade of American upland cotton: Quantities ginned in the United States during specified periods,¹ by States, crops of 1932, 1931, 1930, 1929, and 1928—Continued*

VIRGINIA—Continued

Grade	Ginned during October					Ginned during November				
	1932	1931	1930	1929	1928	1932	1931	1930	1929	1928
	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales
Yellow Tinged										.1
2-S.G.M.										
3-G.M.										
4-S.M.										.1
5-M. ⁴										
6-S.L.M. ⁴										
7-L.M. ⁴										
Gray									.1	
3-G.M.										
4-S.M.									.1	
5-M. ⁴										
Blue Stained										
3-G.M. ⁴										
4-S.M. ⁴										
5-M. ⁴										
Grade	Ginned Dec. 1-Jan. 15, crop of—					Ginned after Jan. 15, crop of—				
	1932	1931	1930	1929	1928	1932	1931	1930	1929	1928
	1,000 bales 3.7	1,000 bales 2.2	1,000 bales 2.4	1,000 bales 10.2	1,000 bales 6.0	1,000 bales 1.3	1,000 bales 0.7	1,000 bales 1.1	1,000 bales 2.1	1,000 bales 2.2
White	2.4	2.1	2.3	8.5	3.2	.6	.6	1.1	1.6	1.9
1-M.F.										
2-S.G.M.										
3-G.M.										
4-S.M.										
5-M. ⁴	.4	1.9	.8	3.5	1.3	.2	.6		.5	.4
6-S.L.M.	1.9	.2	.6	3.7	1.1	.4		.2	1.0	.5
7-L.M.	.1		1.0	.6	.1			.8	.1	.7
8-S.G.O. ⁴			.1		.1			.1		.2
9-G.O. ⁴										.1
Spotted	1.3	.1	.1	1.4	2.2	.7	.1		.4	.1
3-G.M.										
4-S.M.	.2	.1		.7	1.3	.2	.1			
5-M. ⁴	1.1		.1	.6	.7	.4		.4		.1
6-S.L.M. ⁴				.1	.1	.1				
7-L.M. ⁴										
Yellow Tinged										
2-S.G.M.										
3-G.M.										
4-S.M.										
5-M. ⁴										
6-S.L.M. ⁴										
7-L.M. ⁴										
Gray										
3-G.M.										
4-S.M.										
5-M. ⁴										
Blue Stained										
3-G.M. ⁴										
4-S.M. ⁴										
5-M. ⁴										

See footnotes at end of table.

TABLE 5.—*Grade of American upland cotton: Quantities ginned in the United States during specified periods,¹ by States, crops of 1932, 1931, 1930, 1929, and 1928—Continued*ALL OTHER STATES²

Grade	Total, crop of—					Ginned prior to Oct. 1				
	1932	1931	1930	1929	1928	1932	1931	1930	1929	1928
All grades ³	1,000 bales 14.0	1,000 bales 11.7	1,000 bales 6.4	1,000 bales 8.9	1,000 bales 6.2	1,000 bales 2.0	1,000 bales 0.6	1,000 bales 1.6	1,000 bales 0.5	1,000 bales 0.2
Extra White.....	.1									
Above 3-G.M.....										
3-G.M.....										
4-S.M.....										
5-M.....										
6-S.L.M.....										
7-L.M.....	.1									
Below 7-L.M. ⁴										
White.....	11.1	9.3	5.8	6.8	4.5	1.8	.6	1.6	.5	.2
1-M.F.....										
2-S.G.M.....										
3-G.M.....			.2							
4-S.M.....	2.1	1.8	2.0	.5	.6	.6	.4	.8	.2	.1
5-M.....	5.0	3.1	2.5	2.5	.8	1.0	.2	.6	.2	.1
6-S.L.M.....	2.2	2.5	.6	1.7	1.2	.2		.1		
7-L.M.....	.4	.5	.2	.4	.5					
8-S.G.O.....	.4	.2	.2	1.0	.5					
9-G.O. ⁴	1.0	1.2	.1	.7	.9					
Spotted.....	1.7	.4	.1	.8	.5	.2				
3-G.M.....	.1						.1			
4-S.M.....	.5	.1					.1			
5-M.....	.5	.2		.2	.2					
6-S.L.M. ⁴2	.1	.1	.3	.2					
7-L.M. ⁴4			.3	.1					
Yellow Tinged.....				.2						
2-S.G.M.....										
3-G.M.....										
4-S.M.....										
5-M ⁴										
6-S.L.M. ⁴1						
7-L.M. ⁴1						
No grade ⁴	1.1	2.0	.6	1.1	1.2					
 Ginned during October										
Grade	1932	1931	1930	1929	1928	1932	1931	1930	1929	1928
All grades ²	1,000 bales 5.5	1,000 bales 3.9	1,000 bales 2.7	1,000 bales 2.9	1,000 bales 1.5	1,000 bales 2.7	1,000 bales 3.3	1,000 bales 1.2	1,000 bales 2.3	1,000 bales 1.4
Extra White.....										
Above 3-G.M.....										
3-G.M.....										
4-S.M.....										
5-M.....										
6-S.L.M.....										
7-L.M.....										
Below 7-L.M. ⁴										
White.....	5.1	3.8	2.7	2.8	1.4	2.5	3.2	1.2	2.0	1.3
1-M.F.....										
2-S.G.M.....										
3-G.M.....			.1							
4-S.M.....	.9	1.3	1.1	.3	.4	.6		.1		.1
5-M.....	3.0	1.8	1.4	1.6	.4	1.0	.8	.4	.7	.8
6-S.L.M.....	1.0	.6	.1	.8	.3	.6	1.9	.4	.2	.1
7-L.M.....	.2	.1		.1	.2	.1	.4	.2		
8-S.G.O.....					.1	.1	.1	.1	.3	
9-G.O. ⁴1				

See footnotes at end of table.

TABLE 5.—*Grade of American upland cotton: Quantities ginned in the United States during specified periods,¹ by States, crops of 1932, 1931, 1930, 1929, and 1928—Continued*ALL OTHER STATES²—Continued

Grade	Ginned during October					Ginned during November				
	1932	1931	1930	1929	1928	1932	1931	1930	1929	1928
Spotted	1,000 bales 0.4	1,000 bales 0.1	1,000 bales	1,000 bales 0.1	1,000 bales 0.1	1,000 bales 0.1	1,000 bales 0.1	1,000 bales	1,000 bales 0.3	1,000 bales 0.1
3-G.M.	.2	.1								
4-S.M.	.2									
5-M.				.1	.1		.1			.1
6-S.L.M. ⁴										
7-L.M. ⁴										.2
Yellow Tinged										
2-S.G.M.										
3-G.M.										
4-S.M.										
5-M. ⁴										
6-S.L.M. ⁴										
7-L.M. ⁴										
No grade ^{5,6}										

Grade	Ginned Dec. 1-Jan. 15, crop of—					Ginned after Jan. 15, crop of—				
	1932	1931	1930	1929	1928	1932	1931	1930	1929	1928
All grades ¹	1,000 bales 2.6	1,000 bales 2.0	1,000 bales 0.6	1,000 bales 1.4	1,000 bales 1.8	1,000 bales 1.2	1,000 bales 1.0	1,000 bales 0.3	1,000 bales 1.8	1,000 bales 1.3
Extra White										
Above 3-G.M.										
3-G.M.										
4-S.M.										
5-M.										
6-S.L.M.										
7-L.M.										
Below 7-L.M. ⁴										
White	1.4	1.3	.2	.5	1.1	.3	.4	.1	1.0	.5
1-M.F.										
2-S.G.M.										
3-G.M.										
4-S.M.										
5-M.										
6-S.L.M.	.4					.1	.1			
7-L.M.	.1					.1	.2			
8-S.G.O. ⁴	.3					.2	.2			
9-G.O. ⁴	.6	.9				.1	.3	.1	.5	.3
Spotted	.6	.2	.1	.3	.2	.4				
3-G.M.										
4-S.M.										
5-M.	.2		.1							
6-S.L.M. ⁴	.1	.1	.1							
7-L.M. ⁴	.2					.3	.1			
Yellow Tinged						.1				
2-S.G.M.										
3-G.M.										
4-S.M.										
5-M. ⁴										
6-S.L.M. ⁴										
7-L.M. ⁴										
No grade ^{5,6}	.6	.6	.3	.5	.5	.5	1.5	.2	.6	.7

¹ Periods are those used by the Bureau of the Census.² As reported by the Bureau of the Census.³ Included with Strict Good Middling, White.⁴ Untenderable in settlement of futures contracts made subject to sec. 5 of the United States Cotton Futures Act and the regulations of the Secretary of Agriculture thereunder.⁵ Included with Strict Good Ordinary, White.⁶ Includes bales not otherwise classified above.⁷ Includes Illinois, Kansas, and Kentucky.

TABLE 6.—*Grade of American-Egyptian cotton: Quantities ginned in the United States during specified periods,¹ crops of 1932, 1931, 1930, 1929, and 1928*

[Quantities are given in running bales]

Period and grade	Crop of	Crop of	Crop of	Crop of	Crop of
	1932 1,000 bales 8.3	1931 1,000 bales 13.7	1930 1,000 bales 23.3	1929 1,000 bales 28.8	1928 1,000 bales 28.3
All periods and grades ²					
1 and 1½.....	2.0	1.2	6.2	5.5	5.5
2 and 2½.....	3.7	5.9	11.4	16.4	13.8
3 and 3½.....	2.2	4.5	5.1	6.5	8.4
4 and 4½.....	.4	1.9	.6	.4	.7
5.....		.2			.1
Below 5.....					
Prior to Oct. 1.....	1.1	2.3	3.9	3.4	0.2
1 and 1½.....	1.0	.6	1.5	.7	3.8
2 and 2½.....	.1	1.4	2.3	2.6	.22
3 and 3½.....		.3	.1	.1	.4
4 and 4½.....					
5.....					
Below 5.....					
During October.....	2.7	3.1	6.6	7.4	7.2
1 and 1½.....	.7	.3	2.4	2.8	.9
2 and 2½.....	1.5	1.7	4.1	4.2	4.4
3 and 3½.....	.5	.7	.1	.4	1.8
4 and 4½.....		.3			
5.....		.1			
Below 5.....					
During November.....	2.4	1.4	5.0	7.2	0.4
1 and 1½.....	.3	.3	2.2	1.3	.8
2 and 2½.....	1.4	.8	2.2	4.3	3.4
3 and 3½.....	.7	.2	.6	1.5	2.1
4 and 4½.....		.1		.1	.1
5.....					
Below 5.....					
Dec. 1-Jan. 15.....	1.2	4.1	5.7	7.5	5.6
1 and 1½.....					
2 and 2½.....	.6	1.0	2.2	4.5	3.2
3 and 3½.....	.5	2.0	3.2	2.2	1.0
4 and 4½.....	.1	1.0	.2	.1	.3
5.....		.1			
Below 5.....					
After Jan. 15.....	.9	2.8	2.1	3.3	2.9
1 and 1½.....					
2 and 2½.....	.1	1.0	.6	.8	.4
3 and 3½.....	.5	1.3	1.1	2.3	2.2
4 and 4½.....	.3	.5	.4	.2	.2
5.....					
Below 5.....					

¹ Periods are those used by the Bureau of the Census.² As reported by the Bureau of the Census.

TABLE 7.—*Staple length of American upland cotton: Quantities ginned in the United States during specified periods,¹ by States, crops of 1932, 1931, 1930, 1929, and 1928*

[Quantities are given in running bales, except that round bales are counted as half bales. Linters are not included]

UNITED STATES

Staple length (inches)	Total, crop of—					Ginned prior to Oct. 1				
	1932	1931	1930	1929	1928	1932	1931	1930	1929	1928
	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales
All lengths ²	12,701.3	16,615.2	13,732.2	14,519.0	14,268.2	4,834.9	5,407.4	6,300.0	5,899.9	4,955.2
Shorter than $\frac{7}{8}$ ³	837.7	1,019.5	1,829.2	2,921.5	2,072.1	341.8	175.9	882.6	926.7	651.9
$\frac{7}{8}$ and $2\frac{1}{2}$ ₃₂	4,780.5	6,593.3	5,327.7	5,533.7	5,914.8	1,882.3	2,083.6	2,382.4	1,148.4	1,029.7
$\frac{1}{2}$ and $3\frac{1}{2}$ ₂	3,671.0	4,511.9	3,421.6	2,748.2	3,225.7	1,371.8	1,897.7	1,636.6	1,297.2	1,218.4
1 and $1\frac{1}{2}$ ₂	1,822.0	2,557.1	1,725.9	1,693.6	1,575.8	646.9	975.1	900.0	704.1	640.0
$1\frac{1}{2}$ and $1\frac{3}{4}$ ₂	871.8	1,087.8	970.9	938.6	794.2	346.6	262.8	317.2	345.5	232.2
$1\frac{1}{2}$ and $1\frac{1}{2}$ ₂	622.1	590.0	393.3	556.1	489.2	222.4	125.8	151.3	209.3	165.7
$1\frac{1}{2}$ and $1\frac{3}{4}$ ₂	84.5	224.6	60.8	119.4	167.9	21.6	86.0	28.8	83.8	96.8
$1\frac{1}{2}$ and longer.....	5.7	31.0	2.8	7.9	28.5	1.5	20.5	1.1	4.9	20.5
 Ginned during October										
Staple length (inches)	1932	1931	1930	1929	1928	1932	1931	1930	1929	1928
	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales
	4,408.3	6,711.5	4,553.4	4,981.3	5,193.9	2,385.7	2,892.7	1,968.2	1,954.0	2,391.3
All lengths ²	207.0	455.0	654.0	1,293.7	819.6	91.7	207.8	209.9	396.0	211.9
$\frac{7}{8}$ and $2\frac{1}{2}$ ₃₂	1,554.0	2,854.4	1,869.0	1,913.0	2,136.1	932.4	1,120.9	734.6	809.3	1,136.4
$\frac{1}{2}$ and $3\frac{1}{2}$ ₂	1,321.7	1,679.4	1,040.6	794.8	1,203.0	733.3	748.5	503.8	336.2	404.4
1 and $1\frac{1}{2}$ ₂	701.9	904.2	471.7	484.8	551.2	351.9	405.0	224.3	252.9	247.5
$1\frac{1}{2}$ and $1\frac{3}{4}$ ₂	301.6	439.2	341.5	318.9	245.1	160.5	222.4	208.5	113.9	180.6
$1\frac{1}{2}$ and $1\frac{1}{2}$ ₂	271.1	273.5	142.6	160.5	181.7	102.9	148.7	80.3	38.3	101.8
$1\frac{1}{2}$ and $1\frac{3}{4}$ ₂	47.5	97.0	23.6	24.1	50.8	12.1	37.9	6.5	6.4	17.3
$1\frac{1}{2}$ and longer.....	2.9	8.8	1.4	1.9	6.4	.9	1.5	.3	1.0	1.4
 Ginned Dec. 1-Jan. 15, crop of—										
Staple length (inches)	1932	1931	1930	1929	1928	1932	1931	1930	1929	1928
	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales
	778.6	973.9	751.6	1,316.3	1,323.1	293.8	629.7	159.0	367.5	404.7
All lengths ²	82.5	94.0	67.7	250.5	257.9	114.1	86.8	15.0	48.6	130.8
Shorter than $\frac{7}{8}$ ³	334.4	415.9	277.0	535.8	594.0	83.4	138.5	63.8	127.2	118.6
$\frac{7}{8}$ and $2\frac{1}{2}$ ₃₂	191.9	255.0	157.2	241.0	244.9	52.3	131.3	44.4	79.0	65.0
$\frac{1}{2}$ and $3\frac{1}{2}$ ₂	92.9	115.4	108.4	132.7	95.0	28.4	157.4	21.5	45.5	42.1
1 and $1\frac{1}{2}$ ₂	50.1	68.4	61.9	107.3	97.0	13.0	96.0	11.8	53.0	39.3
$1\frac{1}{2}$ and $1\frac{3}{4}$ ₂	23.4	22.0	16.8	38.5	31.5	2.3	19.4	2.3	13.5	8.5
$1\frac{1}{2}$ and $1\frac{1}{2}$ ₂	3.0	2.5	1.7	4.4	2.6	.3	1.2	.2	.7	.4
$1\frac{1}{2}$ and longer.....	.4	.1	—	.1	.2	—	.1	—	—	—

See footnotes at end of table (p. 84).

TABLE 7.—*Staple length of American upland cotton: Quantities ginned in the United States during specified periods,¹ by States, crops of 1932, 1931, 1930, 1929, and 1928—Continued*

ALABAMA

Staple length (inches)	Total, crop of—					Ginned prior to Oct. 1				
	1932	1931	1930	1929	1928	1932	1931	1930	1929	1928
All lengths ²	1,000 bales 933.7	1,000 bales 1,385.0	1,000 bales 1,444.9	1,000 bales 1,307.7	1,000 bales 1,096.6	1,000 bales 412.0	1,000 bales 531.0	1,000 bales 582.6	1,000 bales 578.1	1,000 bales 336.0
Shorter than $\frac{7}{8}$ ³	137.3	136.5	553.3	586.6	264.2	70.4	19.5	252.3	187.8	68.2
$\frac{7}{8}$ and $2\frac{1}{4}$	657.5	1,044.7	802.2	689.0	762.6	205.5	430.4	310.3	369.2	251.8
$\frac{1}{2}\frac{1}{2}$ and $3\frac{1}{2}$	108.7	168.7	77.1	28.3	49.8	35.6	67.8	15.7	18.0	22.2
1 and $1\frac{1}{2}$	26.5	28.0	9.2	3.3	17.6	9.4	10.2	2.9	1.8	2.6
$1\frac{1}{4}$ and $1\frac{3}{4}$	3.1	5.8	2.5	1.4	1.8	1.1	2.4	1.2	1.2	.9
$1\frac{1}{4}$ and $1\frac{5}{8}$5	1.3	.6	.1	.5	.1	.7	.2	.1	.3
$1\frac{1}{4}$ and $1\frac{1}{2}$1	—	—	—	.1	—	—	—	—	—
1 $\frac{1}{4}$ and longer.....	—	—	—	—	—	—	—	—	—	—

Staple length (inches)	Ginned during October					Ginned during November				
	1932	1931	1930	1929	1928	1932	1931	1930	1929	1928
All lengths ²	1,000 bales 330.0	1,000 bales 647.9	1,000 bales 595.4	1,000 bales 511.8	1,000 bales 482.0	1,000 bales 160.6	1,000 bales 185.8	1,000 bales 202.0	1,000 bales 109.0	1,000 bales 206.2
Shorter than $\frac{7}{8}$ ³	55.0	74.4	230.0	280.7	149.4	11.0	41.3	59.0	62.8	43.5
$\frac{7}{8}$ and $2\frac{1}{4}$	216.7	472.1	321.6	222.1	208.6	114.6	126.9	121.1	45.2	165.8
$\frac{1}{2}\frac{1}{2}$ and $3\frac{1}{2}$	44.3	83.6	38.8	7.6	18.3	20.9	14.5	19.1	1.7	6.5
1 and $1\frac{1}{2}$	12.2	14.1	3.9	1.2	14.5	3.6	3.0	2.2	.2	.4
$1\frac{1}{4}$ and $1\frac{3}{4}$	1.5	3.1	1.0	.2	.9	.4	.1	.3	—	—
$1\frac{1}{4}$ and $1\frac{5}{8}$3	.6	.1	—	.2	.1	—	.3	—	—
$1\frac{1}{4}$ and $1\frac{1}{2}$	—	—	—	—	.1	—	—	—	—	—
1 $\frac{1}{4}$ and longer.....	—	—	—	—	—	—	—	—	—	—

Staple length (inches)	Ginned Dec. 1-Jan. 15, crop of—					Ginned after Jan. 15, crop of—				
	1932	1931	1930	1929	1928	1932	1931	1930	1929	1928
All lengths ²	1,000 bales 35.3	1,000 bales 16.0	1,000 bales 60.9	1,000 bales 95.5	1,000 bales 63.1	1,000 bales 5.8	1,000 bales 3.4	1,000 bales 8.0	1,000 bales 12.4	1,000 bales 9.3
Shorter than $\frac{7}{8}$ ³7	1.1	11.3	40.4	11.7	.2	.2	.7	4.9	1.4
$\frac{7}{8}$ and $2\frac{1}{4}$	26.5	13.2	42.2	45.2	48.8	4.2	2.1	7.0	7.3	7.6
$\frac{1}{2}\frac{1}{2}$ and $3\frac{1}{2}$	6.8	1.8	3.2	.8	2.5	1.2	1.0	.3	.2	.3
1 and $1\frac{1}{2}$	1.2	.6	.2	.1	.1	.1	.1	—	—	—
$1\frac{1}{4}$ and $1\frac{3}{4}$1	.2	—	—	—	—	—	—	—	—
$1\frac{1}{4}$ and $1\frac{5}{8}$	—	—	—	—	—	—	—	—	—	—
$1\frac{1}{4}$ and $1\frac{1}{2}$	—	—	—	—	—	—	—	—	—	—
1 $\frac{1}{4}$ and longer.....	—	—	—	—	—	.1	—	—	—	—

ARIZONA

Staple length (inches)	Total, crop of—					Ginned prior to Oct. 1				
	1932	1931	1930	1929	1928	1932	1931	1930	1929	1928
All lengths ²	1,000 bales 58.8	1,000 bales 97.3	1,000 bales 127.2	1,000 bales 120.7	1,000 bales 117.4	1,000 bales 9.8	1,000 bales 10.2	1,000 bales 10.6	1,000 bales 15.1	1,000 bales 21.0
Shorter than $\frac{7}{8}$ ³1	3.3	.3	1.5	2.6	—	—	—	—	.9
$\frac{7}{8}$ and $2\frac{1}{4}$	1.4	.6	5.5	4.3	11.3	—	—	—	—	3.5
$\frac{1}{2}\frac{1}{2}$ and $3\frac{1}{2}$	6.9	14.9	41.2	27.4	17.6	.6	2.4	1.7	2.7	4.2
1 and $1\frac{1}{2}$	23.5	43.4	47.8	65.3	40.2	2.5	3.2	10.9	8.5	6.4
$1\frac{1}{4}$ and $1\frac{3}{4}$	20.3	30.8	29.6	19.4	42.7	4.6	3.4	6.0	2.2	5.3
$1\frac{1}{4}$ and $1\frac{5}{8}$	6.3	4.3	2.8	2.8	3.0	2.1	.9	1.0	1.1	.7
$1\frac{1}{4}$ and $1\frac{1}{2}$3	—	—	—	—	—	—	—	—	—
1 $\frac{1}{4}$ and longer.....	—	—	—	—	—	—	—	—	—	—

See footnotes at end of table.

TABLE 7.—*Staple length of American upland cotton: Quantities ginned in the United States during specified periods,¹ by States, crops of 1932, 1931, 1930, 1929, and 1928—Continued*

ARIZONA—Continued

Staple length (inches)	Ginned during October					Ginned during November				
	1932	1931	1930	1929	1928	1932	1931	1930	1929	1928
	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales
All lengths ²	14.7	20.3	31.4	34.6	27.2	15.8	16.6	28.0	35.5	27.4
Shorter than $\frac{7}{8}$ ³					1.6	.1				
$\frac{7}{8}$ and $\frac{29}{32}$2									.9
$\frac{7}{8}$ and $\frac{3}{4}$	1.1	3.6	7.2	4.1	7.2	1.6	.7	6.9	8.3	2.8
1 and $\frac{1}{3}$	4.0	8.7	12.6	18.2	6.4	7.2	8.3	10.1	23.5	11.0
$\frac{1}{4}$ and $\frac{13}{16}$	6.2	6.7	10.5	10.2	6.3	5.6	6.8	10.2	3.7	11.8
$\frac{1}{4}$ and $\frac{15}{16}$	3.0	1.3	1.0	1.7	.4	1.2	.8	.8		.8
$\frac{3}{4}$ and $\frac{17}{16}$2					.1				
1 and longer.....										

Staple length (inches)	Ginned Dec. 1-Jan. 15, crop of—					Ginned after Jan. 15, crop of—				
	1932	1931	1930	1929	1928	1932	1931	1930	1929	1928
	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales
All lengths ²	12.5	31.9	38.3	28.8	27.7	6.0	18.3	9.9	5.7	14.1
Shorter than $\frac{7}{8}$ ³2	.1	.9				3.1	.2	.6	.1
$\frac{7}{8}$ and $\frac{29}{32}$2	.3	3.1	1.9		1.0		2.3	1.4	.6
$\frac{7}{8}$ and $\frac{3}{4}$	2.6	6.5	19.4	10.7	1.1	1.0	1.7	6.0	1.6	2.3
1 and $\frac{1}{3}$	6.7	15.5	13.0	13.2		10.0	3.1	7.7	1.2	6.4
$\frac{1}{4}$ and $\frac{13}{16}$	3.0	8.5	2.7	3.1	15.6	.9	5.4	.2	.2	4.6
$\frac{1}{4}$ and $\frac{15}{16}$9				1.0		.4			.1
$\frac{3}{4}$ and $\frac{17}{16}$										
1 and longer.....										

ARKANSAS

Staple length (inches)	Total, crop of—					Ginned prior to Oct. 1				
	1932	1931	1930	1929	1928	1932	1931	1930	1929	1928
	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales
All lengths ²	1,283.4	1,836.1	863.5	1,395.9	1,218.2	501.8	268.6	264.2	539.0	361.8
Shorter than $\frac{7}{8}$ ³	25.8	20.9	76.9	177.9	127.0	12.1	4.1	25.8	84.7	51.1
$\frac{7}{8}$ and $\frac{29}{32}$	185.9	307.4	196.1	393.4	359.4	88.2	53.3	64.5	161.4	122.2
$\frac{7}{8}$ and $\frac{3}{4}$	375.0	549.6	252.9	379.1	290.5	160.9	77.2	71.5	124.4	79.7
1 and $\frac{1}{3}$	360.4	614.2	206.8	261.9	223.6	131.8	78.4	66.9	96.2	56.5
$\frac{1}{4}$ and $\frac{13}{16}$	196.8	247.0	106.6	128.0	130.6	70.5	30.7	28.8	49.1	31.3
$\frac{1}{4}$ and $\frac{15}{16}$	120.4	81.4	22.4	45.4	65.9	35.2	15.6	6.1	19.5	15.6
$\frac{3}{4}$ and $\frac{17}{16}$	17.3	14.9	1.7	8.6	16.9	3.0	3.1	.5	3.3	4.7
1 and longer.....	1.8	.7	.1	1.6	2.3	.1	.2	.1	.4	.7

Staple length (inches)	Ginned during October					Ginned during November				
	1932	1931	1930	1929	1928	1932	1931	1930	1929	1928
	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales
All lengths ²	481.0	766.1	368.0	523.4	399.3	181.0	438.3	182.3	180.1	259.1
Shorter than $\frac{7}{8}$ ³	11.2	10.6	41.5	73.3	55.5	1.8	4.9	7.9	16.0	10.5
$\frac{7}{8}$ and $\frac{29}{32}$	58.5	130.7	88.8	151.1	127.1	23.2	68.1	33.4	56.8	61.2
$\frac{7}{8}$ and $\frac{3}{4}$	122.7	198.7	100.5	148.1	78.1	49.0	131.5	58.7	59.7	68.6
1 and $\frac{1}{3}$	143.0	254.0	75.7	92.5	70.5	54.4	153.7	50.5	35.6	57.3
$\frac{1}{4}$ and $\frac{13}{16}$	78.4	116.0	43.5	38.2	34.5	29.3	50.0	27.9	15.8	38.6
$\frac{1}{4}$ and $\frac{15}{16}$	56.0	42.2	11.7	16.0	24.4	19.9	17.0	3.7	3.7	18.9
$\frac{3}{4}$ and $\frac{17}{16}$	11.2	7.7	.9	3.4	8.0	2.3	3.8	.2	1.1	3.6
1 and longer.....	1.1	.2		.8	1.2	.5	.3		.4	.4

See footnotes at end of table.

TABLE 7.—*Staple length of American upland cotton: Quantities ginned in the United States during specified periods,¹ by States, crops of 1932, 1931, 1930, 1929, and 1928—Continued*

ARKANSAS—Continued

Staple length (inches)	Ginned Dec. 1-Jan. 15, crop of—					Ginned after Jan. 15, crop of—				
	1932	1931	1930	1929	1928	1932	1931	1930	1929	1928
	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales
All lengths ²	89.2	102.9	43.0	99.2	136.0	30.4	200.2	4.5	45.2	60.0
Shorter than $\frac{7}{8}$ ³7	1.0	1.6	3.5	8.8	—	—	.3	.1	.4
$\frac{7}{8}$ and $\frac{29}{32}$	13.2	36.1	8.2	16.9	36.1	2.8	13.2	1.2	7.2	12.8
$\frac{7}{8}$ and $\frac{31}{32}$	29.1	78.7	14.4	31.7	44.0	12.7	63.5	1.8	15.2	20.1
1 and $1\frac{1}{2}$	22.1	34.3	13.0	25.0	25.9	8.2	93.8	.7	12.6	13.4
$\frac{11}{16}$ and $1\frac{3}{8}$	14.9	9.6	5.9	16.6	15.4	5.7	25.7	.5	8.3	10.8
$\frac{13}{16}$ and $1\frac{5}{8}$	8.5	3.0	.7	4.8	5.3	.8	3.6	.2	1.4	1.7
$\frac{15}{16}$ and $1\frac{7}{8}$6	.2	.1	.7	.5	.2	.1	—	.1	.1
$1\frac{1}{4}$ and longer.....	.1	—	—	—	—	—	—	—	—	—

CALIFORNIA

Staple length (inches)	Total, crop of—					Ginned prior to Oct. 1				
	1932	1931	1930	1929	1928	1932	1931	1930	1929	1928
	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales
All lengths ²	124.4	171.2	256.3	254.1	171.0	7.4	27.8	15.4	13.5	20.3
Shorter than $\frac{7}{8}$ ³	—	8.9	—	.8	2.8	—	—	—	—	.5
$\frac{7}{8}$ and $\frac{29}{32}$	2.2	.1	1.7	5.0	20.6	—	—	—	.2	2.0
$\frac{7}{8}$ and $\frac{31}{32}$	3.9	2.4	13.9	17.1	29.3	.1	.2	1.3	2.0	—
1 and $1\frac{1}{2}$	13.4	25.6	41.2	119.6	47.6	.6	5.8	2.4	9.0	9.9
$\frac{11}{16}$ and $1\frac{3}{8}$	57.9	98.0	160.0	104.2	60.0	2.1	19.6	11.8	2.8	4.4
$\frac{13}{16}$ and $1\frac{5}{8}$	46.8	33.9	39.2	7.4	10.8	4.5	2.2	1.2	.2	.6
$\frac{15}{16}$ and $1\frac{7}{8}$2	2.3	.3	—	—	.1	—	—	—	—
$1\frac{1}{4}$ and longer.....	—	—	—	—	—	—	—	—	—	—

Staple length (inches)	Ginned during October					Ginned during November				
	1932	1931	1930	1929	1928	1932	1931	1930	1929	1928
	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales
All lengths ²	53.6	66.6	75.3	87.4	61.3	45.4	38.9	72.3	78.0	43.2
Shorter than $\frac{7}{8}$ ³	—	.3	—	.1	2.1	—	—	—	.3	.2
$\frac{7}{8}$ and $\frac{29}{32}$1	—	—	.1	15.8	.1	—	—	—	1.8
$\frac{7}{8}$ and $\frac{31}{32}$8	.4	1.3	1.8	18.4	1.0	.1	.8	0.7	4.1
1 and $1\frac{1}{2}$	1.0	8.9	8.3	35.5	7.3	4.9	3.0	4.2	55.9	16.4
$\frac{11}{16}$ and $1\frac{3}{8}$	21.5	41.1	55.7	43.9	18.1	27.8	21.2	44.8	15.1	18.1
$\frac{13}{16}$ and $1\frac{5}{8}$	30.1	15.0	10.0	6.0	1.6	11.6	13.3	22.2	—	3.6
$\frac{15}{16}$ and $1\frac{7}{8}$1	.9	—	—	—	—	1.8	.3	—	—
$1\frac{1}{4}$ and longer.....	—	—	—	—	—	—	—	—	—	—

Staple length (inches)	Ginned Dec. 1-Jan. 15, crop of—					Ginned after Jan. 15, crop of—				
	1932	1931	1930	1929	1928	1932	1931	1930	1929	1928
	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales
All lengths ²	13.2	23.5	70.2	40.3	24.8	4.8	14.4	23.1	25.9	21.4
Shorter than $\frac{7}{8}$ ³	—	.6	—	.1	—	—	8.0	—	.3	—
$\frac{7}{8}$ and $\frac{29}{32}$1	—	.6	.6	—	1.9	.1	1.1	4.1	.0
$\frac{7}{8}$ and $\frac{31}{32}$	1.3	1.1	4.9	2.8	.8	.7	.6	6.9	4.5	3.8
1 and $1\frac{1}{2}$	5.6	6.5	17.1	13.7	5.9	1.3	1.4	9.2	5.5	9.1
$\frac{11}{16}$ and $1\frac{3}{8}$	5.7	12.7	42.4	31.0	14.3	.8	3.4	5.3	11.4	7.1
$\frac{13}{16}$ and $1\frac{5}{8}$5	2.5	5.2	1.1	4.3	.1	.9	.6	.1	.1
$\frac{15}{16}$ and $1\frac{7}{8}$	—	.1	—	—	—	—	—	—	—	—
$1\frac{1}{4}$ and longer.....	—	—	—	—	—	—	—	—	—	—

See footnotes at end of table.

TABLE 7.—*Staple length of American upland cotton: Quantities ginned in the United States during specified periods,¹ by States, crops of 1932, 1931, 1930, 1929, and 1928—Continued*

FLORIDA

Staple length (inches)	Total, crop of—					Ginned prior to Oct. 1				
	1932	1931	1930	1929	1928	1932	1931	1930	1929	1928
	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales
All lengths ²	15.6	43.4	51.1	29.8	20.1	10.9	32.9	40.3	24.9	11.2
Shorter than $\frac{7}{8}$ ³	2.3	2.9	10.1	11.1	2.3	1.7	2.6	9.0	9.7	1.1
$\frac{7}{8}$ and $2\frac{1}{2}$ ₂₂	12.5	33.9	34.9	17.2	14.5	8.6	25.1	27.8	14.2	8.2
$\frac{1}{2}$ ₁₆ and $3\frac{1}{2}$ ₂₂7	6.1	5.8	1.2	2.6	.5	4.7	3.3	.8	1.5
1 and $1\frac{1}{2}$ ₂₂1	.4	.3	.2	.4	.1	.4	.2	.2	.3
$1\frac{1}{2}$ ₁₆ and $1\frac{3}{4}$ ₂₂1	—	—	.1	—	—	.1	—	—	.1
$1\frac{1}{2}$ ₁₆ and $1\frac{5}{8}$ ₂₂	—	—	—	—	.1	—	—	—	—	—
$1\frac{1}{2}$ ₁₆ and $1\frac{7}{8}$ ₂₂	—	—	—	—	—	—	—	—	—	—
$1\frac{1}{2}$ ₁₆ and longer.....	—	—	—	—	—	—	—	—	—	—

Staple length (inches)	Ginned during October					Ginned during November				
	1932	1931	1930	1929	1928	1932	1931	1930	1929	1928
	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales
All lengths ²	3.4	8.6	8.7	4.0	6.5	0.9	1.5	1.5	0.8	1.9
Shorter than $\frac{7}{8}$ ³4	.3	1.1	1.1	1.2	.2	—	—	.3	—
$\frac{7}{8}$ and $2\frac{1}{2}$ ₂₂	2.9	6.9	6.1	2.4	4.4	.6	1.5	.8	.6	1.6
$\frac{1}{2}$ ₁₆ and $3\frac{1}{2}$ ₂₂1	1.4	1.4	.4	.7	.1	—	.7	—	.3
1 and $1\frac{1}{2}$ ₂₂	—	—	—	.1	—	—	—	—	—	—
$1\frac{1}{2}$ ₁₆ and $1\frac{3}{4}$ ₂₂	—	—	—	.1	—	—	—	—	—	—
$1\frac{1}{2}$ ₁₆ and $1\frac{5}{8}$ ₂₂	—	—	—	—	—	—	—	—	—	—
$1\frac{1}{2}$ ₁₆ and $1\frac{7}{8}$ ₂₂	—	—	—	—	—	—	—	—	—	—
$1\frac{1}{2}$ ₁₆ and longer.....	—	—	—	—	—	—	—	—	—	—

Staple length (inches)	Ginned Dec. 1-Jan. 15, crop of—					Ginned after Jan. 15, crop of—				
	1932	1931	1930	1929	1928	1932	1931	1930	1929	1928
	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales
All lengths ²	0.2	0.2	0.5	0.1	0.3	0.2	0.2	0.1	—	0.2
Shorter than $\frac{7}{8}$ ³2	.2	.2	.1	.2	.2	.2	.1	—	.1
$\frac{7}{8}$ and $2\frac{1}{2}$ ₂₂2	.2	.3	—	—	—	—	—	—	—
$\frac{1}{2}$ ₁₆ and $3\frac{1}{2}$ ₂₂	—	—	—	—	—	—	—	—	—	—
1 and $1\frac{1}{2}$ ₂₂	—	—	—	—	—	—	—	—	—	—
$1\frac{1}{2}$ ₁₆ and $1\frac{3}{4}$ ₂₂	—	—	—	—	—	—	—	—	—	—
$1\frac{1}{2}$ ₁₆ and $1\frac{5}{8}$ ₂₂	—	—	—	—	—	—	—	—	—	—
$1\frac{1}{2}$ ₁₆ and $1\frac{7}{8}$ ₂₂	—	—	—	—	—	—	—	—	—	—
$1\frac{1}{2}$ ₁₆ and longer.....	—	—	—	—	—	—	—	—	—	—

GEORGIA

Staple length (inches)	Total, crop of—					Ginned prior to Oct. 1				
	1932	1931	1930	1929	1928	1932	1931	1930	1929	1928
	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales
All lengths ²	881.8	1,393.7	1,507.5	1,339.8	1,053.2	426.2	649.4	842.2	578.2	308.9
Shorter than $\frac{7}{8}$ ³	43.1	69.4	237.1	282.2	120.5	30.1	22.5	131.9	71.2	28.1
$\frac{7}{8}$ and $2\frac{1}{2}$ ₂₂	502.8	1,015.6	1,006.8	918.8	743.9	315.8	478.2	684.4	436.0	226.0
$\frac{1}{2}$ ₁₆ and $3\frac{1}{2}$ ₂₂	163.1	242.1	218.4	113.7	144.1	57.5	112.4	101.5	54.4	43.3
1 and $1\frac{1}{2}$ ₂₂	53.0	54.8	36.3	21.3	28.3	19.5	28.6	18.1	12.8	8.7
$1\frac{1}{2}$ ₁₆ and $1\frac{3}{4}$ ₂₂	6.7	10.4	7.3	3.6	5.4	2.3	6.7	5.1	2.8	2.1
$1\frac{1}{2}$ ₁₆ and $1\frac{5}{8}$ ₂₂	1.9	1.3	1.4	.2	1.4	.0	1.0	1.1	.1	.6
$1\frac{1}{2}$ ₁₆ and $1\frac{7}{8}$ ₂₂3	—	.2	—	.6	.1	—	.1	—	.1
$1\frac{1}{2}$ ₁₆ and longer.....	—	.1	—	—	—	—	—	—	—	—

See footnotes at end of table.

76 STATISTICAL BULLETIN 47, U. S. DEPT. OF AGRICULTURE

TABLE 7.—*Staple length of American upland cotton: Quantities ginned in the United States during specified periods,¹ by States, crops of 1932, 1931, 1930, 1929, and 1928—Continued*

GEORGIA—Continued

Staple length (inches)	Ginned during October					Ginned during November				
	1932	1931	1930	1929	1928	1932	1931	1930	1929	1928
All lengths ² -----	1,000 bales 275.0	1,000 bales 529.0	1,000 bales 531.3	1,000 bales 452.8	1,000 bales 475.5	1,000 bales 114.4	1,000 bales 166.2	1,000 bales 139.2	1,000 bales 146.0	1,000 bales 186.4
Shorter than $\frac{7}{8}$ ³ -----	10.3	29.2	84.2	139.3	79.9	2.5	15.3	16.7	30.8	13.9
$\frac{7}{8}$ and $\frac{21}{32}$ -----	106.4	378.0	356.7	271.4	305.3	81.3	124.6	93.1	100.2	146.4
$\frac{15}{16}$ and $\frac{31}{32}$ -----	70.3	98.8	76.1	36.0	70.8	23.2	22.4	25.1	12.6	22.6
1 and $1\frac{1}{2}$ -----	23.9	19.6	12.6	5.0	16.0	8.5	3.6	3.0	2.0	2.6
$1\frac{1}{16}$ and $1\frac{3}{4}$ -----	3.1	3.1	1.4	.5	2.6	.7	.3	.6	.3	.6
$1\frac{1}{8}$ and $1\frac{1}{2}$ -----	.9	.2	.2	-----	.4	.1	-----	.1	.1	.3
$1\frac{1}{16}$ and $1\frac{1}{2}$ -----	.1	-----	.1	-----	.5	.1	-----	.1	-----	.3
$1\frac{1}{4}$ and longer-----	-----	.1	-----	-----	-----	-----	-----	-----	-----	-----

Staple length (inches)	Ginned Dec. 1-Jan. 15, crop of—					Ginned after Jan. 15, crop of—				
	1932	1931	1930	1929	1928	1932	1931	1930	1929	1928
All lengths ² -----	1,000 bales 37.2	1,000 bales 35.9	1,000 bales 66.9	1,000 bales 129.6	1,000 bales 67.2	1,000 bales 9.0	1,000 bales 13.2	1,000 bales 17.9	1,000 bales 33.2	1,000 bales 15.2
Shorter than $\frac{7}{8}$ ³ -----	.2	2.0	3.1	35.0	5.5	-----	.4	1.2	5.9	2.1
$\frac{7}{8}$ and $\frac{21}{32}$ -----	25.1	25.7	48.3	87.0	53.3	4.2	9.1	14.3	23.3	12.7
$\frac{15}{16}$ and $\frac{31}{32}$ -----	9.5	6.0	13.5	6.9	7.0	2.6	2.5	2.2	3.2	.4
1 and $1\frac{1}{2}$ -----	2.2	1.9	1.8	.7	1.0	1.8	1.1	.2	.8	-----
$1\frac{1}{16}$ and $1\frac{3}{4}$ -----	.2	.2	.2	-----	.1	.4	.1	-----	-----	-----
$1\frac{1}{8}$ and $1\frac{1}{2}$ -----	-----	.1	-----	-----	.1	-----	-----	-----	-----	-----
$1\frac{1}{16}$ and $1\frac{1}{2}$ -----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
$1\frac{1}{4}$ and longer-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----

LOUISIANA

Staple length (inches)	Total, crop of—					Ginned prior to Oct. 1				
	1932	1931	1930	1929	1928	1932	1931	1930	1929	1928
All lengths ² -----	1,000 bales 599.5	1,000 bales 876.6	1,000 bales 704.8	1,000 bales 797.7	1,000 bales 685.0	1,000 bales 303.9	1,000 bales 300.1	1,000 bales 399.6	1,000 bales 542.4	1,000 bales 370.3
Shorter than $\frac{7}{8}$ ³ -----	62.1	58.6	91.3	110.6	88.3	41.1	20.9	53.7	69.2	38.0
$\frac{7}{8}$ and $\frac{21}{32}$ -----	131.0	157.3	160.1	205.5	203.1	84.4	39.9	76.9	141.0	108.5
$\frac{15}{16}$ and $\frac{31}{32}$ -----	152.6	203.6	167.4	212.7	178.6	93.6	54.4	91.7	143.6	97.5
1 and $1\frac{1}{2}$ -----	156.4	283.0	188.8	170.4	127.6	107.4	99.8	111.7	117.0	73.1
$1\frac{1}{16}$ and $1\frac{3}{4}$ -----	68.7	133.1	67.2	63.7	52.1	45.6	61.6	40.4	45.5	31.1
$1\frac{1}{8}$ and $1\frac{1}{2}$ -----	26.2	35.4	28.0	29.6	27.7	19.8	19.2	21.8	21.7	17.6
$1\frac{1}{16}$ and $1\frac{3}{4}$ -----	2.4	5.3	3.8	5.0	7.8	1.9	4.1	3.2	4.3	6.0
$1\frac{1}{4}$ and longer-----	.1	.3	.2	.2	.7	.1	.2	.2	.1	.5

Staple length (inches)	Ginned during October					Ginned during November				
	1932	1931	1930	1929	1928	1932	1931	1930	1929	1928
All lengths ² -----	1,000 bales 156.0	1,000 bales 367.5	1,000 bales 207.8	1,000 bales 202.8	1,000 bales 215.9	1,000 bales 40.2	1,000 bales 146.1	1,000 bales 74.7	1,000 bales 34.3	1,000 bales 77.3
Shorter than $\frac{7}{8}$ ³ -----	15.8	29.5	28.2	34.3	20.4	4.0	5.8	8.2	6.9	14.9
$\frac{7}{8}$ and $\frac{21}{32}$ -----	37.0	82.0	58.8	54.4	67.6	7.5	27.0	10.2	8.3	23.9
$\frac{15}{16}$ and $\frac{31}{32}$ -----	47.3	88.8	47.9	54.5	50.1	9.2	39.8	21.6	7.9	18.3
1 and $1\frac{1}{2}$ -----	33.0	109.1	51.8	30.4	38.0	12.9	48.4	18.1	7.4	13.4
$1\frac{1}{16}$ and $1\frac{3}{4}$ -----	16.7	47.3	17.6	13.8	15.1	5.9	18.8	5.9	2.5	4.7
$1\frac{1}{8}$ and $1\frac{1}{2}$ -----	5.8	10.0	3.3	5.8	7.7	.6	5.8	1.6	1.2	2.1
$1\frac{1}{16}$ and $1\frac{3}{4}$ -----	.4	.7	.2	.5	1.8	.1	.5	.1	.1	.1
$1\frac{1}{4}$ and longer-----	-----	.1	-----	.1	.2	-----	-----	-----	-----	-----

See footnotes at end of table

TABLE 7.—*Staple length of American upland cotton: Quantities ginned in the United States during specified periods,¹ by States, crops of 1932, 1931, 1930, 1929, and 1928—Continued*

LOUISIANA—Continued

Staple length (inches)	Ginned Dec. 1-Jan. 15, crop of—					Ginned after Jan. 15, crop of—					Ginned after Jan. 15, crop of—					
	1932		1931		1930	1929		1928	1932		1931		1930	1929		1928
	<i>1,000 bales</i>		<i>1,000 bales</i>		<i>1,000 bales</i>	<i>1,000 bales</i>		<i>1,000 bales</i>	<i>1,000 bales</i>		<i>1,000 bales</i>		<i>1,000 bales</i>	<i>1,000 bales</i>		<i>1,000 bales</i>
All lengths ²	7.7	37.0	19.5	13.9	10.0	1.7	25.9	3.2	4.3	3.4						
Shorter than $\frac{7}{8}$ ³8	1.5	.9	.1	5.4	.4	.9	.3	.1	.6						
$\frac{7}{8}$ and $\frac{2\frac{1}{2}}{8}$	1.5	5.6	4.0	1.4	4.7	.6	2.8	1.2	.4	.4						
$\frac{7}{8}$ and $\frac{3\frac{1}{2}}{8}$	2.2	14.0	5.8	4.6	5.0	.3	6.6	.4	2.1	1.7						
1 and $1\frac{1}{2}$	2.8	13.3	5.0	5.0	2.5	.3	12.4	.2	1.6	.6						
$\frac{1\frac{1}{2}}{8}$ and $\frac{1\frac{3}{4}}{8}$4	2.5	2.4	1.8	1.1	.1	2.9	.9	.1	.1						
$\frac{1\frac{1}{2}}{8}$ and $\frac{1\frac{1}{2}}{8}$1	1.1	.9	.3	—	.3	—	.2	—	—						
$\frac{1\frac{1}{2}}{8}$ and $\frac{1\frac{1}{2}}{8}$3	.1	—	—	—	—	—	—	—	—						
$1\frac{1}{4}$ and longer.....	—	—	—	—	—	—	—	—	—	—						

MISSISSIPPI

Staple length (inches)	Total, crop of—					Ginned prior to Oct. 1										
	1932		1931		1930	1929		1928	1932		1931		1930	1929		1928
	<i>1,000 bales</i>		<i>1,000 bales</i>		<i>1,000 bales</i>	<i>1,000 bales</i>		<i>1,000 bales</i>	<i>1,000 bales</i>		<i>1,000 bales</i>		<i>1,000 bales</i>	<i>1,000 bales</i>		<i>1,000 bales</i>
All lengths ²	1,161.2	1,719.4	1,458.5	1,876.0	1,462.0	511.7	358.5	532.1	908.4	560.3						
Shorter than $\frac{7}{8}$ ³	32.2	65.4	114.9	189.7	254.3	20.2	16.3	39.6	92.3	81.9						
$\frac{7}{8}$ and $\frac{2\frac{1}{2}}{8}$	148.6	285.4	290.2	352.3	258.9	71.5	76.3	132.6	191.7	114.8						
$\frac{7}{8}$ and $\frac{3\frac{1}{2}}{8}$	162.3	238.0	264.5	227.0	111.9	63.3	49.2	95.2	103.6	47.8						
1 and $1\frac{1}{2}$	170.9	234.8	209.4	202.7	127.8	70.2	38.1	68.6	78.7	44.4						
$\frac{1\frac{1}{2}}{8}$ and $\frac{1\frac{3}{4}}{8}$	271.5	311.0	312.9	388.9	259.8	146.3	20.7	103.5	136.1	68.2						
$\frac{1\frac{1}{2}}{8}$ and $\frac{1\frac{1}{2}}{8}$	328.2	368.1	216.6	412.4	305.9	128.6	57.7	77.6	228.6	105.5						
$\frac{1\frac{1}{2}}{8}$ and $\frac{1\frac{1}{2}}{8}$	45.3	188.7	30.4	97.7	121.7	11.1	71.7	14.7	73.2	79.4						
$1\frac{1}{4}$ and longer.....	2.2	28.0	1.6	5.3	21.7	.5	19.5	.3	4.2	18.3						

Staple length (inches)	Ginned during October					Ginned during November										
	1932		1931		1930	1929		1928	1932		1931		1930	1929		1928
	<i>1,000 bales</i>		<i>1,000 bales</i>		<i>1,000 bales</i>	<i>1,000 bales</i>		<i>1,000 bales</i>	<i>1,000 bales</i>		<i>1,000 bales</i>		<i>1,000 bales</i>	<i>1,000 bales</i>		<i>1,000 bales</i>
All lengths ²	385.2	781.2	594.0	632.8	530.7	198.3	378.2	251.0	151.3	247.0						
Shorter than $\frac{7}{8}$ ³	7.4	41.1	66.7	83.8	132.7	4.1	7.2	5.7	9.7	28.8						
$\frac{7}{8}$ and $\frac{2\frac{1}{2}}{8}$	40.3	149.0	120.8	118.4	81.4	26.6	47.3	34.7	23.2	48.4						
$\frac{7}{8}$ and $\frac{3\frac{1}{2}}{8}$	57.7	107.7	101.0	78.1	31.2	31.1	56.8	52.8	22.6	23.2						
1 and $1\frac{1}{2}$	56.9	94.2	79.1	79.9	48.8	29.5	55.1	43.2	21.5	21.8						
$\frac{1\frac{1}{2}}{8}$ and $\frac{1\frac{3}{4}}{8}$	66.8	121.1	113.6	137.8	98.1	41.1	77.1	70.6	43.9	50.6						
$\frac{1\frac{1}{2}}{8}$ and $\frac{1\frac{1}{2}}{8}$	128.8	178.9	93.3	116.7	114.9	58.3	101.9	38.4	26.6	63.2						
$\frac{1\frac{1}{2}}{8}$ and $\frac{1\frac{1}{2}}{8}$	20.0	82.1	18.5	17.3	29.7	7.3	31.6	5.3	3.6	10.6						
$1\frac{1}{4}$ and longer.....	1.3	7.1	1.0	.8	2.9	.3	1.2	.3	.2	.4						

Staple length (inches)	Ginned Dec. 1-Jan. 15, crop of—					Ginned after Jan. 15, crop of—										
	1932		1931		1930	1929		1928	1932		1931		1930	1929		1928
	<i>1,000 bales</i>		<i>1,000 bales</i>		<i>1,000 bales</i>	<i>1,000 bales</i>		<i>1,000 bales</i>	<i>1,000 bales</i>		<i>1,000 bales</i>		<i>1,000 bales</i>	<i>1,000 bales</i>		<i>1,000 bales</i>
All lengths ²	53.6	89.0	72.4	121.2	81.3	12.4	112.5	9.0	62.3	33.7						
Shorter than $\frac{7}{8}$ ³5	.8	2.9	2.8	8.7	—	—	—	—	—	1.1	2.2				
$\frac{7}{8}$ and $\frac{2\frac{1}{2}}{8}$	7.8	10.2	8.2	15.0	12.4	2.4	2.6	2.9	4.0	1.9						
$\frac{7}{8}$ and $\frac{3\frac{1}{2}}{8}$	7.4	16.8	13.9	16.2	6.9	2.8	7.5	1.6	6.5	2.8						
1 and $1\frac{1}{2}$	10.1	16.5	17.2	13.7	7.1	4.2	30.9	1.3	8.9	5.7						
$\frac{1\frac{1}{2}}{8}$ and $\frac{1\frac{3}{4}}{8}$	14.7	26.9	23.1	41.5	28.0	2.6	56.2	2.1	29.6	14.9						
$\frac{1\frac{1}{2}}{8}$ and $\frac{1\frac{1}{2}}{8}$	12.1	15.5	6.3	28.8	16.4	.4	14.1	1.0	11.7	5.9						
$\frac{1\frac{1}{2}}{8}$ and $\frac{1\frac{1}{2}}{8}$9	2.2	.8	3.1	1.7	—	1.1	.1	.5	.3						
$1\frac{1}{4}$ and longer.....	.1	.1	—	.1	.1	—	—	—	—	—						

See footnotes at end of table.

78 STATISTICAL BULLETIN 47, U. S. DEPT. OF AGRICULTURE

TABLE 7.—*Staple length of American upland cotton: Quantities ginned in the United States during specified periods,¹ by States, crops of 1932, 1931, 1930, 1929, and 1928—Continued*

MISSOURI

Staple length (inches)	Total, crop of—					Ginned prior to Oct. 1										
	1932		1931		1930	1929		1928	1932		1931		1930	1929		
	<i>1,000 bales</i>	<i>1,000 bales</i>	<i>1,000 bales</i>	<i>1,000 bales</i>	<i>1,000 bales</i>	<i>1,000 bales</i>	<i>1,000 bales</i>	<i>1,000 bales</i>	<i>1,000 bales</i>	<i>1,000 bales</i>						
All lengths ²	300.7	280.4	153.3	220.9	146.9	80.7	32.5	52.9	23.1	13.6						
Shorter than $\frac{7}{8}$ ³	2.0	2.6	11.2	3.1	8.8	.5					4.6	2.0	.4			
$\frac{7}{8}$ and $\frac{29}{32}$	52.1	85.9	56.0	57.5	30.2	17.1	9.0	24.4	11.4	5.3						
$\frac{15}{16}$ and $\frac{31}{32}$	114.9	128.0	55.6	88.4	52.2	29.0	11.7	16.4	7.5	4.8						
1 and $\frac{11}{32}$	107.7	53.9	22.4	56.2	39.3	27.9	8.9	6.2	2.0	2.8						
$\frac{13}{16}$ and $\frac{13}{32}$	23.0	9.5	7.1	13.7	14.3	6.0	2.7	1.3	.2	.5						
$\frac{13}{16}$ and $\frac{15}{32}$	1.0	.5	1.0	1.8	1.7	.2										
$\frac{13}{16}$ and $\frac{17}{32}$.2	.4											
$\frac{1}{4}$ and longer																

Staple length (inches)	Ginned during October					Ginned during November										
	1932		1931		1930	1929		1928	1932		1931		1930	1929		
	<i>1,000 bales</i>	<i>1,000 bales</i>	<i>1,000 bales</i>	<i>1,000 bales</i>	<i>1,000 bales</i>	<i>1,000 bales</i>	<i>1,000 bales</i>	<i>1,000 bales</i>	<i>1,000 bales</i>	<i>1,000 bales</i>	<i>1,000 bales</i>	<i>1,000 bales</i>	<i>1,000 bales</i>	<i>1,000 bales</i>	<i>1,000 bales</i>	
All lengths ²	118.3	102.7	64.4	79.2	35.6	55.6	65.9	28.8	58.3	45.4						
Shorter than $\frac{7}{8}$ ³	1.2	1.0	4.3	.3	.2	.2	.8	.5	.1	.2						
$\frac{7}{8}$ and $\frac{29}{32}$	10.8	23.7	26.3	17.5	7.3	7.5	17.7	4.2	10.9	5.0						
$\frac{15}{16}$ and $\frac{31}{32}$	41.7	42.1	23.7	28.0	12.7	21.7	33.2	12.7	27.2	13.0						
1 and $\frac{11}{32}$	50.8	30.1	7.0	26.5	11.9	22.8	12.9	7.7	14.8	18.0						
$\frac{13}{16}$ and $\frac{13}{32}$	13.0	5.5	2.5	6.1	3.0	3.4	1.3	3.3	4.7	9.6						
$\frac{13}{16}$ and $\frac{15}{32}$.8	.3	.6	.7	.4				.4	.7						
$\frac{13}{16}$ and $\frac{17}{32}$.1	.1					.1						
$\frac{1}{4}$ and longer																

Staple length (inches)	Ginned Dec. 1-Jan. 15, crop of—					Ginned after Jan. 15, crop of—										
	1932		1931		1930	1929		1928	1932		1931		1930	1929		
	<i>1,000 bales</i>	<i>1,000 bales</i>	<i>1,000 bales</i>	<i>1,000 bales</i>	<i>1,000 bales</i>	<i>1,000 bales</i>	<i>1,000 bales</i>	<i>1,000 bales</i>	<i>1,000 bales</i>	<i>1,000 bales</i>	<i>1,000 bales</i>	<i>1,000 bales</i>	<i>1,000 bales</i>	<i>1,000 bales</i>	<i>1,000 bales</i>	
All lengths ²	35.2	48.4	5.9	37.7	40.4	10.9	30.9	1.3	22.6	11.0						
Shorter than $\frac{7}{8}$ ³	.1	.7	1.5	.2	5.5						.1	.3	.5	.2	.5	2.5
$\frac{7}{8}$ and $\frac{29}{32}$	12.8	21.6	.9	8.4	9.7	3.9	13.9	.2	9.3	2.9						
$\frac{15}{16}$ and $\frac{31}{32}$	16.3	24.9	2.3	16.2	18.1	6.2	16.1	.5	9.5	3.8						
1 and $\frac{11}{32}$	5.4	1.2	1.2	10.3	6.3	.8	.8	.3	2.8	2.3						
$\frac{13}{16}$ and $\frac{13}{32}$.6			2.3	.8						.4					
$\frac{13}{16}$ and $\frac{15}{32}$.3							.1					
$\frac{13}{16}$ and $\frac{17}{32}$																
$\frac{1}{4}$ and longer																

NEW MEXICO

Staple length (inches)	Total, crop of—					Ginned prior to Oct. 1										
	1932		1931		1930	1929		1928	1932		1931		1930	1929		
	<i>1,000 bales</i>	<i>1,000 bales</i>	<i>1,000 bales</i>	<i>1,000 bales</i>	<i>1,000 bales</i>	<i>1,000 bales</i>	<i>1,000 bales</i>	<i>1,000 bales</i>	<i>1,000 bales</i>	<i>1,000 bales</i>						
All lengths ²	67.5	93.8	95.8	86.3	82.2	4.6	5.6	17.0	8.4	6.1						
Shorter than $\frac{7}{8}$ ³	2.1	5.1	.5	4.8	4.9											
$\frac{7}{8}$ and $\frac{29}{32}$	2.6	1.4	1.3	2.7	6.2											
$\frac{15}{16}$ and $\frac{31}{32}$	3.4	13.7	7.8	6.3	8.6											
1 and $\frac{11}{32}$	23.6	36.7	35.6	40.2	22.1	.4	1.3	6.6	4.3	1.7						
$\frac{13}{16}$ and $\frac{13}{32}$	25.1	31.4	42.5	27.7	34.0	2.1	3.5	9.6	3.8	2.8						
$\frac{13}{16}$ and $\frac{15}{32}$	10.7	5.5	8.0	4.6	6.4	2.1	.8	.7	.2	.1						
$\frac{13}{16}$ and $\frac{17}{32}$.1													
$\frac{1}{4}$ and longer																

See footnotes at end of table.

TABLE 7.—*Staple length of American upland cotton: Quantities ginned in the United States during specified periods,¹ by States, crops of 1932, 1931, 1930, 1929, and 1928—Continued*

NEW MEXICO—Continued

Staple length (inches)	Ginned during October					Ginned during November				
	1932	1931	1930	1929	1928	1932	1931	1930	1929	1928
	1,000 bales 23.1	1,000 bales 34.0	1,000 bales 32.2	1,000 bales 28.6	1,000 bales 27.1	1,000 bales 25.7	1,000 bales 29.2	1,000 bales 28.6	1,000 bales 27.6	1,000 bales 24.1
All lengths ²										
Shorter than $\frac{7}{8}$ ³3					
$\frac{7}{8}$ and $\frac{29}{32}$					4.0					
$\frac{15}{16}$ and $\frac{31}{32}$		2.1	.2	.4	5.4	.3	5.5	1.9	1.1	1.2
1 and $\frac{1}{2}$	3.5	15.5	12.7	5.6	8.5	14.2	11.8	9.4	21.9	7.1
$\frac{17}{16}$ and $\frac{19}{32}$	12.2	13.9	16.8	18.2	7.6	10.0	10.0	12.5	4.6	13.0
$\frac{19}{16}$ and $\frac{19}{32}$	7.4	2.5	2.5	4.4	1.3	1.2	1.9	4.6		2.7
$\frac{13}{16}$ and $\frac{17}{32}$1	
$\frac{1}{4}$ and longer.....										

Staple length (inches)	Ginned Dec. 1-Jan. 15, crop of—					Ginned after Jan. 15, crop of—				
	1932	1931	1930	1929	1928	1932	1931	1930	1929	1928
	1,000 bales 10.7	1,000 bales 17.6	1,000 bales 16.2	1,000 bales 19.7	1,000 bales 19.7	1,000 bales 3.4	1,000 bales 7.4	1,000 bales 1.8	1,000 bales 2.0	1,000 bales 5.2
All lengths ²										
Shorter than $\frac{7}{8}$ ³	1.9	.8	.4	4.0	2.2	.2	4.3	.1	.8	2.4
$\frac{7}{8}$ and $\frac{29}{32}$5	1.2	1.0	2.0	.6	2.1	.2	.2	.7	1.4
$\frac{15}{16}$ and $\frac{31}{32}$	2.7	4.3	4.2	4.3	.9	.4	1.8	1.4	.4	.7
1 and $\frac{1}{2}$	4.8	7.3	6.8	8.3	4.3	.7	.8	.1	.1	.5
$\frac{17}{16}$ and $\frac{19}{32}$8	3.8	3.6	1.1	10.4		.2			
$\frac{19}{16}$ and $\frac{19}{32}$2			1.3		.1			
$\frac{13}{16}$ and $\frac{17}{32}$										
$\frac{1}{4}$ and longer.....										

NORTH CAROLINA

Staple length (inches)	Total, crop of—					Ginned prior to Oct. 1				
	1932	1931	1930	1929	1928	1932	1931	1930	1929	1928
	1,000 bales 680.3	1,000 bales 771.2	1,000 bales 800.6	1,000 bales 767.0	1,000 bales 869.3	1,000 bales 226.1	1,000 bales 215.6	1,000 bales 198.5	1,000 bales 50.2	1,000 bales 60.1
All lengths ²										
Shorter than $\frac{7}{8}$ ³	7.8	7.8	40.4	81.9	87.1	5.5	1.9	12.4	2.8	10.8
$\frac{7}{8}$ and $\frac{29}{32}$	227.8	340.1	410.2	479.4	605.5	72.4	92.8	112.3	31.5	38.4
$\frac{15}{16}$ and $\frac{31}{32}$	223.6	264.6	236.5	151.2	129.7	67.4	69.3	51.2	11.2	7.4
1 and $\frac{1}{2}$	160.4	126.0	81.6	45.7	30.5	60.3	40.2	16.0	3.5	2.6
$\frac{17}{16}$ and $\frac{19}{32}$	47.5	28.5	25.6	7.0	9.0	16.7	10.1	4.9	.9	.5
$\frac{19}{16}$ and $\frac{19}{32}$	11.0	3.8	5.6	1.5	4.9	3.4	1.2	1.4	.2	.3
$\frac{13}{16}$ and $\frac{17}{32}$	2.1	.4	.7	.3	2.2	.4	.1	.3	.1	.1
$\frac{1}{4}$ and longer.....	.1				.4					

Staple length (inches)	Ginned during October					Ginned during November				
	1932	1931	1930	1929	1928	1932	1931	1930	1929	1928
	1,000 bales 264.1	1,000 bales 382.2	1,000 bales 391.0	1,000 bales 340.4	1,000 bales 437.0	1,000 bales 125.2	1,000 bales 134.7	1,000 bales 133.8	1,000 bales 207.7	1,000 bales 243.2
All lengths ²										
Shorter than $\frac{7}{8}$ ³	1.5	3.9	24.5	49.6	50.3	.6	2.0	2.7	18.1	12.0
$\frac{7}{8}$ and $\frac{29}{32}$	71.8	153.2	199.9	211.2	276.0	55.6	76.8	58.9	122.8	185.5
$\frac{15}{16}$ and $\frac{31}{32}$	92.3	138.3	107.0	50.0	76.6	41.4	42.1	50.8	48.4	34.1
1 and $\frac{1}{2}$	69.6	69.1	40.9	20.2	18.3	19.7	11.2	16.8	15.5	7.4
$\frac{17}{16}$ and $\frac{19}{32}$	22.1	15.2	15.2	2.7	5.4	5.7	2.3	3.9	2.5	2.3
$\frac{19}{16}$ and $\frac{19}{32}$	5.5	2.2	3.1	.6	2.9	1.7	.3	.7	.4	1.2
$\frac{13}{16}$ and $\frac{17}{32}$	1.2	.3	.4	.1	1.3	.5				.6
$\frac{1}{4}$ and longer.....	.1				.2					.1

See footnotes at end of table.

TABLE 7.—*Staple length of American upland cotton: Quantities ginned in the United States during specified periods,¹ by States, crops of 1932, 1931, 1930, 1929, and 1928—Continued*

NORTH CAROLINA—Continued

Staple length (inches)	Ginned Dec. 1-Jan. 15, crop of—					Ginned after Jan. 15, crop of—										
	1932		1931		1930	1929		1928	1932		1931		1930	1929		
	<i>1,000 bales</i>		<i>1,000 bales</i>		<i>1,000 bales</i>	<i>1,000 bales</i>		<i>1,000 bales</i>	<i>1,000 bales</i>		<i>1,000 bales</i>		<i>1,000 bales</i>	<i>1,000 bales</i>		
All lengths ²	51.9		31.2		50.7	139.4		104.9	13.0		7.5		20.6	20.3		24.1
Shorter than $\frac{7}{8}$ ³	.2				.4	8.3		5.7					.4	3.1		2.3
$\frac{7}{8}$ and $2\frac{1}{2}$ ₂	21.9		14.5		28.4	97.3		85.2	6.1		2.8		10.7	16.6		20.4
$1\frac{1}{16}$ and $3\frac{1}{2}$ ₂	18.2		11.6		20.3	28.5		10.6	4.3		3.3		7.2	7.1		1.0
1 and $1\frac{1}{2}$ ₂	8.8		4.3		5.0	4.4		1.9			2.0		1.2	2.0		.3
$1\frac{1}{16}$ and $1\frac{3}{4}$ ₂	2.5		.7		1.3	.6		.7	.5		.2		.3	.3		.1
$1\frac{1}{8}$ and $1\frac{3}{4}$ ₂	.3		.1		.4	.2		.5			.1					.1
$1\frac{1}{16}$ and $1\frac{1}{2}$ ₂						.1										
1 $\frac{1}{4}$ and longer																

OKLAHOMA

Staple length (inches)	Total, crop of—					Ginned prior to Oct. 1										
	1932		1931		1930	1929		1928	1932		1931		1930	1929		
	<i>1,000 bales</i>		<i>1,000 bales</i>		<i>1,000 bales</i>	<i>1,000 bales</i>		<i>1,000 bales</i>	<i>1,000 bales</i>		<i>1,000 bales</i>		<i>1,000 bales</i>	<i>1,000 bales</i>		
All lengths ²	1,072.0		1,235.9		856.8	1,125.6		1,187.0	290.2		281.1		276.6	255.1		288.1
Shorter than $\frac{7}{8}$ ³	91.9		137.9		126.7	321.5		162.3	34.4		6.6		49.3	63.6		34.6
$\frac{7}{8}$ and $2\frac{1}{2}$ ₂	535.8		557.4		348.0	500.1		421.7	137.5		74.9		105.3	83.0		72.1
$1\frac{1}{16}$ and $3\frac{1}{2}$ ₂	386.1		414.5		295.1	207.9		395.5	98.2		136.2		85.5	69.5		87.8
1 and $1\frac{1}{2}$ ₂	52.4		110.8		71.6	67.1		172.3	19.7		55.5		28.1	26.8		50.1
$1\frac{1}{16}$ and $1\frac{3}{4}$ ₂	5.2		14.0		12.2	22.6		25.3	2.3		7.3		6.6	9.6		11.8
$1\frac{1}{8}$ and $1\frac{3}{4}$ ₂	.5		1.3		2.5	6.1		8.2	.1		.6		1.8	3.5		4.8
$1\frac{1}{16}$ and $1\frac{1}{2}$ ₂	.1				.1	.3		1.6								.1
1 $\frac{1}{4}$ and longer									.1							

Staple length (inches)	Ginned during October					Ginned during November										
	1932		1931		1930	1929		1928	1932		1931		1930	1929		
	<i>1,000 bales</i>		<i>1,000 bales</i>		<i>1,000 bales</i>	<i>1,000 bales</i>		<i>1,000 bales</i>	<i>1,000 bales</i>		<i>1,000 bales</i>		<i>1,000 bales</i>	<i>1,000 bales</i>		
All lengths ²	464.5		465.0		313.0	450.6		484.8	241.5		307.5		208.9	254.4		198.3
Shorter than $\frac{7}{8}$ ³	21.5		47.1		28.2	141.3		53.2	8.1		38.9		33.0	73.3		8.5
$\frac{7}{8}$ and $2\frac{1}{2}$ ₂	235.8		215.6		116.8	188.2		150.0	124.2		158.0		97.9	147.5		83.0
$1\frac{1}{16}$ and $3\frac{1}{2}$ ₂	184.4		157.6		126.2	77.9		185.4	96.2		93.1		71.0	28.1		74.9
1 and $1\frac{1}{2}$ ₂	20.4		39.0		35.9	29.4		82.7	12.0		15.0		6.6	4.4		25.3
$1\frac{1}{16}$ and $1\frac{3}{4}$ ₂	2.0		5.2		5.2	11.1		10.0	.9		1.5		.3	1.0		3.2
$1\frac{1}{8}$ and $1\frac{3}{4}$ ₂	.4		.5		.6	2.5		2.9			.1		.1	.1		.4
$1\frac{1}{16}$ and $1\frac{1}{2}$ ₂					.1	.2		.5			.1					.1
1 $\frac{1}{4}$ and longer									.1							

Staple length (inches)	Ginned Dec. 1-Jan. 15, crop of—					Ginned after Jan. 15, crop of—										
	1932		1931		1930	1929		1928	1932		1931		1930	1929		
	<i>1,000 bales</i>		<i>1,000 bales</i>		<i>1,000 bales</i>	<i>1,000 bales</i>		<i>1,000 bales</i>	<i>1,000 bales</i>		<i>1,000 bales</i>		<i>1,000 bales</i>	<i>1,000 bales</i>		
All lengths ²	55.6		145.3		53.1	144.7		202.2	20.2		37.0		5.2	20.8		36.6
Shorter than $\frac{7}{8}$ ³	11.9		29.6		13.5	37.2		40.7	16.0		15.7		2.7	6.1		26.4
$\frac{7}{8}$ and $2\frac{1}{2}$ ₂	34.4		88.9		26.8	71.7		108.6	3.9		19.1		1.8	9.7		7.1
$1\frac{1}{16}$ and $3\frac{1}{2}$ ₂	9.0		25.4		11.8	28.2		44.3	.3		2.2		.6	4.2		3.1
1 and $1\frac{1}{2}$ ₂	.3		1.3		.9	6.7		8.2					.1	.8		
$1\frac{1}{16}$ and $1\frac{3}{4}$ ₂					.1	.9		.3								
$1\frac{1}{8}$ and $1\frac{3}{4}$ ₂					.1			.1								
$1\frac{1}{16}$ and $1\frac{1}{2}$ ₂																
1 $\frac{1}{4}$ and longer																

See footnotes at end of table.

TABLE 7.—*Staple length of American upland cotton: Quantities ginned in the United States during specified periods,¹ by States, crops of 1932, 1931, 1930, 1929, and 1928—Continued*

SOUTH CAROLINA

Staple length (inches)	Total, crop of—					Ginned prior to Oct. 1				
	1932	1931	1930	1929	1928	1932	1931	1930	1929	1928
All lengths ²	1,000 bales 722.2	1,000 bales 1,010.2	1,000 bales 1,015.3	1,000 bales 833.1	1,000 bales 744.4	1,000 bales 295.0	1,000 bales 370.6	1,000 bales 377.4	1,000 bales 162.6	1,000 bales 120.7
Shorter than $\frac{7}{8}$ ³	2.8	12.0	20.9	93.1	51.8	1.7	4.1	10.8	9.5	6.3
$\frac{7}{8}$ and $2\frac{1}{2}$	181.2	395.6	426.0	440.4	409.9	81.8	143.3	155.2	81.4	67.9
$\frac{1}{16}$ and $3\frac{1}{2}$	225.6	315.2	288.6	157.4	144.9	84.6	106.2	101.5	36.4	23.9
1 and $1\frac{1}{2}$	160.5	169.8	140.1	81.0	68.5	69.9	61.4	53.8	19.3	10.9
$1\frac{1}{2}$ and $1\frac{3}{4}$	81.8	67.8	83.9	42.6	33.9	34.5	28.0	31.7	10.3	6.0
$1\frac{1}{2}$ and $1\frac{1}{2}$	44.0	36.0	38.4	14.4	23.4	16.9	20.3	17.9	4.3	3.3
$1\frac{1}{2}$ and $1\frac{7}{8}$	16.8	12.0	10.6	3.7	9.6	4.8	6.7	6.1	1.3	1.7
$1\frac{1}{4}$ and longer	1.5	1.8	.8	.5	2.4	.8	.6	.4	.1	.7

Staple length (inches)	Ginned during October					Ginned during November				
	1932	1931	1930	1929	1928	1932	1931	1930	1929	1928
All lengths ²	1,000 bales 258.8	1,000 bales 459.4	1,000 bales 430.7	1,000 bales 371.2	1,000 bales 392.7	1,000 bales 112.1	1,000 bales 144.7	1,000 bales 128.2	1,000 bales 168.8	1,000 bales 168.5
Shorter than $\frac{7}{8}$ ³	.9	6.5	15.1	53.8	35.4	.1	1.3	.8	16.0	6.7
$\frac{7}{8}$ and $2\frac{1}{2}$	48.3	172.5	198.9	193.2	184.9	33.6	65.6	46.2	86.0	112.3
$\frac{1}{16}$ and $3\frac{1}{2}$	80.2	146.2	117.2	66.1	87.3	39.7	48.9	44.2	32.3	26.2
1 and $1\frac{1}{2}$	68.6	84.3	53.5	34.3	43.5	21.4	17.2	21.1	19.5	10.5
$1\frac{1}{2}$ and $1\frac{3}{4}$	32.3	30.9	34.7	16.8	18.8	10.8	8.5	11.4	10.3	7.0
$1\frac{1}{2}$ and $1\frac{1}{2}$	20.1	13.0	13.5	5.7	15.2	4.9	2.7	4.0	3.4	4.0
$1\frac{1}{2}$ and $1\frac{7}{8}$	8.0	4.8	3.4	1.2	6.2	1.5	.5	.5	1.0	1.5
$1\frac{1}{4}$ and longer	.4	1.2	.4	.1	1.4	.1	—	—	.3	.3

Staple length (inches)	Ginned Dec. 1-Jan. 15, crop of—					Ginned after Jan. 15, crop of—				
	1932	1931	1930	1929	1928	1932	1931	1930	1929	1928
All lengths ²	1,000 bales 42.0	1,000 bales 25.1	1,000 bales 60.0	1,000 bales 100.7	1,000 bales 49.7	1,000 bales 14.3	1,000 bales 10.4	1,000 bales 17.0	1,000 bales 23.8	1,000 bales 12.8
Shorter than $\frac{7}{8}$ ³	.1	.1	.2	12.4	2.9	—	—	—	1.4	.5
$\frac{7}{8}$ and $2\frac{1}{2}$	12.4	10.9	19.0	66.3	34.9	5.1	3.3	6.7	13.5	9.9
$\frac{1}{16}$ and $3\frac{1}{2}$	15.8	9.9	18.7	17.2	6.2	5.3	4.0	7.0	5.4	1.3
1 and $1\frac{1}{2}$	7.3	3.8	9.5	6.0	3.0	2.3	3.1	2.2	1.9	.6
$1\frac{1}{2}$ and $1\frac{3}{4}$	3.2	.4	5.4	3.7	1.7	1.0	—	.7	1.5	.4
$1\frac{1}{2}$ and $1\frac{1}{2}$	1.5	—	2.7	.9	.8	.6	—	.3	.1	.1
$1\frac{1}{2}$ and $1\frac{7}{8}$	1.5	—	.5	.2	.2	—	—	.1	—	—
$1\frac{1}{4}$ and longer	.2	—	—	—	—	—	—	—	—	—

TENNESSEE

Staple length (inches)	Total, crop of—					Ginned prior to Oct. 1				
	1932	1931	1930	1929	1928	1932	1931	1930	1929	1928
All lengths ²	1,000 bales 467.5	1,000 bales 578.0	1,000 bales 371.4	1,000 bales 504.3	1,000 bales 423.5	1,000 bales 61.3	1,000 bales 48.4	1,000 bales 87.4	1,000 bales 71.5	1,000 bales 64.7
Shorter than $\frac{7}{8}$ ³	12.4	39.9	59.7	80.0	88.0	2.2	2.3	20.3	15.3	26.9
$\frac{7}{8}$ and $2\frac{1}{2}$	115.0	218.6	128.1	173.4	173.0	19.8	12.8	37.4	24.9	24.0
$\frac{1}{16}$ and $3\frac{1}{2}$	175.3	211.3	123.8	163.8	108.0	21.3	17.5	20.9	20.9	9.6
1 and $1\frac{1}{2}$	134.7	93.5	48.0	74.9	42.1	15.6	13.4	7.6	8.8	3.3
$1\frac{1}{2}$ and $1\frac{3}{4}$	24.0	11.7	11.0	10.3	10.4	2.2	2.0	1.1	1.4	.8
$1\frac{1}{2}$ and $1\frac{1}{2}$	4.3	2.5	.8	1.6	1.4	.2	.3	.1	.2	.1
$1\frac{1}{2}$ and $1\frac{7}{8}$.3	.5	.3	.5	.5	.1	—	—	—	—
$1\frac{1}{4}$ and longer	—	—	—	—	—	—	—	—	—	—

See footnotes at end of table.

TABLE 7.—Staple length of American upland cotton: Quantities ginned in the United States during specified periods,¹ by States, crops of 1932, 1931, 1930, 1929, and 1928—Continued

TENNESSEE—Continued

Staple length (inches)	Ginned during October					Ginned during November										
	1932		1931		1930	1929		1928	1932		1931		1930	1929		1928
	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	
All lengths ²	201.1	288.9	169.0	226.0	154.0	123.0	167.6	85.2	98.5	110.8						
Shorter than $\frac{7}{8}$ ³	5.0	24.3	33.4	42.1	46.0	3.9	11.2	3.7	13.0	7.0						
$\frac{7}{8}$ and $\frac{29}{32}$	35.8	102.2	58.0	72.7	62.5	32.2	62.6	23.9	38.0	40.1						
$\frac{1}{4}$ and $\frac{31}{32}$	70.6	99.0	53.5	67.2	30.1	45.5	68.3	30.8	32.6	37.2						
1 and $1\frac{1}{2}$	73.9	54.9	19.1	38.2	11.9	34.2	21.9	15.7	12.3	19.6						
$\frac{1}{4}$ and $1\frac{1}{2}$	13.6	6.8	4.6	4.6	2.7	5.7	2.7	4.8	2.4	6.0						
$\frac{1}{4}$ and $1\frac{1}{2}$	2.0	1.5	.4	1.0	.5	1.4	.7	.3	.3	.6						
$\frac{1}{4}$ and $1\frac{1}{2}$.2	.2		.2	.3	.1	.2			.2						
$1\frac{1}{4}$ and longer																.1
All lengths ²																
Shorter than $\frac{7}{8}$ ³	1.3	1.4	2.3	5.5	7.5	.7	.8	4.1	.6							
$\frac{7}{8}$ and $\frac{29}{32}$	24.1	27.7	8.0	26.4	37.2	4.0	13.3	11.4	9.2							
$\frac{1}{4}$ and $\frac{31}{32}$	29.2	16.8	11.0	31.7	23.4	8.7	9.7	1.6	11.5	7.7						
1 and $1\frac{1}{2}$	8.8	2.1	4.0	11.6	6.1	2.2	1.2	1.6	4.0	1.2						
$\frac{1}{4}$ and $1\frac{1}{2}$	2.1	.1	.3	1.7	.8	1.0	.1	.2	.2	.1						.1
$\frac{1}{4}$ and $1\frac{1}{2}$.5				.1	.2										
$\frac{1}{4}$ and $1\frac{1}{2}$																
$1\frac{1}{4}$ and longer																
All lengths ²	66.0	48.1	25.6	77.0	75.2	16.1	25.0	4.2	31.3	18.8						
Shorter than $\frac{7}{8}$ ³	1.3	1.4	2.3	5.5	7.5	.7	.8	4.1	.6							
$\frac{7}{8}$ and $\frac{29}{32}$	24.1	27.7	8.0	26.4	37.2	4.0	13.3	11.4	9.2							
$\frac{1}{4}$ and $\frac{31}{32}$	29.2	16.8	11.0	31.7	23.4	8.7	9.7	1.6	11.5	7.7						
1 and $1\frac{1}{2}$	8.8	2.1	4.0	11.6	6.1	2.2	1.2	1.6	4.0	1.2						
$\frac{1}{4}$ and $1\frac{1}{2}$	2.1	.1	.3	1.7	.8	1.0	.1	.2	.2	.1						.1
$\frac{1}{4}$ and $1\frac{1}{2}$.5				.1	.2										
$1\frac{1}{4}$ and $1\frac{1}{2}$																
$1\frac{1}{4}$ and longer																

TEXAS

Staple length (inches)	Total, crop of—					Ginned prior to Oct. 1										
	1932		1931		1930	1929		1928	1932		1931		1930	1929		1928
	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	
All lengths ²	4,307.4	5,008.8	3,886.1	3,803.2	4,941.5	1,595.2	2,208.3	2,584.7	2,128.6	2,430.9						
Shorter than $\frac{7}{8}$ ³	414.3	446.5	475.5	974.2	794.6	121.9	75.1	271.8	318.6	312.6						
$\frac{7}{8}$ and $\frac{29}{32}$	1,918.6	2,123.4	1,330.4	1,260.5	1,856.8	686.5	624.1	744.7	600.8	880.5						
$\frac{1}{4}$ and $\frac{31}{32}$	1,554.8	1,719.3	1,363.2	955.9	1,557.1	659.2	985.5	979.6	702.6	785.6						
1 and $1\frac{1}{2}$	362.3	678.3	586.3	480.8	586.8	110.4	529.6	490.5	406.1	360.7						
$\frac{1}{4}$ and $1\frac{1}{2}$	37.5	86.9	101.0	103.2	113.0	9.9	48.8	64.8	79.5	66.4						
$\frac{1}{4}$ and $1\frac{1}{2}$	19.7	13.9	25.7	26.0	26.9	8.1	5.0	20.3	19.5	15.1						
$\frac{1}{4}$ and $1\frac{1}{2}$.2	.4	3.9	2.4	5.8	.2	.2	3.9	1.4	3.7						
$1\frac{1}{4}$ and longer		.1	.1	.2	.6				.1	.1						
All lengths ²	1,362.8	1,766.1	709.7	1,016.0	1,434.1	944.3	656.0	393.4	372.9	637.2						
Shorter than $\frac{7}{8}$ ³	77.2	186.2	93.9	391.9	175.2	54.2	77.9	71.6	148.0	65.4						
$\frac{7}{8}$ and $\frac{29}{32}$	623.3	950.9	302.2	398.2	528.8	419.6	335.5	194.1	156.2	256.0						
$\frac{1}{4}$ and $\frac{31}{32}$	502.4	500.4	235.8	165.5	522.4	348.4	188.5	98.4	42.9	159.3						
1 and $1\frac{1}{2}$	137.4	100.7	57.3	41.8	172.6	107.2	39.2	14.5	17.9	39.5						
$\frac{1}{4}$ and $1\frac{1}{2}$	12.8	22.7	18.4	13.9	24.6	13.0	11.9	11.7	6.3	14.5						
$\frac{1}{4}$ and $1\frac{1}{2}$	9.7	4.9	2.1	4.2	8.4	1.9	3.9	3.1	1.2	2.3						.2
$\frac{1}{4}$ and $1\frac{1}{2}$.2		.5	1.0				.3	.1						
$1\frac{1}{4}$ and longer		.1			.2					.1						
All lengths ²																

See footnotes at end of table.

TABLE 7.—*Staple length of American upland cotton: Quantities ginned in the United States during specified periods,¹ by States, crops of 1932, 1931, 1930, 1929, and 1928—Continued*

TEXAS—Continued

Staple length (inches)	Ginned Dec. 1-Jan. 15, crop of—					Ginned after Jan. 15, crop of—				
	1932	1931	1930	1929	1928	1932	1931	1930	1929	1928
	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales
All lengths ²	202.0	256.7	166.5	240.9	403.8	143.1	120.8	31.8	44.8	135.5
Shorter than $\frac{7}{8}$ ³	63.7	54.2	29.3	90.4	152.8	97.3	53.1	8.9	19.3	88.6
$\frac{7}{8}$ and $2\frac{1}{3}$ ²	150.4	158.4	77.1	88.1	156.7	39.8	54.5	12.3	17.2	28.8
$\frac{7}{8}$ ² and $3\frac{1}{3}$ ²	39.9	35.2	42.7	39.1	74.0	4.9	9.7	6.7	5.8	15.8
1 and $1\frac{1}{2}$ ²	6.2	6.1	12.7	13.3	12.1	1.1	2.7	2.3	1.7	1.9
$1\frac{1}{2}$ ² and $1\frac{3}{4}$ ²	1.8	2.7	4.5	2.7	7.1	.8	1.6	.8	.8	.4
$1\frac{1}{2}$ ² and $1\frac{5}{8}$ ²1	.2	1.1	1.1						
$1\frac{1}{2}$ ² and $1\frac{1}{2}$ ²2							
$1\frac{1}{2}$ ² and longer.....										

VIRGINIA

Staple length (inches)	Total, crop of—					Ginned prior to Oct. 1				
	1932	1931	1930	1929	1928	1932	1931	1930	1929	1928
	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales
All lengths ²	31.3	42.5	42.7	48.0	44.8	6.1	6.2	7.9	0.3	1.0
Shorter than $\frac{7}{8}$ ³	1.5	1.7	3.9	3.3	2.8					.6
$\frac{7}{8}$ and $2\frac{1}{3}$ ²	19.4	22.0	29.6	34.1	37.1	3.9	3.2	6.4	.2	.4
$\frac{7}{8}$ ² and $3\frac{1}{3}$ ²	8.7	16.2	8.4	9.7	4.7	1.6	2.8	.5	.1	
1 and $1\frac{1}{2}$ ²	1.5	1.7	.7	.8	.2	.5	.2			
$1\frac{1}{2}$ ² and $1\frac{3}{4}$ ²2		.1	.1		.1				
$1\frac{1}{2}$ ² and $1\frac{1}{2}$ ²										
$1\frac{1}{2}$ ² and $1\frac{5}{8}$ ²										
$1\frac{1}{2}$ ² and longer.....										

Staple length (inches)	Ginned during October					Ginned during November				
	1932	1931	1930	1929	1928	1932	1931	1930	1929	1928
	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales
All lengths ²	11.2	22.1	22.2	16.8	19.7	9.0	11.3	9.1	18.6	15.9
Shorter than $\frac{7}{8}$ ³2	.6	2.8	2.1	1.2	.9	1.1	.1	.7	.3
$\frac{7}{8}$ and $2\frac{1}{3}$ ²	5.9	10.7	13.4	11.7	16.1	5.7	7.7	6.9	13.6	13.5
$\frac{7}{8}$ ² and $3\frac{1}{3}$ ²	4.3	9.0	5.3	2.5	2.3	2.2	2.4	2.0	4.1	2.0
1 and $1\frac{1}{2}$ ²7	1.2	.6	.4	.1	.2	.1	.1	.2	.1
$1\frac{1}{2}$ ² and $1\frac{3}{4}$ ²1		.1	.1						
$1\frac{1}{2}$ ² and $1\frac{1}{2}$ ²										
$1\frac{1}{2}$ ² and $1\frac{5}{8}$ ²										
$1\frac{1}{2}$ ² and longer.....										

Staple length (inches)	Ginned Dec. 1-Jan. 15, crop of—					Ginned after Jan. 15, crop of—				
	1932	1931	1930	1929	1928	1932	1931	1930	1929	1928
	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales
All lengths ²	3.7	2.2	2.4	10.2	6.0	1.3	0.7	1.1	2.1	2.2
Shorter than $\frac{7}{8}$ ³4			.5	.3					.4
$\frac{7}{8}$ and $2\frac{1}{3}$ ²	2.8	1.0	1.8	7.5	5.3	1.1	.3	1.1	1.1	1.8
$\frac{7}{8}$ ² and $3\frac{1}{3}$ ²4	1.0	.6	2.0	.4	.2	.4			
1 and $1\frac{1}{2}$ ²1	.2		.2						
$1\frac{1}{2}$ ² and $1\frac{3}{4}$ ²										
$1\frac{1}{2}$ ² and $1\frac{1}{2}$ ²										
$1\frac{1}{2}$ ² and $1\frac{5}{8}$ ²										
$1\frac{1}{2}$ ² and longer.....										

See footnotes at end of table.

TABLE 7.—*Staple length of American upland cotton: Quantities ginned in the United States during specified periods,¹ by States, crops of 1932, 1931, 1930, 1929, and 1928—Continued*ALL OTHER STATES⁴

Staple length (inches)	Total, crop of—					Ginned prior to Oct. 1				
	1932	1931	1930	1929	1928	1932	1931	1930	1929	1928
	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales
All lengths ²	14.0	11.7	6.4	8.9	6.2	2.0	0.6	1.6	0.5	0.2
Shorter than $\frac{7}{8}$ ³1	.5	.2	.8				.1	
$\frac{7}{8}$ and $2\frac{1}{2}$	1.2	3.0	1.0	.1	.2	.3			.2	
$1\frac{1}{2}$ and $3\frac{1}{2}$	5.4	3.7	1.4	1.1	.6	.5	.2	.3	.1	
1 and $1\frac{1}{2}$	4.8	2.2	1.8	2.2	.9	.7	.1	.5	.1	
$1\frac{1}{2}$ and $3\frac{1}{2}$	1.9	1.8	1.4	2.1	1.7	.3	.2	.4	.1	
$1\frac{1}{2}$ and $1\frac{1}{2}$6	.8	.3	2.2	1.0	.2	.1	.1	.1	.1
$1\frac{1}{2}$ and $1\frac{1}{2}$1				.7				.1	
1 $\frac{1}{2}$ and longer.....				.1	.3					
All lengths ²	5.5	3.9	2.7	2.9	1.5	2.7	3.3	1.2	2.3	1.4
Shorter than $\frac{7}{8}$ ³1					.1		
$\frac{7}{8}$ and $2\frac{1}{2}$2	.9	.6			.1	.7	.1	.1	
$1\frac{1}{2}$ and $3\frac{1}{2}$	1.5	1.1	.5			1.2	.7	.3	.1	.1
1 and $1\frac{1}{2}$	2.1	.8	.6	.3	.1	1.2	.6	.5	.5	.1
$1\frac{1}{2}$ and $3\frac{1}{2}$	1.3	.6	.7	.7	.4	.2	.9	.3	.8	.5
$1\frac{1}{2}$ and $1\frac{1}{2}$3	.4	.2	1.2	.4		.3		.6	.4
$1\frac{1}{2}$ and $1\frac{1}{2}$1	.1			.6	.4			.2	.2
1 $\frac{1}{2}$ and longer.....					.1	.2				.1
All lengths ²	2.6	2.0	0.6	1.4	1.8	1.2	1.9	0.3	1.8	1.3
Shorter than $\frac{7}{8}$ ³2	.2	.2			.1		.6
$\frac{7}{8}$ and $2\frac{1}{2}$5	.4	.1		.1	.1	1.0			.1
$1\frac{1}{2}$ and $3\frac{1}{2}$	1.5	1.0	.2	.1	.2	.7	.1	.8	.3	
1 and $1\frac{1}{2}$5	.5	.1	.5	.6	.3	.2	.8	.1	
$1\frac{1}{2}$ and $3\frac{1}{2}$1	.1		.3	.6		.1		.2	.2
$1\frac{1}{2}$ and $1\frac{1}{2}$3	.1					
1 $\frac{1}{2}$ and $1\frac{1}{2}$										
All lengths ²	2.6	2.0	0.6	1.4	1.8	1.2	1.9	0.3	1.8	1.3

¹ Periods are those used by the Bureau of the Census.² As reported by the Bureau of the Census.³ Untenderable in settlement of futures contracts made subject to sec. 5 of the United States Cotton Futures Act and the regulations of the Secretary of Agriculture thereunder.⁴ Includes Illinois, Kansas, and Kentucky.

TABLE 8.—*Staple length of American-Egyptian cotton: Quantities ginned in the United States during specified periods,¹ crops of 1932, 1931, 1930, 1929, and 1928*

[Quantities are given in running bales]

Staple length (inches)	Total, crop of—					Ginned prior to Oct. 1				
	1932	1931	1930	1929	1928	1932	1931	1930	1929	1928
All lengths ²	1,000 bales 8.3	1,000 bales 13.7	1,000 bales 23.3	1,000 bales 28.8	1,000 bales 28.3	1,000 bales 1.1	1,000 bales 2.3	1,000 bales 3.0	1,000 bales 3.4	1,000 bales 6.2
Shorter than 1½.....	1.7				.7					.2
1½ and 1½½.....	2.9	2.4	2.6	5.8	18.4	.1	.8	.6	.4	2.2
1½ and 1½½.....	2.6	8.4	16.2	17.1	12.5	.5	1.4	2.7	1.8	3.0
1½ to 1½½.....	1.1	2.9	4.6	6.0	1.6	.5	.1	.6	1.2	.8
1½ and longer.....				.4	.1					
Staple length (inches)	Ginned during October					Ginned during November				
	1932	1931	1930	1929	1928	1932	1931	1930	1929	1928
All lengths ²	1,000 bales 2.7	1,000 bales 3.1	1,000 bales 6.6	1,000 bales 7.4	1,000 bales 7.2	1,000 bales 2.4	1,000 bales 1.4	1,000 bales 5.0	1,000 bales 7.2	1,000 bales 6.4
Shorter than 1½.....	.1				.3	.7				
1½ and 1½½.....	.7	.2	.2	.9	3.0	1.1				1.8
1½ and 1½½.....	1.4	1.9	4.9	3.4	3.3	.5	.7	3.3	4.6	4.0
1½ to 1½½.....	.5	1.0	1.5	2.7	.5	.1	.7	1.7	.8	.2
1½ and longer.....				.4	.1					
Staple length (inches)	Ginned Dec. 1-Jan. 15, crop of—					Ginned after Jan. 15, crop of—				
	1932	1931	1930	1929	1928	1932	1931	1930	1929	1928
All lengths ²	1,000 bales 1.2	1,000 bales 4.1	1,000 bales 5.7	1,000 bales 7.5	1,000 bales 5.6	1,000 bales 0.9	1,000 bales 2.8	1,000 bales 2.1	1,000 bales 3.3	1,000 bales 2.9
Shorter than 1½.....	.3				.1	.6				.1
1½ and 1½½.....	.7	.4	.8	.9	3.4	.3	1.0	.9	1.3	2.6
1½ and 1½½.....	.2	2.8	4.1	5.8	2.0		1.6	1.2	2.0	.2
1½ to 1½½.....	.9	.8	1.3	.1		.2				
1½ and longer.....										

¹ Periods are those used by the Bureau of the Census.² As reported by the Bureau of the Census.

86 STATISTICAL BULLETIN 47, U. S. DEPT. OF AGRICULTURE

TABLE 9.—*Staple length of American upland cotton ginned in designated districts of Alabama, crops of 1932, 1931, 1930, 1929, and 1928 (fig. 2)*

[Quantities are given in running bales, except that round bales are counted as half bales. Linters are not included]

Staple length (inches)	Crop year	Total	District 1	District 2	District 3	District 4	District 5	District 6			
		1,000 bales	Per- cent								
All lengths...	1932	933.7	100.0	215.2	100.0	182.3	100.0	94.5	100.0	364.0	100.0
	1931	1,385.0	100.0	283.8	100.0	268.8	100.0	115.5	100.0	594.9	100.0
	1930	1,444.9	100.0	248.0	100.0	274.9	100.0	118.2	100.0	680.6	100.0
	1929	1,307.7	100.0	278.5	100.0	289.3	100.0	115.2	100.0	541.9	100.0
	1928	1,096.6	100.0	253.5	100.0	235.0	100.0	109.9	100.0	422.1	100.0
										41.1	100.0
Shorter than $\frac{7}{8}$...	1932	137.3	14.7	42.9	19.9	15.9	8.7	3.4	3.6	66.0	18.1
	1931	136.5	9.9	33.6	11.8	30.5	11.4	12.4	10.8	49.3	8.3
	1930	553.3	33.3	111.7	45.1	116.8	42.5	29.1	24.6	274.8	40.4
	1929	585.6	44.8	131.4	47.2	164.2	61.0	48.7	42.3	205.6	38.0
	1928	264.2	24.1	61.5	24.3	62.9	26.8	17.5	15.9	107.2	25.4
										9.2	22.4
$\frac{7}{8}$ and $\frac{21}{32}$...	1932	657.5	70.4	129.0	60.0	127.8	70.1	67.5	71.4	269.3	74.0
	1931	1,044.7	75.4	189.2	66.7	199.0	74.0	95.2	82.4	465.7	78.3
	1930	802.2	55.5	121.1	48.8	136.6	49.7	79.6	67.4	374.1	55.0
	1929	689.0	52.7	141.8	50.9	100.6	37.4	63.3	54.9	320.6	59.2
	1928	762.6	69.5	176.0	69.4	159.3	67.8	79.9	72.7	291.4	69.0
										28.7	89.8
$\frac{15}{16}$ and $\frac{31}{32}$...	1932	108.7	11.7	32.6	15.1	30.2	16.6	19.1	20.2	22.8	6.3
	1931	168.7	12.2	47.0	16.6	34.1	12.7	7.4	6.4	66.4	11.1
	1930	77.1	5.3	13.9	5.6	18.7	6.8	9.0	7.6	25.2	3.7
	1929	28.3	2.2	4.9	1.8	3.8	1.4	3.0	2.6	12.6	2.3
	1928	49.8	4.5	9.2	3.6	9.1	3.9	10.2	9.3	16.8	4.0
										2.8	6.8
1 and $1\frac{1}{2}$...	1932	28.5	2.8	9.8	4.6	6.9	3.8	4.2	4.5	4.9	1.3
	1931	28.0	2.0	11.2	3.9	4.8	1.8	5.5	4.4	9.7	1.6
	1930	9.2	.6	1.1	.4	2.5	.9	.5	.4	4.3	.6
	1929	3.3	.2	.3	.1	.6	.2	.2	2.1	.3	.1
	1928	17.6	1.6	6.5	2.6	3.4	1.4	2.0	1.8	5.3	1.2
										.7	.1
$1\frac{1}{2}$ and $1\frac{3}{4}$...	1932	3.1	.3	.8	.4	1.3	.7	.3	.7	.2	
	1931	5.8	.4	2.5	.9	.4	.1				
	1930	2.5	.2	.2	.1	.3	.1				
	1929	1.4	.1	.1	(3)	.1	(3)				
	1928	1.8	.2	.2	1	.3	.1	.1			
$1\frac{3}{4}$ and $1\frac{5}{8}$...	1932	.5	.1	.1	(3)	.2	.1				
	1931	1.3	.1	.3	.1						
	1930	.6	(3)								
	1929	.1	(3)								
	1928	.5	(3)	.1	(3)						
		.1	(3)								
$1\frac{1}{4}$ and longer...	1931										
	1930										
	1929										
	1928										

¹ As reported by the Bureau of the Census.

² Untenderable in settlement of futures contracts made subject to sec. 5 of the United States Cotton Futures Act and the regulations of the Secretary of Agriculture thereunder.

³ Less than 0.05 percent.

COTTON GRADE AND STAPLE DISTRICTS OF ALABAMA




FIGURE 2.

TABLE 10.—*Staple length of American upland cotton ginned in designated districts of Arkansas, crops of 1932, 1931, 1930, 1929, and 1928 (fig. 3)*

[Quantities are given in running bales, except that round bales are counted as half bales. Linters are not included]

Staple length (inches)	Crop year	Total	District 1	District 2	District 3	District 4	District 5
		1,000 bales	Per- cent	1,000 bales	Per- cent	1,000 bales	Per- cent
All lengths-----	1932	1,283.4	100.0	43.3	100.0	174.3	100.0
	1931	1,836.1	100.0	46.7	100.0	269.4	100.0
	1930	1,863.5	100.0	22.4	100.0	128.9	100.0
	1929	1,395.9	100.0	48.6	100.0	201.9	100.0
	1928	1,216.2	100.0	38.3	100.0	237.2	100.0
Shorter than $\frac{7}{8}$ ¹ -----	1932	25.8	2.0	7.1	16.4	6.4	3.7
	1931	20.9	1.1	1.6	3.4	6.5	2.4
	1930	76.9	8.9	8.4	37.5	28.5	22.5
	1929	177.9	12.7	24.8	53.2	59.4	29.4
	1928	127.0	10.4	12.2	31.9	45.8	19.3
	1932	185.9	14.5	11.6	26.8	55.0	31.6
	1931	307.4	16.7	15.5	33.2	74.1	27.5
	1930	196.1	22.7	8.4	37.5	46.1	36.3
	1929	393.4	28.2	17.8	38.2	81.1	40.2
	1928	359.4	29.6	18.6	48.5	97.7	41.2
	1932	375.0	29.2	7.4	17.1	79.9	45.8
	1931	549.6	29.9	19.9	42.6	105.0	39.0
	1930	252.9	29.3	4.8	21.4	34.6	22.3
	1929	379.1	27.2	3.3	7.1	42.1	20.9
	1928	290.5	23.9	5.6	14.6	60.0	25.3
	1932	360.4	28.1	11.9	27.5	30.4	17.4
	1931	614.2	33.5	8.0	17.2	71.5	26.6
	1930	206.8	24.0	.7	3.1	14.0	11.0
	1929	261.9	18.8	.7	1.5	15.5	7.7
	1928	223.6	18.4	1.4	3.7	25.7	10.8
	1932	196.8	15.3	5.1	11.8	2.5	1.4
	1931	247.0	13.5	1.6	3.4	10.8	4.0
	1930	106.6	12.3	.1	.5	3.4	2.7
	1929	128.0	9.2	—	—	3.3	1.6
	1928	130.6	10.7	.1	.3	6.2	2.6
	1932	120.4	9.4	.2	.4	.1	.2
	1931	81.4	4.4	.1	.2	1.4	.5
	1930	22.4	2.6	—	—	.3	.2
	1929	45.4	3.2	—	—	.3	.1
	1928	65.9	5.4	—	—	1.3	.5
	1932	17.3	1.4	—	—	—	—
	1931	14.9	.8	—	—	.1	(?)
	1930	1.7	.2	—	—	—	—
	1929	8.6	.6	—	—	.1	(?)
	1928	16.9	1.4	.2	.5	.4	.2
	1932	1.8	.1	—	—	—	—
	1931	.7	(?)	—	—	—	—
	1930	.1	(?)	—	—	—	—
	1929	1.6	.1	—	—	—	—
	1928	2.3	.2	.2	.5	.1	(?)

¹ As reported by the Bureau of the Census.² Untenderable in settlement of futures contracts made subject to sec. 5 of the United States Cotton Futures Act and the regulations of the Secretary of Agriculture thereunder.³ Less than 0.05 percent.




FIGURE 3.

TABLE 11.—*Staple length of American upland cotton ginned in designated districts of Georgia, crops of 1932, 1931, 1930, 1929, and 1928 (fig. 4)*

[Quantities are given in running bales, except that round bales are counted as half bales. Linters are not included]

Staple length (inches)	Crop year	Total		District 1		District 2		District 3		District 4	
		1,000 bales	Per- cent								
All lengths	1932	1,861.8	100.0	84.9	100.0	424.5	100.0	51.4	100.0	301.0	100.0
	1931	1,393.7	100.0	103.3	100.0	567.3	100.0	113.3	100.0	609.8	100.0
	1930	1,597.5	100.0	97.8	100.0	630.3	100.0	171.2	100.0	608.2	100.0
	1929	1,339.8	100.0	111.6	100.0	586.3	100.0	130.2	100.0	511.7	100.0
	1928	1,053.2	100.0	102.1	100.0	518.4	100.0	92.6	100.0	340.1	100.0
		43.1	*5.0	2.5	.2.9	8.0	1.9	2.0	3.9	30.6	10.2
Shorter than $\frac{7}{8}$	1932	69.4	5.0	10.6	10.2	21.5	3.8	3.7	3.3	33.6	5.5
	1931	237.1	14.8	27.0	27.6	58.0	9.2	29.9	17.5	122.2	17.5
	1930	282.2	21.0	25.7	23.0	123.1	21.0	24.9	19.1	108.5	21.2
	1929	129.5	12.3	6.3	6.2	24.6	4.7	13.8	14.9	84.8	24.9
	1928	592.8	68.8	57.6	67.9	254.4	59.8	38.6	75.0	242.2	80.4
	1931	1,015.0	72.9	75.7	73.3	306.3	69.8	84.8	74.8	458.8	75.2
$\frac{7}{8}$ and $\frac{29}{32}$	1930	1,096.8	68.7	54.9	56.1	417.0	66.2	125.3	73.2	499.6	71.6
	1929	918.8	68.6	73.8	66.1	386.3	65.9	96.7	74.3	302.0	70.7
	1928	743.9	70.6	76.0	74.0	371.9	71.7	70.7	76.3	225.7	60.4
	1932	163.1	18.9	14.5	17.1	120.7	28.4	6.2	12.1	21.7	7.2
	1931	242.1	17.4	12.7	12.3	121.7	21.5	18.7	16.5	89.0	14.6
	1930	218.4	13.7	13.0	13.3	129.3	20.5	14.1	8.2	62.0	8.9
$\frac{15}{16}$ and $\frac{31}{32}$	1929	113.7	8.5	10.8	9.7	64.4	11.0	6.7	5.2	31.8	6.2
	1928	144.1	13.7	15.0	14.7	90.0	19.1	7.2	7.8	22.9	6.7
	1932	53.9	6.3	9.1	10.7	35.5	8.4	3.9	7.6	5.4	1.8
	1931	54.8	3.9	3.7	3.6	24.6	4.3	4.2	3.7	22.3	1.3
	1930	36.3	2.3	2.5	2.6	22.2	3.5	1.5	9.9	10.1	1.4
	1929	21.3	1.6	1.1	1.0	11.1	1.9	1.8	1.2	7.5	1.5
$\frac{13}{16}$ and $\frac{13}{16}$	1928	28.3	2.7	4.5	4.4	18.5	3.6	.8	.9	4.5	.5
	1932	6.7	.8	1.2	1.4	4.1	.6	1.2	.8	.3	
	1931	10.4	.7	.5	.5	2.8	.5	1.7	1.5	5.4	.9
	1930	7.3	.4	.4	.4	3.3	.5	.4	.2	3.2	.5
	1929	3.6	.3	.1	.1	1.4	.2	.2	.2	1.8	.4
	1928	5.4	.5	.6	.6	3.2	.6	.1	.1	1.5	.4
$\frac{13}{16}$ and $\frac{15}{16}$	1932	1.9	.2			1.5	.4	.1	.2	.3	.1
	1931	1.3	.1	.1	.1	.4	.1	.2	.2	.6	.1
	1930	1.4	.1			.5	.1			.9	.1
	1929	.2	(?)	.1	.1					.1	(?)
	1928	1.4	.1	.1	.1	.8	.2			.5	.2
	1932	.3	(?)			.3	.1				
$\frac{13}{16}$ and $\frac{17}{16}$	1931										
	1930	.2	(?)							.2	(?)
	1929										
	1928	.6	.1			.4	.1			.2	.1
	1932										
	1931	.1	(?)							.1	(?)
$\frac{1}{4}$ and longer	1930										
	1929										
	1928										
	1928										

¹ As reported by the Bureau of the Census.

² Untenderable in settlement of futures contracts made subject to sec. 5 of the United States Cotton Futures Act and the regulations of the Secretary of Agriculture thereunder.

³ Less than 0.05 percent.

COTTON GRADE AND STAPLE DISTRICTS OF GEORGIA




FIGURE 4.

TABLE 12.—*Staple length of American upland cotton ginned in designated districts of Louisiana, crops of 1932, 1931, 1930, 1929, and 1928 (fig. 5)*

[Quantities are given in running bales, except that round bales are counted as half bales. Linters are not included]

Staple length (inches)	Crop year	Total	District 1	District 2	District 3	District 4			
		1,000 bales	Per- cent	1,000 bales	Per- cent	1,000 bales	Per- cent	1,000 bales	Per- cent
All lengths	1932	589.5	100.0	150.6	100.0	52.0	100.0	379.0	100.0
	1931	876.6	100.0	208.8	100.0	77.7	100.0	559.5	100.0
	1930	1,704.8	100.0	143.9	100.0	92.1	100.0	433.1	100.0
	1929	1,797.7	100.0	216.3	100.0	76.5	100.0	481.5	100.0
	1928	1,685.9	100.0	185.5	100.0	79.8	100.0	399.4	100.0
Shorter than $\frac{7}{8}$	1932	62.1	10.4	53.6	35.6	.1	.2	4.6	1.2
	1931	58.6	6.7	45.8	22.2	.2	.3	7.8	1.4
	1930	91.3	13.0	59.3	41.2	2.7	2.9	21.3	4.9
	1929	110.6	13.9	70.7	32.7	3.0	3.9	33.9	7.0
	1928	88.3	12.9	42.5	22.9	3.5	4.4	39.3	8.8
$\frac{7}{8}$ and $2\frac{1}{2}$	1932	131.0	21.8	64.3	42.7	8.4	16.1	48.8	12.9
	1931	157.3	17.9	85.9	41.5	7.6	9.8	51.4	9.2
	1930	180.1	22.7	44.7	31.1	17.4	18.9	84.6	19.5
	1929	205.5	25.8	89.2	41.2	15.0	19.6	92.9	19.3
	1928	203.1	29.7	83.6	45.1	17.3	21.7	95.4	23.9
$1\frac{1}{2}$ and $3\frac{1}{2}$	1932	152.6	25.4	23.6	15.6	22.1	42.5	103.6	27.3
	1931	203.6	23.2	45.0	21.8	20.9	28.9	128.1	22.9
	1930	167.4	23.8	24.8	17.2	26.7	29.0	108.3	25.0
	1929	212.7	26.7	44.4	20.5	31.5	41.2	130.9	27.2
	1928	178.6	26.0	39.0	21.5	26.2	32.8	105.9	26.5
$1\frac{1}{2}$ and $3\frac{1}{2}$	1932	156.4	26.1	8.1	5.4	19.3	37.1	127.9	33.8
	1931	283.0	32.3	27.2	13.2	35.6	45.8	216.0	38.6
1 and $1\frac{1}{2}$	1930	186.8	26.5	13.8	9.6	35.8	38.9	132.1	30.5
	1929	170.4	21.3	11.1	5.1	22.6	29.5	132.5	27.5
	1928	127.6	18.6	14.5	7.8	20.6	25.8	89.4	22.4
	1932	68.7	11.5	.8	.5	1.9	3.7	65.8	17.4
	1931	133.1	15.2	2.7	1.3	12.3	15.8	116.7	20.9
$1\frac{1}{2}$ and $1\frac{3}{2}$	1930	67.2	9.5	1.2	.8	8.0	8.7	56.9	13.1
	1929	63.7	8.0	.8	.4	3.9	5.1	57.5	12.0
	1928	52.1	7.6	2.6	1.4	8.6	10.8	39.9	10.0
	1932	28.2	4.4	.1	.1	.2	.4	25.9	6.8
	1931	35.4	4.0	.1	(3)	.9	1.2	34.2	6.1
	1930	28.0	4.0	.1	.1	1.5	1.6	28.0	6.0
	1929	29.6	3.7	.1	(3)	.5	.7	28.6	5.9
	1928	27.7	4.0	1.2	.6	3.3	4.1	22.7	5.7
$1\frac{1}{2}$ and $1\frac{3}{2}$	1932	2.4	.4					2.4	.6
	1931	5.3	.6	.1	(3)	.1	.1	5.1	.9
	1930	3.8	.5					3.7	.9
	1929	5.0	.6					5.0	1.0
	1928	7.8	1.1	1.1	.6	.3	.4	6.2	1.6
	1932	.1	(3)	.1	.1			.2	(3)
	1931	.3	(3)			.1	.1	.2	
	1930	.2	(3)					.2	.1
	1929	.2	(3)					.2	(3)
	1928	.7	.1	.1	.1			.6	.1
$1\frac{1}{2}$ and longer									

¹ As reported by the Bureau of the Census.

² Untenderable in settlement of futures contracts made subject to sec. 5 of the United States Cotton Futures Act and the regulations of the Secretary of Agriculture thereunder.

³ Less than 0.05 percent.




FIGURE 5.

TABLE 13.—*Staple length of American upland cotton ginned in designated districts of Mississippi, crops of 1932, 1931, 1930, 1929, and 1928 (fig. 6)*

[Quantities are given in running bales, except that round bales are counted as half bales. Linters are not included]

Staple length (inches)	Crop year	Total	District 1	District 2	District 3	District 4			
		1,000 bales	Per- cent	1,000 bales	Per- cent	1,000 bales	Per- cent	1,000 bales	Per- cent
All lengths	1932	1,161.2	100.0	489.4	100.0	301.4	100.0	333.7	100.0
	1931	1,719.4	100.0	800.5	100.0	382.7	100.0	400.5	100.0
	1930	1,458.5	100.0	535.0	100.0	334.2	100.0	547.5	100.0
	1929	1,876.0	100.0	877.8	100.0	400.5	100.0	546.4	100.0
	1928	1,482.0	100.0	671.7	100.0	310.9	100.0	434.1	100.0
Shorter than $\frac{7}{8}$ ²	1932	32.2	2.8	.1	(3)	1.5	.5	20.7	8.0
	1931	65.4	3.8	.3	(3)	2.9	.8	59.2	12.1
	1930	114.9	7.9	.1	(3)	14.0	4.2	96.9	17.7
	1929	180.7	10.1	.3	(3)	18.0	4.5	167.2	30.6
	1928	264.3	17.4	1.1	2	24.4	7.6	215.3	49.6
$\frac{7}{8}$ and $2\frac{1}{2}$ ²	1932	148.0	12.8	.3	(3)	14.3	4.8	124.8	37.4
	1931	285.4	16.6	2.6	3	39.6	10.3	224.9	46.8
	1930	299.2	20.5	1.8	3	45.6	13.7	236.3	43.2
	1929	352.3	18.8	2.3	3	83.6	20.9	240.7	45.7
	1928	258.9	17.7	8.1	1.2	67.2	21.0	166.7	38.4
$1\frac{1}{16}$ and $2\frac{1}{2}$ ²	1932	162.3	14.0	1.4	3	45.3	15.0	103.7	31.1
	1931	238.0	13.8	8.4	1.1	83.4	21.8	129.8	20.5
	1930	264.5	18.1	15.6	2.9	78.3	23.4	155.1	28.3
	1929	227.0	12.1	8.9	1.0	110.2	27.5	91.9	16.8
	1928	111.9	7.7	16.7	2.5	55.5	17.3	35.3	8.1
1 and $1\frac{1}{2}$ ²	1932	170.9	14.7	21.3	4.2	88.4	29.4	50.8	17.0
	1931	234.8	13.7	65.6	8.2	102.0	26.7	60.6	12.4
	1930	209.4	14.4	65.9	12.3	91.5	27.4	46.3	8.5
	1929	202.7	10.8	69.2	7.9	97.6	24.4	20.6	4.9
	1928	127.8	8.7	59.4	8.8	55.3	17.3	12.0	2.8
$1\frac{1}{16}$ and $1\frac{3}{4}$ ²	1932	271.5	23.4	174.8	35.0	82.0	27.2	14.4	4.3
	1931	311.0	18.1	214.3	26.8	81.9	21.4	13.6	2.8
	1930	312.9	21.5	225.8	42.2	74.9	22.4	11.1	2.0
	1929	388.9	20.7	321.3	36.6	54.4	13.6	9.1	1.7
	1928	259.8	17.8	204.2	30.4	51.7	16.2	3.6	.8
$1\frac{1}{8}$ and $1\frac{1}{2}$ ²	1932	328.2	28.2	262.7	52.6	61.6	20.4	3.9	1.2
	1931	368.1	21.4	315.1	39.4	50.6	13.2	2.2	.4
	1930	210.6	14.8	188.6	35.3	26.1	7.8	1.8	.3
	1929	412.4	22.0	379.1	43.2	30.5	7.6	1.8	.3
	1928	305.0	20.9	264.2	30.4	40.7	12.7	1.0	.2
$1\frac{1}{8}$ and $1\frac{1}{2}$ ²	1932	45.3	3.9	37.0	7.4	7.9	2.6	.4	.1
	1931	188.7	11.0	169.2	21.1	19.3	5.0	.2	(3)
	1930	30.4	2.7	35.9	6.7	3.5	1.0		
	1929	97.7	5.2	91.9	10.5	5.7	1.4	.1	(3)
	1928	121.7	8.3	101.0	15.0	20.4	6.4	.2	.1
$1\frac{1}{4}$ and longer	1932	2.2	.2	1.8	.4	.4	.1		
	1931	28.0	1.6	25.0	3.1	3.0	.8		
	1930	1.6	.1	1.3	.3	.3	.1		
	1929	5.3	.3	4.8	.5	.5	.1		
	1928	21.7	1.5	17.0	2.5	4.7	1.5		

¹ As reported by the Bureau of the Census.² Untenderable in settlement of futures contracts made subject to sec. 5 of the United States Cotton Futures Act and the regulations of the Secretary of Agriculture thereunder.³ Less than 0.05 percent.




FIGURE 6.

TABLE 14.—*Staple length of American upland cotton ginned in designated districts of North Carolina, crops of 1932, 1931, 1930, 1929, and 1928 (fig. 7)*

[Quantities are given in running bales, except that round bales are counted as half bales. Linters are not included]

Staple length (inches)	Crop year	Total		District 1		District 2		District 3		District 4	
		1,000 bales	Per- cent								
All lengths	1932	1,680.3	100.0	241.5	100.0	55.9	100.0	333.8	100.0	49.1	100.0
	1931	1,771.2	100.0	318.3	100.0	74.7	100.0	341.2	100.0	37.0	100.0
	1930	1,800.6	100.0	315.8	100.0	73.4	100.0	360.7	100.0	50.7	100.0
	1929	1,767.0	100.0	322.4	100.0	68.1	100.0	324.5	100.0	52.0	100.0
	1928	1,800.3	100.0	322.8	100.0	77.9	100.0	300.1	100.0	72.5	100.0
	1932	7.8	1.2	3.9	1.6	.1	.2	3.3	1.0	.5	1.0
Shorter than $\frac{7}{8}$ ^a	1931	7.8	1.0	2.8	.9			4.9	1.4	.1	.3
	1930	40.4	5.0	17.5	5.5	.2	.3	21.1	5.8	1.6	3.2
	1929	81.9	10.7	32.2	10.0	1.6	2.4	34.1	10.5	14.0	26.9
	1928	87.1	10.0	26.2	8.1	1.5	1.9	48.6	12.3	10.8	14.9
	1932	227.8	33.4	104.6	43.3	6.8	12.1	92.2	27.6	24.2	49.3
	1931	340.1	44.1	163.0	51.2	16.5	22.1	145.2	42.5	15.4	41.6
$\frac{7}{8}$ and $2\frac{1}{2}$ ^b	1930	410.2	51.2	174.6	55.3	23.3	31.8	183.3	50.8	29.0	57.1
	1929	479.4	62.5	207.4	64.3	34.2	50.2	210.3	64.8	27.5	52.9
	1928	605.5	69.7	224.9	69.6	46.5	58.7	281.7	71.1	52.4	72.2
	1932	223.6	32.9	77.6	32.1	19.4	34.7	112.2	33.6	14.4	29.3
	1931	264.6	34.3	97.6	30.7	34.8	46.5	117.9	34.6	14.3	38.6
	1930	236.5	29.6	88.6	28.1	33.6	45.8	100.1	27.8	14.2	28.0
$1\frac{1}{16}$ and $3\frac{1}{2}$ ^b	1929	151.2	19.7	62.5	19.4	22.7	33.3	57.8	17.8	8.2	15.8
	1928	129.7	14.9	58.3	18.1	23.2	29.8	41.9	10.6	6.3	8.7
	1932	160.4	23.6	43.1	17.9	23.5	42.0	86.9	26.0	6.9	14.1
	1931	126.0	16.3	41.9	13.2	19.9	26.8	59.0	17.3	5.2	14.1
	1930	81.6	10.2	26.5	8.4	13.1	17.8	37.9	10.5	4.1	8.1
	1929	45.7	6.0	17.2	5.3	8.9	13.1	17.8	5.4	2.0	3.8
$1\frac{1}{16}$ and $1\frac{3}{2}$ ^b	1928	30.5	3.5	9.6	3.0	6.0	7.7	12.9	3.3	2.0	2.8
	1932	47.5	7.0	10.2	4.2	4.9	8.8	30.6	9.2	1.8	3.7
	1931	28.5	3.7	11.6	3.6	3.3	4.4	12.2	3.6	1.4	3.8
	1930	25.6	3.2	7.6	2.4	2.8	3.8	13.8	3.8	1.4	2.8
	1929	7.0	.9	2.4	.7	.7	1.0	3.6	1.1	.3	.6
	1928	9.0	1.0	1.8	.6	.5	.8	6.2	1.6	.5	.7
$1\frac{1}{8}$ and $1\frac{1}{2}$ ^b	1932	11.0	1.6	1.3	.5	1.1	2.0	7.6	2.3	1.0	2.0
	1931	3.8	.5	1.3	.4	.2	.3	1.9	.6	.4	1.1
	1930	5.0	.7	.9	.3	.4	.5	3.9	1.1	.4	.8
	1929	1.5	.2	.6	.2			.9	.3		
	1928	4.9	.6	1.2	.4	.2	.3	3.3	.8	.2	.6
	1932	2.1	.3	.7	.3	.1	.2	1.0	.3	.3	.6
$1\frac{1}{4}$ and $1\frac{1}{2}$ ^b	1931	.4	.1	.1	(2)			.1	(3)	.2	.6
	1930	.7	.1	.1	(2)			.6	.2		
	1929	.3	(3)	.1	(2)			.2	.1		
	1928	2.2	.3	.7	.2			1.3	.3	.2	.3
	1932	.1	(3)	.1	.1						
	1931										
$1\frac{1}{4}$ and longer	1930										
	1929										
	1928	.4	(3)	.1	(2)			.2	(3)	.1	.1

^a As reported by the Bureau of the Census.

^b Untenderable in settlement of futures contracts made subject to sec. 5 of the United States Cotton Futures Act and the regulations of the Secretary of Agriculture thereunder.

^c Less than 0.05 percent.

COTTON GRADE AND STAPLE DISTRICTS OF NORTH CAROLINA




FIGURE 7.

TABLE 15.—*Staple length of American upland cotton ginned in designated districts of Oklahoma, crops of 1932, 1931, 1930, 1929, and 1928 (fig. 8)*

[Quantities are given in running bales, except that round bales are counted as half bales. Linters are not included]

Staple length (inches)	Crop year	Total		District 1		District 2		District 3		District 4		District 5	
		1,000 bales	Per- cent										
All lengths...	1932	1,072.0	100.0	732.8	100.0	260.3	100.0	36.9	100.0	13.2	100.0	28.8	100.0
	1931	1,235.9	100.0	751.8	100.0	361.3	100.0	58.7	100.0	25.3	100.0	38.8	100.0
	1930	1,858.8	100.0	527.4	100.0	260.4	100.0	31.4	100.0	18.4	100.0	21.2	100.0
	1929	1,125.6	100.0	804.5	100.0	246.7	100.0	31.7	100.0	18.9	100.0	23.8	100.0
	1928	1,187.0	100.0	787.4	100.0	311.3	100.0	43.0	100.0	21.3	100.0	24.0	100.0
Shorter than 7/16 ² ...	1932	.01.9	8.6	80.3	11.0	7.9	3.0	1.1	3.0	.2	1.5	2.4	8.3
	1931	137.9	11.2	127.5	17.0	8.4	2.3	.7	1.2	.1	.4	1.2	3.1
	1930	128.7	14.8	111.7	21.2	10.8	4.1	2.0	6.4			2.2	10.4
	1929	321.5	28.6	278.4	34.6	28.2	11.4	6.3	19.9	.3	1.6	8.3	34.9
	1928	182.3	13.7	148.2	18.8	8.9	2.9	1.3	3.0	.1	.5	3.8	15.8
7/16 and 29/32...	1932	535.8	50.0	360.3	49.2	131.6	50.6	19.4	52.6	3.9	29.6	20.6	71.6
	1931	657.4	45.1	384.1	51.1	132.8	36.8	19.8	33.7	2.5	9.9	18.2	40.0
	1930	348.6	40.7	241.6	45.8	83.3	32.0	10.1	32.2	1.0	6.1	12.6	50.4
	1929	500.1	44.4	380.2	47.3	93.0	37.7	12.8	40.4	2.7	14.3	11.4	47.9
	1928	421.7	35.5	325.1	41.3	69.3	22.3	12.2	28.4	2.0	9.4	13.1	54.6
15/16 and 31/32...	1932	386.1	36.0	255.9	34.9	108.2	40.8	13.2	35.8	5.0	37.9	5.8	20.2
	1931	414.5	33.5	201.6	26.8	163.3	45.2	25.8	44.0	7.7	30.4	16.1	41.5
	1930	295.1	34.4	150.2	28.5	122.0	46.9	12.9	41.0	3.8	23.2	6.2	29.3
	1929	207.9	18.5	119.5	14.8	73.0	29.6	7.8	24.6	3.8	20.1	3.8	10.0
	1928	395.5	33.3	236.3	30.0	130.2	41.8	18.9	43.9	4.6	21.6	5.5	22.9
1 and 13/16...	1932	52.4	4.9	32.6	4.4	13.7	5.3	2.6	7.0	3.5	26.5		
	1931	110.8	9.0	36.0	4.8	52.8	14.6	9.9	16.9	9.2	36.4	2.9	7.5
	1930	71.6	8.4	22.3	4.2	39.1	15.0	5.3	16.9	4.7	28.7	.2	.9
	1929	67.1	6.0	22.3	2.8	38.2	14.7	2.7	8.5	5.6	29.6	.3	1.2
	1928	172.3	14.5	72.6	9.2	82.3	26.4	9.7	22.6	6.5	30.5	1.2	5.0
1 1/4 and 13/16...	1932	.5	.5	3.1	.4	.9	.3	.6	1.6	.6	4.5		
	1931	14.0	1.1	2.4	.3	4.0	1.1	2.2	3.7	5.1	20.1	.3	.8
	1930	12.2	1.4	1.6	.3	4.6	1.8	1.0	3.2	5.0	30.4		
	1929	22.6	2.0	3.9	.5	13.0	5.3	1.5	4.7	4.2	22.2		
	1928	25.3	2.1	4.6	.6	15.3	4.9	.7	1.6	4.4	20.7	.3	1.3
1 1/4 and 13/16...	1932	.5	(3)	.5	.1								
	1931	1.3	.1	.2	(3)			.3	.5	.7	2.8	.1	.2
	1930	2.5	.3			.6	.2	.1	.3	1.8	11.0		
	1929	6.1	.5	.2	(3)	3.1	1.2	.6	1.9	2.2	11.7		
	1928	8.2	.7	.4	.1	4.3	1.4	.2	.6	3.2	15.0	.1	.4
1 1/4 and longer...	1930												
	1929												
	1928												

¹ As reported by the Bureau of the Census.² Untenderable in settlement of futures contracts made subject to sec. 5 of the United States Cotton Futures Act and the regulations of the Secretary of Agriculture thereunder.³ Less than 0.05 percent.

COTTON GRADE AND STAPLE DISTRICTS OF OKLAHOMA

- 1 Red lands and plains
- 2 Oklahoma prairies
- 3 Southern Ozarks
- 4 Sandy coastal plains
- 5 Cross timbers




FIGURE 8.

TABLE 16.—*Staple length of American upland cotton ginned in designated districts of South Carolina, crops of 1932, 1931, 1930, 1929, and 1928 (fig. 9)*

[Quantities are given in running bales, except that round bales are counted as half bales. Linters are not included]

Staple length (inches)	Crop year	Total		District 1		District 2		District 3		District 4	
		1,000 bales	Per- cent								
All lengths.....	1932	722.2	100.0	385.8	100.0	67.6	100.0	243.1	100.0	25.7	100.0
	1931	1,010.2	100.0	559.7	100.0	94.0	100.0	310.9	100.0	39.6	100.0
	1930	1,015.3	100.0	540.3	100.0	94.1	100.0	336.3	100.0	44.6	100.0
	1929	833.1	100.0	482.6	100.0	63.9	100.0	251.4	100.0	35.2	100.0
	1928	744.4	100.0	435.2	100.0	72.5	100.0	214.2	100.0	22.5	100.0
Shorter than $\frac{7}{8}$	1932	2.8	.4	1.9	.5	.1	.1	.6	.2	.2	.8
	1931	12.0	1.2	7.8	1.4	.2	.2	3.8	1.2	.2	.5
	1930	26.9	2.6	16.1	3.0	.8	.8	9.0	2.7	1.0	2.2
	1929	93.1	11.2	54.2	11.2	2.9	4.6	30.8	12.3	5.2	14.8
	1928	51.8	7.0	13.0	3.0	5.2	7.2	29.0	13.5	4.6	20.5
$\frac{7}{8}$ and $\frac{29}{32}$	1932	181.2	25.1	107.3	27.8	12.0	17.8	51.8	21.3	10.1	30.3
	1931	395.6	39.1	228.5	40.8	17.6	18.8	131.0	41.3	18.5	46.7
	1930	426.0	42.0	251.2	46.5	21.3	22.6	129.8	38.5	23.7	53.2
	1929	440.4	52.9	277.7	57.5	25.0	30.1	117.5	46.7	20.2	57.4
	1928	409.9	55.0	282.0	60.2	35.4	48.8	97.7	45.6	14.8	65.8
$\frac{15}{16}$ and $\frac{31}{32}$	1932	225.6	31.2	142.3	36.9	13.3	19.7	60.3	24.8	9.7	37.7
	1931	315.2	31.2	194.2	34.7	22.1	23.5	81.4	25.7	17.5	44.2
	1930	288.6	28.4	166.7	30.0	20.3	21.6	87.8	26.1	13.8	31.0
	1929	157.4	18.9	94.1	19.5	10.8	18.9	40.1	18.3	6.4	18.2
	1928	144.9	19.5	106.0	24.4	11.5	15.9	25.6	12.0	1.8	8.0
1 and $1\frac{1}{2}$	1932	169.5	23.5	95.0	24.8	14.8	21.9	54.6	22.5	4.5	17.5
	1931	169.8	16.8	105.6	18.9	15.4	16.4	45.9	14.5	2.9	7.3
	1930	140.1	13.8	71.8	13.3	20.0	21.3	44.0	13.1	4.3	9.6
	1929	81.0	9.7	33.7	8.0	9.8	15.3	30.0	11.9	2.6	7.1
	1928	68.5	9.2	37.1	8.5	9.3	12.8	21.2	9.9	.9	4.0
$\frac{1}{2}$ and $1\frac{1}{2}$	1932	81.8	11.3	29.7	7.7	12.0	19.1	38.3	15.8	.9	3.5
	1931	67.8	6.7	20.0	3.6	15.8	16.8	31.7	10.0	.3	.8
	1930	83.9	8.3	28.2	5.2	16.8	17.9	37.6	11.2	1.3	2.9
	1929	42.6	5.1	14.4	3.0	8.0	13.9	18.5	7.4	.8	2.2
	1928	33.9	4.6	10.0	2.3	5.1	7.0	18.6	8.7	.2	.9
$\frac{1}{2}$ and $1\frac{1}{2}$	1932	44.0	6.1	7.6	2.0	9.3	13.7	26.8	11.0	.3	1.2
	1931	36.0	3.6	2.9	.5	14.5	15.4	18.4	5.8	.2	.5
	1930	38.4	3.8	5.5	1.0	10.5	11.2	22.0	6.5	.4	.9
	1929	14.4	1.7	2.9	.6	5.1	8.0	6.3	2.5	.1	.3
	1928	23.4	3.1	4.8	1.1	4.0	5.5	14.5	6.8	.1	.4
$1\frac{1}{2}$ and $1\frac{1}{2}$	1932	15.8	2.2	1.3	.3	4.8	7.1	9.7	4.0		
	1931	12.0	1.2	.7	.1	6.7	7.1	4.6	1.5		
	1930	10.6	1.0	.8	.1	3.8	4.0	5.9	1.8	.1	.2
	1929	3.7	.4	.5	.1	1.3	2.0	1.9	.8		
	1928	9.6	1.3	1.8	.4	1.6	2.2	6.1	2.8	.1	.4
1 $\frac{1}{4}$ and longer.....	1932	1.5	.2	.1	(3)	.4	.6	1.0	.4		
	1931	1.8	.2			1.7	1.8	.1	(3)		
	1930	.8	.1			.6	.6	.2			
	1929	.5	.1	.1	(3)	.1	.2	.3			
	1928	2.4	.3	.5	.1	.4	.6	1.5	.7		

¹ As reported by the Bureau of the Census.

² Untenderable in settlement of futures contracts made subject to sec. 5 of the United States Cotton Futures Act and the regulations of the Secretary of Agriculture thereunder.

³ Less than 0.05 percent.




FIGURE 9.

TABLE 17.—*Staple length of American upland cotton ginned in designated districts of Tennessee, crops of 1932, 1931, 1930, 1929, and 1928 (fig. 10)*

[Quantities are given in running bales, except that round bales are counted as half bales. Linters are not included]

Staple length (inches)	Crop year	Total		District 1		District 2		District 3		District 4	
		1,000 bales	Per- cent								
All lengths -----	1932	467.5	100.0	193.1	100.0	207.4	100.0	18.2	100.0	48.8	100.0
	1931	578.0	100.0	246.2	100.0	250.2	100.0	30.2	100.0	51.4	100.0
	1930	371.4	100.0	136.2	100.0	157.2	100.0	29.5	100.0	48.5	100.0
	1929	504.3	100.0	184.6	100.0	225.5	100.0	33.5	100.0	60.7	100.0
	1928	423.5	100.0	150.9	100.0	184.3	100.0	27.8	100.0	60.5	100.0
Shorter than $\frac{7}{8}$" -----	1932	12.4	2.7	2.3	1.2	7.5	3.6	2.0	11.0	.6	1.2
	1931	39.9	6.9	3.9	1.6	17.7	7.1	7.6	25.2	10.7	20.8
	1930	59.7	16.1	9.2	6.8	26.6	16.9	12.1	41.0	11.8	24.3
	1929	80.0	15.9	4.5	2.4	44.2	19.6	16.4	48.9	14.9	24.6
	1928	88.0	20.8	10.5	7.0	50.5	27.4	14.8	53.3	12.2	20.2
$\frac{7}{8}$ and $2\frac{1}{2}$" -----	1932	115.9	24.8	31.3	16.2	53.2	25.6	13.6	74.7	17.8	36.5
	1931	218.6	37.8	74.5	30.2	99.1	39.6	17.2	56.9	27.8	54.1
	1930	128.1	34.5	40.5	29.7	57.4	36.6	11.7	39.7	18.5	38.2
	1929	173.4	34.4	50.3	27.2	85.6	38.0	14.5	43.3	23.0	37.9
	1928	173.0	40.9	56.6	37.5	77.3	41.9	10.6	38.1	28.5	47.1
$2\frac{1}{2}$ and $3\frac{1}{2}$" -----	1932	175.3	37.5	71.6	37.1	80.7	38.0	2.3	12.6	20.7	42.4
	1931	211.3	38.6	100.6	40.8	96.2	38.5	4.9	16.2	9.6	18.7
	1930	123.8	33.3	50.7	37.2	54.0	34.3	4.9	16.6	14.2	29.3
	1929	163.8	32.5	77.8	42.1	87.9	30.1	2.3	6.9	15.8	26.0
	1928	108.0	25.5	48.2	31.9	42.7	23.1	2.2	7.9	14.9	24.6
$1\frac{1}{2}$ and $1\frac{3}{4}$" -----	1932	134.7	28.8	73.0	37.8	52.8	25.5	-2	1.1	8.7	17.8
	1931	93.5	16.2	58.1	23.6	31.7	12.7	.5	1.7	3.2	6.2
	1930	48.0	12.9	27.2	20.0	16.5	10.5	.8	2.7	3.5	7.2
	1929	74.9	14.8	42.4	23.0	25.7	11.4	.3	.9	6.5	10.7
	1928	42.1	9.9	26.1	17.3	11.4	6.2	.2	.7	4.4	7.3
$1\frac{1}{2}$ and $1\frac{3}{4}$" -----	1932	24.6	5.3	12.8	6.6	10.7	5.2	.1	.6	1.0	2.1
	1931	11.7	2.0	7.3	3.0	4.3	1.7			.1	.2
	1930	11.0	3.0	7.9	5.8	2.6	1.7			.5	1.0
	1929	10.3	2.0	7.9	4.3	1.9	.8			.5	.8
	1928	10.4	2.5	7.9	5.2	2.0	1.1			.5	
$1\frac{1}{2}$ and $1\frac{3}{4}$" -----	1932	4.3	.9	1.9	1.0	2.4	1.2				
	1931	2.5	.4	1.4	.6	1.1	.4				
	1930	.8	.2	.7	.5	.1	.1				
	1929	1.6	.3	1.4	.8	.2	.1				
	1928	1.4	.3	1.2	.8	.2	.1				
$1\frac{3}{4}$ and $1\frac{7}{8}$" -----	1932	.3	(3)	.2	.1	.1	(4)				
	1931	.5	.1	.4	.2	.1	(3)				
	1930										
	1929	.3	.1	.3	.2						
	1928	.5	.1	.4	.3	.1	.1				
$1\frac{1}{4}$ and longer -----	1931										
	1930										
	1929										
	1928	.1	(3)			.1	.1				

¹ As reported by the Bureau of the Census.

² Untenderable in settlement of futures contracts made subject to sec. 5 of the United States Cotton Futures Act and the regulations of the Secretary of Agriculture thereunder.

³ Less than 0.05 percent.




FIGURE 10.

TABLE 18.—*Staple length of American upland cotton ginned in designated districts of Texas, crops of 1932, 1931, 1930, 1929, and 1928 (fig. 11)*

[Quantities are given in running bales, except that round bales are counted as half bales. Linters are not included]

Staple length (inches)	Crop year	Total		District 1		District 2		District 3		District 4		District 5	
		1,000 bales	Percent										
All lengths	1932	1,4307.4	100.0	35.9	100.0	501.5	100.0	1,122.0	100.0	142.5	100.0	120.3	100.0
	1931	1,5068.8	100.0	55.0	100.0	558.2	100.0	780.8	100.0	146.2	100.0	136.5	100.0
	1930	1,8886.1	100.0	58.7	100.0	287.8	100.0	415.0	100.0	96.4	100.0	79.1	100.0
	1929	1,3803.2	100.0	54.4	100.0	359.5	100.0	605.3	100.0	120.0	100.0	99.0	100.0
	1928	1,4941.5	100.0	66.7	100.0	324.9	100.0	752.6	100.0	175.7	100.0	130.8	100.0
	1932	414.3	9.6			94.6	16.8	133.8	11.9	2.0	1.4	7.5	6.2
Shorter than 7/8	1931	446.5	8.8	1.2	2.2	154.3	27.7	149.5	19.1	6.9	4.7	14.7	10.8
	1930	475.5	12.2	.6	1.0	88.0	30.6	122.0	29.3	25.1	26.0	18.7	23.6
	1929	974.2	25.6	1.6	2.9	245.8	68.4	304.3	50.3	35.7	29.7	49.2	49.7
	1928	794.6	16.1	2.6	3.0	107.6	33.1	202.4	26.9	23.7	13.5	21.1	15.1
	1932	1,918.6	44.5			220.5	39.3	448.6	40.1	61.8	43.4	67.6	56.6
	1931	2,123.4	41.9	.9	1.6	336.6	60.3	459.9	58.9	91.5	62.0	84.5	61.9
7/8 and 2 1/2	1930	1,330.4	34.2	.2	.3	140.4	48.8	200.2	49.6	54.3	56.3	35.7	45.1
	1929	1,280.5	33.2	1.7	3.1	97.8	27.2	237.2	39.2	60.2	50.2	25.2	35.6
	1928	1,856.8	37.0	1.8	2.7	143.3	44.1	287.5	38.2	68.5	39.0	56.4	40.4
	1932	1,554.8	36.1	.9	2.5	193.5	34.5	370.4	33.0	87.4	47.3	40.3	33.5
	1931	1,719.3	33.9	5.6	10.2	61.5	11.0	156.6	20.1	43.2	29.6	35.2	25.8
	1930	1,303.2	35.1	3.6	6.1	47.5	16.5	78.8	18.9	14.8	15.4	21.6	27.3
1 1/2 and 2 1/2	1929	955.9	25.1	3.6	6.6	14.1	3.9	55.4	9.2	21.7	18.1	12.9	13.0
	1928	1,557.1	31.5	4.9	7.3	63.6	19.6	192.9	25.6	62.4	35.5	49.4	35.3
	1932	362.3	8.4	9.4	26.2	50.2	8.9	157.3	14.0	11.2	7.8	4.9	4.1
	1931	678.3	13.4	12.5	22.7	5.3	.9	14.2	1.8	4.6	3.1	2.1	1.5
	1930	586.3	15.1	10.7	18.3	11.3	3.0	8.6	2.1	2.1	2.2	2.0	3.7
	1929	480.8	12.6	23.1	42.5	1.1	.3	5.6	9.4	2.4	2.0	1.7	1.7
1 1/2 and 1 3/2	1928	588.8	11.9	16.7	25.1	0.7	3.0	50.5	7.9	19.9	11.3	12.5	8.9
	1932	37.5	.9	12.1	33.7	2.6	.5	10.8	1.0	.1	.1		
	1931	86.9	1.7	26.8	48.7	.4	.1	.5	.1				
	1930	101.0	2.6	38.1	64.9	.5	.2	.2	.1	.1	.1	.2	.3
	1929	103.2	2.7	18.5	34.0	.3	.1	1.6	.3				
	1928	113.0	2.3	33.1	49.6	.6	.2	4.9	.7	1.1	.6	.4	.3
1 3/4 and 2 1/2	1932	19.7	.5	13.5	37.6	.1	(3)	.3	(3)				
	1931	13.9	.3	8.0	14.6	.1	(3)	.1	(3)				
	1929	25.7	.7	5.5	9.4	.4		.1	(3)				
	1928	26.0	.7	5.9	10.9	.4		.9	1				
	1932	26.9	.5	7.5	11.2	.1	(3)	3.7	.5	.1	.1		
	1931	.2	(3)										
1 3/4 and 2 1/2	1930	.4	(3)										
	1929	3.9	.1			.1	(3)						
	1928	2.4	.1					.3	(3)				
	1932	5.8	.1	.1	.2			1.6	.2				
	1931	.1	(3)										
	1930	.1	(3)										
1 3/4 and longer	1929	.2	(3)										
	1928	.5	(3)					.1	(3)				

Staple length (inches)	Crop year	District 6		District 7		District 8		District 9		District 10		District 11	
		1,000 bales	Percent										
All lengths	1932	123.8	100.0	1,634.3	100.0	504.4	100.0	163.0	100.0	174.2	100.0	324.9	100.0
	1931	141.9	100.0	1,548.9	100.0	711.6	100.0	227.8	100.0	216.8	100.0	545.1	100.0
	1930	103.1	100.0	1,118.4	100.0	553.3	100.0	212.9	100.0	350.4	100.0	610.1	100.0
	1929	116.3	100.0	1,064.2	100.0	684.8	100.0	120.9	100.0	153.7	100.0	425.1	100.0
	1928	143.4	100.0	1,386.6	100.0	898.1	100.0	255.9	100.0	244.8	100.0	553.0	100.0
	1932	3.9	3.1	18.8	1.8	130.3	25.8	7.2	4.4	4.5	2.6	11.9	3.7
Shorter than 7/8	1931	6.6	4.7	30.2	1.9	63.7	9.0	6.8	3.0	2.6	1.2	10.0	1.8
	1930	18.1	17.6	37.1	3.3	128.9	22.9	10.8	5.1	10.0	2.9	18.2	3.0
	1929	28.8	24.8	63.8	6.0	230.1	33.6	6.6	5.5	2.5	1.6	5.8	1.4
	1928	27.2	19.0	112.4	8.1	203.5	22.7	14.1	5.5	34.2	14.0	45.8	8.3
	1932	67.7	54.7	482.8	40.7	283.4	50.2	85.0	51.9	82.6	47.4	117.0	35.2
	1931	75.9	53.5	513.2	33.1	328.4	46.2	75.2	33.0	46.6	21.5	110.7	20.3
7/8 and 2 1/2	1930	55.4	53.7	324.6	29.0	233.6	42.2	72.4	34.0	96.6	27.6	111.0	18.2
	1929	59.8	51.4	387.5	36.4	271.4	39.6	45.2	37.4	32.1	20.9	32.4	7.6
	1928	65.1	45.4	509.2	36.7	356.7	30.7	90.9	39.0	107.4	43.8	161.0	29.1
	1932	50.0	40.4	480.8	46.5	84.0	16.7	48.7	29.8	77.1	44.2	141.7	43.6
	1931	52.1	36.7	739.8	47.8	240.0	33.7	99.8	43.8	90.6	41.8	194.9	35.7
	1930	24.6	23.9	577.7	51.7	147.7	26.7	91.3	42.9	100.1	45.6	105.5	32.0
1 3/4 and 2 1/2	1929	25.6	22.0	442.8	41.6	140.1	20.5	50.0	41.3	70.5	45.9	119.2	28.0
	1928	40.6	28.3	500.3	40.4	236.6	26.3	93.8	36.6	75.9	31.0	176.8	32.0

¹ As reported by the Bureau of the Census.

² Untenderable in settlement of futures contracts made subject to sec. 5 of the United States Cotton Futures Act and the regulations of the Secretary of Agriculture thereunder.

³ Less than 0.05 percent.




FIGURE 11.

TABLE 18.—*Staple length of American upland cotton ginned in designated districts of Texas, crops of 1932, 1931, 1930, 1929, and 1928 (fig. 11)—Continued*

Staple length (inches)	Crop year	District 6		District 7		District 8		District 9		District 10		District 11	
		1,000 bales	Per- cent										
1 and 1½2	1932	2.2	1.8	51.3	4.9	6.2	1.2	13.6	8.3	9.9	5.7	46.1	14.2
	1931	7.3	5.1	250.9	16.6	71.5	10.0	41.3	18.1	73.0	33.7	189.6	34.8
	1930	5.0	4.8	177.1	16.8	42.8	7.7	37.1	17.4	80.3	22.9	208.4	34.2
	1929	2.1	1.8	160.8	15.1	37.7	5.5	18.4	15.2	45.4	29.5	182.5	42.9
	1928	9.8	6.8	187.3	13.5	84.1	9.4	42.4	16.6	25.5	10.4	119.4	21.6
	1932	—	—	—	—	—	—	—	—	—	—	—	—
1½ and 1¾2	1931	—	—	—	—	—	—	—	—	—	—	—	—
	1930	—	—	—	—	—	—	—	—	—	—	—	—
	1929	—	—	—	—	—	—	—	—	—	—	—	—
	1928	—	—	—	—	—	—	—	—	—	—	—	—
	1932	—	—	—	—	—	—	—	—	—	—	—	—
	1931	—	—	—	—	—	—	—	—	—	—	—	—
1¾ and 1½2	1930	—	—	—	—	—	—	—	—	—	—	—	—
	1929	—	—	—	—	—	—	—	—	—	—	—	—
	1928	—	—	—	—	—	—	—	—	—	—	—	—
	1932	—	—	—	—	—	—	—	—	—	—	—	—
	1931	—	—	—	—	—	—	—	—	—	—	—	—
	1930	—	—	—	—	—	—	—	—	—	—	—	—
1½ and 1¾2	1929	—	—	—	—	—	—	—	—	—	—	—	—
	1928	—	—	—	—	—	—	—	—	—	—	—	—
	1932	—	—	—	—	—	—	—	—	—	—	—	—
	1931	—	—	—	—	—	—	—	—	—	—	—	—
	1930	—	—	—	—	—	—	—	—	—	—	—	—
	1929	—	—	—	—	—	—	—	—	—	—	—	—
1¾ and longer	1928	—	—	—	—	—	—	—	—	—	—	—	—
	1931	—	—	—	—	—	—	—	—	—	—	—	—
	1930	—	—	—	—	—	—	—	—	—	—	—	—
	1929	—	—	—	—	—	—	—	—	—	—	—	—
	1928	—	—	—	—	—	—	—	—	—	—	—	—

³ Less than 0.05 percent.TABLE 19.—*Tenderability of American upland cotton ginned in the United States: Quantities, by States, crops of 1932, 1931, 1930, 1929, and 1928*

[Quantities are given in running bales, except that round bales are counted as half bales. Linters are not included]

State	Crop year	Total Ameri- can upland cotton ginned ¹	Tenderable ²								Untender- able	
			All staple lengths		Staple lengths ½ to 1½2 inches		Staple lengths 1½ to 2 inches and longer		All staple lengths			
			1,000 bales	Per- cent	1,000 bales	Per- cent	1,000 bales	Per- cent	1,000 bales	Per- cent		
United States	1932	12,701.3	100	11,489.1	90.5	9,920.4	78.2	1,559.7	12.3	1,212.2	9.6	
	1931	16,615.2	100	14,833.9	89.3	12,989.4	78.2	1,844.6	11.1	1,781.3	10.7	
	1930	13,732.2	100	11,623.2	84.6	10,207.2	74.3	1,416.0	10.3	2,109.0	15.4	
	1929	14,519.0	100	10,992.5	75.7	9,433.5	65.0	1,559.0	10.7	3,520.5	24.3	
	1928	14,268.2	100	11,664.7	81.8	10,231.2	71.7	1,433.5	10.1	2,003.5	18.2	
	1932	933.7	100	792.0	84.8	788.3	84.4	3.7	.4	141.7	15.2	
Alabama	1931	1,385.0	100	1,247.0	90.1	1,240.5	89.6	7.1	.5	137.4	9.9	
	1930	1,444.9	100	873.0	60.4	869.9	60.2	3.1	.2	571.9	39.6	
	1929	1,307.7	100	705.0	54.0	704.4	53.9	1.5	.1	601.8	40.0	
	1928	1,096.6	100	830.4	75.7	828.0	76.5	2.4	.2	206.2	24.3	
	1932	58.8	100	57.9	98.5	31.0	52.7	26.9	45.8	0.9	1.5	
	1931	97.3	100	92.9	95.5	57.8	59.4	35.1	36.1	4.4	4.5	
Arizona	1930	127.2	100	119.8	94.2	87.4	68.7	32.4	25.5	7.4	5.8	
	1929	120.7	100	114.8	95.1	92.6	76.7	22.2	18.4	5.9	4.9	
	1928	117.4	100	112.6	95.8	66.9	57.0	45.6	38.8	4.9	4.2	
	1932	1,283.4	100	1,187.5	92.5	864.9	67.4	322.6	25.1	95.9	7.5	
	1931	1,836.1	100	1,616.0	88.0	1,298.4	70.7	318.2	17.3	219.5	12.0	
	1930	863.6	100	773.5	89.6	643.8	74.6	129.7	15.0	90.0	10.4	
Arkansas	1929	1,395.9	100	1,150.3	82.4	978.3	70.1	172.0	12.3	245.6	17.6	
	1928	1,216.2	100	1,008.3	82.9	807.8	66.4	200.5	16.5	207.9	17.1	
	1932	124.4	100	120.9	97.2	16.2	13.0	104.7	84.2	3.5	15.4	
	1931	171.2	100	161.4	94.3	27.8	16.2	133.0	78.1	9.8	5.7	
	1930	256.3	100	245.6	95.8	46.1	18.0	109.5	77.8	10.7	4.2	
	1929	254.1	100	246.4	97.0	135.1	53.2	111.3	43.8	7.7	3.0	
California	1928	171.0	100	161.3	94.3	92.0	53.8	69.3	40.5	9.7	5.7	
	1932	15.6	100	13.2	84.0	13.2	84.6	—	—	2.4	15.4	
	1931	43.4	100	40.2	92.6	40.1	92.4	.1	.2	3.2	7.4	
	1930	51.1	100	40.7	79.6	40.7	79.6	—	—	10.4	20.4	
	1929	29.8	100	18.6	62.4	18.5	62.1	.1	.3	11.2	37.6	
	1928	20.1	100	17.0	84.6	10.7	83.1	.3	1.5	3.1	15.4	
Florida	1932	861.8	100	809.3	93.9	800.4	92.9	8.9	1.0	52.6	6.1	
	1931	1,393.7	100	1,222.0	94.9	1,310.2	94.0	11.8	.9	71.7	5.1	
	1930	1,597.5	100	1,317.8	82.5	1,309.0	81.9	8.8	.6	279.7	17.5	
	1929	1,239.8	100	996.7	74.4	992.9	74.1	2.8	.3	343.1	25.6	
	1928	1,053.2	100	908.9	86.3	901.5	86.6	7.4	.7	144.3	13.7	

See footnotes at end of table.

TABLE 19.—*Tenderability of American upland cotton ginned in the United States: Quantities, by States, crops of 1932, 1931, 1930, 1929, and 1928—Continued*

State	Crop year	Total Ameri-can upland cotton ginned ¹	Tenderable ²						Untender-able	
			All staple lengths		Staple lengths $\frac{1}{8}$ to $\frac{1}{2}$ inches		Staple lengths $\frac{1}{4}$ inches and longer			
			1,000 bales	Per-cent	1,000 bales	Per-cent	1,000 bales	Per-cent	1,000 bales	Per-cent
Louisiana.....	1932	599.5 100	535.8	89.4	438.8	73.2	97.0	16.2	63.7	10.6
	1931	876.6 100	795.4	90.7	623.2	71.1	172.2	19.6	81.2	9.3
	1930	704.8 100	598.2	84.9	500.6	71.0	97.6	13.9	106.6	15.1
	1929	797.7 100	674.4	84.5	577.5	72.4	98.9	12.1	123.3	15.5
	1928	685.9 100	592.2	86.3	505.0	73.6	87.2	12.7	93.7	13.7
Mississippi.....	1932	1,161.2 100	1,114.0	95.9	473.8	40.8	640.2	55.1	47.2	4.1
	1931	1,719.4 100	1,561.8	90.8	728.2	42.2	835.6	48.6	157.6	9.2
	1930	1,458.5 100	1,322.9	90.7	759.3	52.1	563.6	38.6	135.6	9.3
	1929	1,876.0 100	1,625.1	86.6	762.2	40.6	862.9	46.0	250.9	13.4
	1928	1,462.0 100	1,176.8	80.5	490.1	33.5	686.7	47.0	285.2	10.5
Missouri.....	1932	300.7 100	243.1	80.8	219.7	73.0	23.4	7.8	57.6	19.2
	1931	280.4 100	204.5	72.9	194.5	69.3	10.0	3.6	75.9	27.1
	1930	153.3 100	133.9	87.3	126.3	82.3	7.6	5.0	19.4	12.7
	1929	220.9 100	152.5	69.0	140.3	63.5	12.2	5.5	68.4	31.0
	1928	146.9 100	94.0	64.0	79.6	54.2	14.4	9.8	52.9	36.0
Missouri.....	1932	67.5 100	60.1	89.0	24.3	36.0	35.8	53.0	7.4	11.0
	1931	93.8 100	88.1	93.9	51.2	54.6	36.9	39.3	5.7	6.1
	1930	95.8 100	90.0	93.9	39.4	41.1	50.6	52.8	5.8	6.1
	1929	86.3 100	75.4	87.4	43.2	50.1	32.2	37.3	10.9	12.6
	1928	82.2 100	76.1	92.6	35.7	43.4	40.4	49.2	6.1	7.4
New Mexico.....	1932	680.3 100	668.3	98.2	607.8	80.3	60.5	8.9	12.0	1.8
	1931	771.2 100	761.7	98.8	729.0	94.5	32.7	4.3	9.5	1.2
	1930	800.6 100	747.0	93.3	715.3	89.3	31.7	4.0	53.6	6.7
	1929	767.0 100	673.1	87.8	664.3	86.6	8.8	1.2	93.9	12.2
	1928	869.3 100	771.5	88.8	755.1	86.9	16.4	1.9	97.8	11.2
North Carolina.....	1932	1,072.0 100	959.6	89.5	953.8	89.0	5.8	.5	112.4	10.5
	1931	1,235.9 100	994.1	78.0	948.9	76.8	15.2	1.2	271.8	22.0
	1930	856.8 100	700.2	81.7	685.5	80.0	14.7	1.7	156.6	18.3
	1929	1,125.6 100	708.3	62.9	679.5	60.4	28.8	2.5	417.3	37.1
	1928	1,187.0 100	900.2	75.8	865.7	72.9	34.5	2.9	286.8	24.2
Oklahoma.....	1932	722.2 100	714.8	99.0	572.4	70.3	142.4	19.7	7.4	1.0
	1931	1,010.2 100	997.1	98.7	879.5	87.1	117.6	11.6	13.1	1.3
	1930	1,015.3 100	975.2	96.1	842.4	83.0	132.8	13.1	40.1	3.9
	1929	833.1 100	711.4	85.4	650.5	78.1	60.9	7.3	121.7	14.6
	1928	744.4 100	667.2	89.6	598.4	80.4	68.8	9.2	77.2	10.4
South Carolina.....	1932	467.5 100	423.4	90.6	395.1	84.5	28.3	6.1	44.1	9.4
	1931	578.0 100	508.8	88.0	494.1	85.5	14.7	2.5	69.2	12.0
	1930	371.4 100	301.6	81.2	289.8	78.0	11.8	3.2	69.8	18.8
	1929	504.3 100	381.4	75.6	370.1	73.4	11.3	2.2	122.9	24.4
	1928	423.5 100	304.5	71.9	292.7	69.1	11.8	2.8	119.0	28.1
Tennessee.....	1932	4,307.4 100	3,748.6	87.0	3,691.6	85.7	57.0	1.3	558.8	13.0
	1931	5,068.8 100	4,422.8	87.3	4,321.7	85.3	101.1	2.0	646.0	12.7
	1930	3,886.1 100	3,340.1	85.9	3,209.7	82.5	130.4	3.4	546.0	14.1
	1929	3,803.2 100	2,708.5	71.2	2,578.8	67.8	120.7	3.4	1,094.7	28.8
	1928	4,941.5 100	3,999.1	80.9	3,853.8	78.0	145.3	2.9	942.4	19.1
Texas.....	1932	31.3 100	29.7	94.9	29.5	94.3	0.2	0.6	1.6	5.1
	1931	42.5 100	40.8	96.0	40.8	90.0	-----	-----	1.7	4.0
	1930	42.7 100	38.5	90.2	38.4	90.0	.1	.2	4.2	9.8
	1929	48.0 100	44.4	92.5	44.3	92.3	.1	.2	3.6	7.5
	1928	44.8 100	41.4	92.4	41.4	92.4	-----	-----	3.4	7.6
Virginia.....	1932	14.0 100	10.9	77.9	8.6	61.5	2.3	16.4	3.1	22.1
	1931	11.7 100	8.1	69.2	5.5	47.0	2.6	22.2	3.6	30.8
	1930	6.4 100	5.2	81.3	3.6	56.3	1.6	25.0	1.2	18.7
	1929	8.9 100	5.3	59.0	1.0	11.2	4.3	48.4	3.6	40.4
	1928	6.2 100	3.3	53.2	.8	12.9	2.5	40.3	2.9	46.8
All other ³										

¹ As reported by the Bureau of the Census.² Tenderable in settlement of futures contracts made subject to sec. 5 of the United States Cotton Futures Act and the regulations of the Secretary of Agriculture thereunder.³ Includes Illinois, Kansas, and Kentucky.

TABLE 20.—*Tenderability of American upland cotton on hand in the United States, Aug. 1, 1933, 1932, 1931, 1930, 1929, and 1928*

[Quantities are given in running bales, except that round bales are counted as half bales. Linters are not included]

Year	Total Ameri-can upland cotton on hand Aug. 1 ¹	Tenderable ²								Untenderable	
		All staple lengths		Staple lengths $\frac{3}{4}$ to $1\frac{1}{2}$ inches		Staple lengths $1\frac{1}{4}$ inches and longer					
		1,000 bales	Percent	1,000 bales	Percent	1,000 bales	Percent	1,000 bales	Percent		
1933	8,089.7	100	7,437.4	92.2	6,085.8	75.2	1,371.0	17.0	632.3	7.8	
1932	9,500.3	100	8,882.7	92.9	7,418.4	77.6	1,464.3	15.3	677.6	7.1	
1931	6,240.0	100	5,543.3	88.7	4,773.0	70.4	770.3	12.3	702.7	11.3	
1930	4,313.6	100	3,416.3	79.2	2,660.6	61.8	749.7	17.4	807.3	20.8	
1929	2,122.6	100	1,747.0	82.3	1,251.0	58.9	496.0	23.4	375.6	17.7	
1928	2,410.8	100	2,198.0	90.8	1,459.5	60.3	738.5	30.5	221.8	9.2	

¹ As reported by the Bureau of the Census.

² Tenderable in settlement of futures contracts made subject to sec. 5 of the United States Cotton Futures Act and the regulations of the Secretary of Agriculture thereunder.

TABLE 21.—*Grade and staple length of American upland cotton: Quantities on hand in the United States, Aug. 1, 1932*

[Quantities are given in running bales, except that round bales are counted as half bales. Linters are not included]

Grade	All staple lengths		Shorter than $\frac{3}{8}$ inch ¹	$\frac{3}{4}$ and $2\frac{1}{2}$ inch	$1\frac{1}{4}$ and $3\frac{1}{2}$ inch	1 and $1\frac{1}{2}$ inches	$1\frac{1}{4}$ and $1\frac{1}{2}$ inches	$1\frac{1}{2}$ and $1\frac{1}{2}$ inches	$1\frac{1}{4}$ and $1\frac{1}{2}$ inches	$1\frac{1}{4}$ inches and longer
	1,000 bales	Percent								
	1,000 bales	Percent								
All grades	29,560.3	100.0	298.3	3,302.6	2,704.0	1,657.6	754.5	546.7	174.0	32.0
Extra White	104.2	1.1	.4	5.9	10.5	28.3	39.6	18.0	1.5	-----
Above 3-G.M.	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
3-G.M.	33.3	.3	-----	.3	1.2	12.5	13.4	5.5	.4	-----
4-S.M.	40.4	.4	-----	.9	3.3	9.8	16.2	9.5	.9	-----
5-M.	19.0	.2	-----	.1	2.4	3.2	3.5	7.3	.2	-----
6-S.L.M.	8.9	.1	-----	.1	1.7	2.3	1.8	2.3	.7	-----
7-L.M.	2.5	(3)	-----	.1	.6	.5	.9	.4	-----	-----
Below 7-L.M. ¹	.1	(3)	-----	.1	-----	-----	-----	-----	-----	-----
White	8,507.1	89.0	230.4	2,880.2	2,445.5	1,536.0	688.8	522.3	171.5	32.4
1-M.F.	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
2-S.G.M.	3.2	(3)	-----	.2	.9	1.4	.3	.1	.2	.1
3-G.M.	454.7	4.8	24.0	162.0	143.5	50.3	22.7	30.1	17.6	4.5
4-S.M.	3,183.5	33.3	80.7	1,155.9	992.7	484.7	197.9	177.5	75.7	18.4
5-M.	3,292.2	34.4	72.6	1,104.3	875.6	655.8	292.0	218.2	64.8	8.9
6-S.L.M.	1,083.3	11.3	32.6	333.7	275.7	234.7	123.8	70.6	11.6	.6
7-L.M.	243.1	2.5	8.0	62.6	75.0	56.7	20.3	13.1	1.4	-----
8-S.G.O.	148.6	1.6	6.7	32.7	47.5	36.6	15.7	9.1	.3	-----
9-G.O. ¹	98.5	1.0	5.6	28.1	34.1	16.9	10.3	3.5	-----	-----
Spotted	829.4	8.7	30.9	460.9	222.7	84.4	23.7	5.7	1.0	.1
3-G.M.	102.4	1.1	5.4	62.3	23.9	7.2	2.8	.7	.1	-----
4-S.M.	392.3	4.1	11.2	217.8	107.4	40.6	11.8	2.9	.6	.1
5-M.	244.3	2.6	6.6	136.8	64.7	26.7	7.4	1.8	.2	-----
6-S.L.M. ¹	59.0	.6	3.7	28.6	17.8	7.3	1.3	.2	.1	-----
7-L.M. ¹	31.4	.3	4.0	15.4	8.9	2.6	.4	.1	-----	-----
Yellow Tinged	55.5	.6	4.8	31.8	14.0	4.2	.5	.2	-----	-----
2-S.G.M.	.1	(3)	-----	.1	-----	-----	-----	-----	-----	-----
3-G.M.	3.2	(3)	-----	.1	2.2	.5	.4	-----	-----	-----
4-S.M.	16.5	.2	1.0	9.7	4.2	1.3	.2	.1	-----	-----
5-M. ¹	19.5	.2	.7	10.9	5.0	1.7	.2	.1	-----	-----
6-S.L.M. ¹	11.3	.1	1.4	6.5	2.7	.6	.1	-----	-----	-----
7-L.M. ¹	4.9	.1	1.6	2.4	.7	.2	-----	-----	-----	-----

See footnotes at end of table.

TABLE 21.—Grade and staple length of American upland cotton: Quantities on hand in the United States, Aug. 1, 1932—Continued

Grade	All staple lengths		Shorter than $\frac{7}{8}$ inch ¹	$\frac{7}{8}$ and $2\frac{1}{2}$ inch	$1\frac{1}{8}$ and $3\frac{1}{2}$ inch	1 and $1\frac{1}{2}$ inches	$1\frac{1}{8}$ inches and longer			
	1,000 bales	Percent (%)								
Light Yellow Stained.....	0.8		1,000 bales 0.2	1,000 bales 0.4	1,000 bales 0.2	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales
3-G.M.....	.1	(2)		.1						
4-S.M. ¹2	(2)		.1	.1					
5-M. ¹5	(2)	.2	.2	.1					
Yellow Stained.....	1.3	(2)	.4	.8	.1					
3-G.M.....	.4	(2)		.1	.3					
4-S.M. ¹4	(2)	.1	.3						
5-M. ¹9	(2)	.3	.6	.1					
Gray.....	4.6	(2)	.4	1.2	1.3	1.2	0.4	0.1		
3-G.M.....	.4	(2)		.2	.1	.1				
4-S.M.....	2.1	(2)	.1	.4	.6	.6	.3	.1		
5-M. ¹	2.1	(2)	.3	.6	.6	.5	.1			
Blue Stained.....	.2	(2)		.1	.1					
3-G.M. ¹										
4-S.M. ¹										
5-M. ¹2	(2)		.1	.1					
No grade ¹	57.2	0.6	30.8	11.3	9.6	3.5	1.5	.4		0.1

¹ Untenderable in settlement of futures contracts made subject to sec. 5 of the United States Cotton Futures Act and the regulations of the Secretary of Agriculture thereunder.

² As reported by the Bureau of the Census.

³ Less than 0.05 percent.

⁴ Includes bales not otherwise classified above.

TABLE 22.—Grade and staple length of American upland cotton: Quantities on hand in the United States, Aug. 1, 1933

[Quantities are given in running bales, except that round bales are counted as half bales. Linters are not included]

Grade	All staple lengths		Shorter than $\frac{7}{8}$ inch ¹	$\frac{7}{8}$ and $2\frac{1}{2}$ inch	$1\frac{1}{8}$ and $3\frac{1}{2}$ inch	1 and $1\frac{1}{2}$ inches	$1\frac{1}{8}$ inches and longer			
	1,000 bales	Percent (%)								
All grades.....	18,069.7	100.0	1,000 bales 188.4	1,000 bales 2,503.6	1,000 bales 2,199.3	1,000 bales 1,774.6	1,000 bales 671.7	1,000 bales 562.9	1,000 bales 143.6	1,000 bales 26.6
Extra White.....	135.3	1.7	2.2	22.1	28.2	31.2	27.6	23.0	1.0	
Above 3-G.M.....	.1	(2)							.1	
3-G.M.....	21.7	.3		.1	.4	8.3	5.3	7.3	.3	
4-S.M.....	32.1	.4	.2	.8	1.6	5.0	10.8	12.3	.5	
5-M.....	34.2	.4	1.5	0.6	0.7	6.9	7.2	2.2	.1	
6-S.L.M.....	32.2	.4	.4	9.6	9.7	7.8	3.7	.9	.1	
7-L.M.....	14.1	.2	.1	4.9	6.4	1.9	.6	.2		
Below 7-L.M. ¹9	(2)		.1	.4					
White.....	6,630.0	82.2	131.4	1,916.9	1,735.2	1,653.3	597.9	528.3	141.5	25.5
1-M.F.....										
2-S.G.M.....	2.2	(2)	.2	.9	.8	.2				
3-G.M.....	202.1	2.5	12.6	68.4	51.5	38.2	15.5	9.3	4.4	2.2
4-S.M.....	1,931.7	24.0	48.3	601.1	478.0	414.5	141.1	121.3	53.5	13.9
5-M.....	2,801.6	34.7	44.0	781.4	698.4	682.2	204.0	257.9	65.1	8.6
6-S.L.M.....	1,210.1	15.0	17.2	283.0	347.1	300.0	130.0	115.6	16.4	.8
7-L.M.....	255.0	3.2	4.5	59.3	79.3	64.0	28.5	17.7	1.7	
8-S.G.O.I.....	144.7	1.8	2.6	34.1	49.0	40.2	13.2	5.2	.4	
9-G.O.I.....	82.6	1.0	2.0	28.7	31.1	14.0	5.6	1.2		

See footnotes at end of table.

TABLE 22.—*Grade and staple length of American upland cotton: Quantities on hand in the United States, Aug. 1, 1933—Continued*

[Quantities are given in running bales, except that round bales are counted as half bales. Linters are not included]

Grade	All staple lengths		Shorter than $\frac{7}{8}$ inch ¹	$\frac{7}{8}$ and $\frac{21}{32}$ inch	$1\frac{1}{16}$ and $1\frac{1}{32}$ inch	1 and $1\frac{1}{32}$ inches	$1\frac{1}{16}$ and $1\frac{1}{32}$ inches	$1\frac{1}{8}$ and $1\frac{1}{32}$ inches	$1\frac{1}{16}$ and $1\frac{1}{32}$ inches	$1\frac{1}{4}$ inches and longer
Spotted	<i>1,000 bales</i> 1,193.9	<i>Percent</i> 14.8	<i>1,000 bales</i> 33.6	<i>1,000 bales</i> 514.5	<i>1,000 bales</i> 410.1	<i>1,000 bales</i> 180.1	<i>1,000 bales</i> 43.3	<i>1,000 bales</i> 11.1	<i>1,000 bales</i> 1.1	<i>1,000 bales</i> 0.1
3-G.M.	102.6	1.3	4.7	52.5	27.0	14.1	3.4	.8	.1	
4-S.M.	547.3	6.8	13.5	226.1	101.8	87.8	21.5	5.9	.6	
5-M.	385.9	4.8	6.4	144.9	151.4	64.0	15.2	3.7	.3	
6-S.L.M. ¹	101.3	1.2	6.5	50.4	24.4	10.8	2.5	.6	.1	
7-L.M. ¹	56.8	.7	2.5	34.6	15.5	3.4	.7	.1		
Yellow Tinged	45.2	.6	4.6	26.0	10.2	3.9	.5			
2-S.G.M.										
3-G.M.	2.7	(3)	.1	1.7	.5	.4				
4-S.M.	14.2	.2	.9	8.4	3.8	.9	.2			
5-M. ¹	16.0	.2	.8	9.9	3.4	1.7	.2			
6-S.L.M. ¹	8.3	.1	1.1	4.4	2.0	.7	.1			
7-L.M. ¹	4.0	(3)	1.7	1.6	.5	.2				
Light Yellow Stained	.5	(3)	.1	.2	.2					
3-G.M.	.1	(3)			.1					
4-S.M. ¹	.2	(3)		.1	.1					
5-M. ¹	.2	(3)	.1	.1						
Yellow Stained	.2	(3)		.2						
3-G.M.										
4-S.M. ¹	.1	(3)		.1						
5-M. ¹	.1	(3)		.1						
Gray	4.0	(3)	.2	.6	1.3	1.0	.6	.3		
3-G.M.	.2	(3)			.1	.1				
4-S.M.	2.1	(3)	.2	.5	.3	.5	.4	.2		
5-M. ¹	1.7	(3)		.1	.9	.4	.2	.1		
No grade ^{1,4}	60.6	.7	16.3	23.1	14.1	5.1	1.8	.2		

Untenderable in settlement of futures contracts made subject to sec. 5 of the United States Cotton Futures Act and the regulations of the Secretary of Agriculture thereunder.

² As reported by the Bureau of the Census.

³ Less than 0.06 percent.

⁴ Includes bales not otherwise classified above.

TABLE 23.—*Grade and staple length of American-Egyptian cotton: Quantities on hand in the United States, Aug. 1, 1933, 1932, 1931, 1930, 1929, and 1928*

[Quantities are given in running bales]

Grade	Year	All staple lengths		Shorter than $1\frac{1}{2}$ inches	$1\frac{1}{2}$ and $1\frac{1}{32}$ inches	$1\frac{1}{16}$ and $1\frac{1}{32}$ inches	$1\frac{1}{8}$ to $1\frac{1}{32}$ inches	$1\frac{1}{4}$ inches and longer
All grades		<i>1,000 bales</i>	<i>Percent</i>	<i>1,000 bales</i>	<i>1,000 bales</i>	<i>1,000 bales</i>	<i>1,000 bales</i>	<i>1,000 bales</i>
	(1933)	19.8	100.0	0.7	2.9	5.1	1.1	
	1932	16.5	100.0		3.1	12.4	1.0	
	1931	16.7	100.0		2.0	13.6	1.1	
	1930	8.1	100.0		.2	7.1	.8	
	1929	7.2	100.0	.3	.7	5.1	1.1	
	(1928)	5.8	100.0	.1	1.6	2.4	1.7	
1 and $1\frac{1}{2}$	(1933)	1.8	18.4		.1	1.3	.4	
	1932	2.9	17.6		.3	2.5	.1	
	1931	4.7	28.1		.3	4.2	.2	
	1930	2.4	20.7			2.1	.3	
	1929	1.7	23.6	.2		1.0	.5	
	(1928)	1.3	22.4		.2	.8	.3	

¹ As reported by the Bureau of the Census.

TABLE 23.—*Grade and staple length of American-Egyptian cotton: Quantities on hand in the United States, Aug. 1, 1933, 1932, 1931, 1930, 1929, and 1928—Continued*

Grade	Year	All staple lengths		Shorter than 1½ inches	1½ and 1⅓ inches	1⅔ and 1⅘ inches	1⅕ to 1⅖ inches	1⅗ to 1⅘ inches	1⅘ inches and longer
		1,000 bales	Per- cent	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales
2 and 2½.	1933	5.5	56.1	0.2	1.7	3.0	0.6	-----	-----
	1932	9.1	55.1	-----	1.5	7.0	.6	-----	-----
	1931	8.6	51.5	-----	.6	7.2	.8	-----	-----
	1930	3.5	43.2	-----	.1	3.0	.4	-----	-----
	1929	4.0	55.5	.1	.1	3.6	.2	-----	-----
	1928	1.1	19.0	-----	.6	.1	.4	-----	-----
	1933	2.0	20.4	.4	1.0	.5	.1	-----	-----
	1932	3.5	21.2	-----	1.1	2.2	.2	-----	-----
3 and 3½.	1931	2.9	17.4	-----	.9	1.9	.1	-----	-----
	1930	2.1	25.0	-----	.1	1.9	.1	-----	-----
	1929	1.2	16.7	-----	.3	.5	.4	-----	-----
	1928	2.7	40.6	-----	.4	1.5	.8	-----	-----
	1933	.5	5.1	.1	.1	.3	-----	-----	-----
	1932	1.0	6.1	-----	.2	.7	.1	-----	-----
	1931	.5	3.0	-----	.2	.3	-----	-----	-----
4 and 4½.	1930	.1	1.2	-----	-----	.1	-----	-----	-----
	1929	.2	2.8	-----	.2	-----	-----	-----	-----
	1928	.4	6.0	.1	.2	-----	.1	-----	-----
5.	1933	-----	-----	-----	-----	-----	-----	-----	-----
	1932	-----	-----	-----	-----	-----	-----	-----	-----
	1931	-----	-----	-----	-----	-----	-----	-----	-----
Below 5.	1930	-----	-----	-----	-----	-----	-----	-----	-----
	1929	-----	-----	-----	-----	-----	-----	-----	-----
	1928	.1	1.7	-----	-----	-----	.1	-----	-----
	1933	-----	-----	-----	-----	-----	-----	-----	-----
	1932	-----	-----	-----	-----	-----	-----	-----	-----
	1931	-----	-----	-----	-----	-----	-----	-----	-----
	1930	-----	-----	-----	-----	-----	-----	-----	-----
	1929	.1	1.4	-----	.1	-----	-----	-----	-----
	1928	.2	3.4	-----	.2	-----	-----	-----	-----

TABLE 24.—*Staple length of American upland cotton: Quantities on hand in the United States, Aug. 1, 1933, 1932, 1931, 1930, 1929, and 1928*

(Quantities are given in running bales, except that round bales are counted as half bales. Linters are not included]

Year	All staple lengths ¹		Shorter than 5/8 inch ²		5/8 and 2 1/32 inch		1 1/8 and 3 1/32 inch		1 and 1 1/2 inches	
	1,000 bales	Per- cent	1,000 bales	Per- cent	1,000 bales	Per- cent	1,000 bales	Per- cent	1,000 bales	Per- cent
1933	8,069.7	100	188.4	2.3	2,503.6	31.0	2,190.3	27.3	1,774.6	22.0
1932	9,580.3	100	298.3	3.1	3,392.6	35.5	2,704.0	28.3	1,657.6	17.4
1931	6,246.0	100	463.2	7.4	2,615.7	41.9	1,528.2	24.5	849.2	13.6
1930	4,313.6	100	446.8	10.4	1,445.6	33.5	825.4	19.1	783.0	18.1
1929	2,122.6	100	155.0	7.3	650.9	30.8	397.4	18.7	395.1	18.6
1928	2,419.8	100	87.8	3.6	485.0	20.0	426.8	17.6	650.8	27.3
Year		1 1/8 and 3 1/32 inches		1 1/8 and 3 1/32 inches		1 1/8 and 3 1/32 inches		1 1/4 inches and longer		
		1,000 bales	Per- cent	1,000 bales	Per- cent	1,000 bales	Per- cent	1,000 bales	Per- cent	
1933	671.7	8.3	562.9	7.0	143.6	1.8	25.6	0.3		
1932	754.5	7.9	546.7	5.7	174.0	1.8	32.8	.3		
1931	414.8	6.6	269.5	4.3	89.7	1.4	15.7	.3		
1930	389.3	9.0	283.4	6.6	115.8	2.7	24.3	.6		
1929	221.3	10.4	170.1	8.0	98.4	4.5	36.4	1.7		
1928	268.5	11.1	287.2	11.9	157.0	6.5	47.7	2.0		

¹ As reported by the Bureau of the Census.² Untenderable in settlement of futures contracts made subject to sec. 5 of the United States Cotton Futures Act and the regulations of the Secretary of Agriculture thereunder.

TABLE 25.—*Staple length of Egyptian cotton: ¹ Quantities on hand in the United States, Aug. 1, 1933, 1932, 1931, 1930, 1929, and 1928*

[Quantities are given in equivalent 500-pound bales]

Year	All staple lengths ²	Shorter than $1\frac{1}{4}$ inches	$1\frac{1}{4}$ and $1\frac{1}{2}\frac{1}{2}$ inches	$1\frac{1}{2}\frac{1}{2}$ and $1\frac{1}{2}\frac{1}{2}$ inches	$1\frac{1}{4}$ to $1\frac{1}{2}\frac{1}{2}$ inches	$1\frac{1}{2}\frac{1}{2}$ to $1\frac{1}{2}\frac{1}{2}$ inches	$1\frac{1}{2}$ to $1\frac{1}{2}\frac{1}{2}$ inches	$1\frac{1}{2}\frac{1}{2}$ to $1\frac{1}{2}\frac{1}{2}$ inches	$1\frac{1}{2}$ inches and longer
	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales
1933	54.8	7.9	12.5	2.0	2.4	22.7	6.6	0.6	0.1
1932	68.1	8.0	13.0	.7	3.1	30.3	11.5	1.4	.1
1931	63.9	1.1	21.5	.8	6.5	28.6	5.2	—	.2
1930	145.4	.3	47.3	11.5	9.9	53.5	22.3	.8	—
1929	129.2	.3	70.8	6.6	2.7	34.1	12.9	1.8	—
1928	65.3	1.4	23.4	1.7	7.5	21.3	7.0	2.6	.4

¹ Foreign cottons are not deliverable in settlement of futures contracts made subject to the United States Cotton Futures Act.² As reported by the Bureau of the Census.TABLE 26.—*Staple length of foreign cotton ¹ other than Egyptian: Quantities on hand in the United States, Aug. 1, 1933, 1932, 1931, 1930, 1929, and 1928*

[Quantities are given in equivalent 500-pound bales]

Growth	Year	All staple lengths ²	Shorter than $\frac{1}{2}$ inch	$\frac{3}{8}$ and $2\frac{1}{2}\frac{1}{2}$ inch	$1\frac{1}{16}$ and $3\frac{1}{2}\frac{1}{2}$ inch	1 and $1\frac{1}{2}\frac{1}{2}$ inches	$1\frac{1}{8}$ and $1\frac{1}{2}\frac{1}{2}$ inches	$1\frac{1}{8}$ and $1\frac{1}{2}\frac{1}{2}$ inches	$1\frac{1}{16}$ and $1\frac{1}{2}\frac{1}{2}$ inches	$1\frac{1}{4}$ inches and longer
		1,000 bales	Percent	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales	1,000 bales
Foreign, other than Egyptian	1933	30.3	100.0	26.7	—	—	0.2	0.4	0.4	2.6
	1932	32.9	100.0	25.4	0.8	—	—	.1	.2	6.4
	1931	43.4	100.0	39.5	—	—	.1	.3	1.3	2.2
	1930	63.3	100.0	55.6	—	—	.3	1.3	.1	0.0
	1929	53.0	100.0	42.1	.3	5.5	.7	.1	.4	3.9
	1928	45.6	100.0	39.3	—	0.3	.1	.1	.9	5.0
Peruvian	1933	1.6	5.3	—	—	—	.1	.3	.4	.8
	1932	2.6	7.9	—	—	—	—	.1	.2	2.3
	1931	3.2	7.4	—	—	—	.1	.3	1.3	1.5
	1930	7.1	11.2	—	—	—	.3	1.3	.1	5.4
	1929	4.6	8.7	—	—	—	.7	.1	.3	3.5
	1928	5.5	12.1	—	—	—	—	.1	.9	4.6
Chinese	1933	21.4	70.6	21.4	—	—	—	—	—	—
	1932	10.5	31.9	10.2	.3	—	—	—	—	—
	1931	21.1	48.6	21.1	—	—	—	—	—	—
	1930	25.5	40.3	25.5	—	—	—	—	—	—
	1929	22.7	42.8	18.1	—	4.6	—	—	—	—
	1928	29.9	65.5	20.9	—	—	—	—	—	—
British Indian	1933	5.3	17.5	5.3	—	—	—	—	—	—
	1932	15.4	46.8	15.0	.4	—	—	—	—	—
	1931	18.3	42.2	18.3	—	—	—	—	—	—
	1930	20.9	47.2	29.9	—	—	—	—	—	—
	1929	25.0	47.2	23.9	.3	.8	—	—	—	—
	1928	9.4	20.6	9.4	—	—	—	—	—	—
Other	1933	2.0	6.6	—	—	—	.1	.1	—	1.8
	1932	4.4	13.4	.2	.1	—	—	—	—	4.1
	1931	.8	1.8	.1	—	—	—	—	—	.7
	1930	.8	1.3	.2	—	—	—	—	—	.6
	1929	.7	1.3	.1	—	.1	—	—	.1	.4
	1928	.8	1.8	—	—	—	.3	—	—	.6

¹ Foreign cottons are not deliverable in settlement of futures contracts made subject to the United States Cotton Futures Act.² As reported by the Bureau of the Census.

STATISTICAL BULLETIN SERIES

At the present time (November 1934) statistical bulletins have been issued as follows:

1. Cold-storage holdings, year ended January 1923.
2. Seed statistics.
3. Sheep, lamb, mutton, and wool statistics.
4. Cold-storage holdings, to October 1924.
5. Horses, mules, and motor vehicles.
6. Grain futures; daily data.
7. Shipments and unloads of certain fruits and vegetables.
8. Carload shipments of fruits and melons.
9. Carload shipments of vegetables.
10. Statistics of potatoes and sweetpotatoes.
11. Hay and feed statistics.
12. Wheat and rye statistics.
13. Cold-storage holdings, year ended December 1925.
14. Prices of farm products received by producers, North Atlantic States.
15. Prices of farm products received by producers, North Central States.
16. Prices of farm products received by producers, South Atlantic and South Central States.
17. Prices of farm products received by producers, Mountain and Pacific States.
18. Statistics of hogs, pork, and pork products.
19. Carload shipments of fruits and vegetables from stations in the United States.
20. Cattle, calves, beef, veal, hides, and skins statistics.
21. American forests and forest products.
22. Vegetable statistics.
23. Carload shipments and unloads of certain fruits and vegetables, 1924-26.
24. Statistics of fats, oils, and oleaginous raw materials.
25. Dairy statistics.
26. Cold-storage holdings, year ended December 31, 1927.
27. Car-lot shipments of fruits and vegetables from stations in the United States for the calendar years 1926 and 1927.
28. Corn statistics.
29. Statistics of oats, barley, and grain sorghums.
30. Car-lot shipments and unloads of important fruits and vegetables for the calendar years 1927 and 1928.
31. Wheat futures: Volume of trading, open commitments, and prices.
32. Stumpage and log prices for the calendar year 1928.
33. Cold-storage holdings, year ended December 31, 1929.
34. Corn futures.
35. Car-lot shipments of fruits and vegetables from stations in the United States for the calendar years 1928 and 1929.
36. Stumpage and log prices for the calendar year 1929.
37. Stumpage and log prices for the calendar year 1930.
38. Car-lot shipments and unloads of important fruits and vegetables for the calendar years 1929 and 1930.
39. Cold-storage holdings, year ended December 31, 1931.
40. Grade, staple length, and tenderability of cotton in the United States, 1928-29 to 1931-32.
41. Wheat futures.
42. Car-lot shipments of fruits and vegetables from stations in the United States for the calendar years 1930 and 1931.
43. Corn futures.
44. Stumpage and log prices for the calendar years 1931 and 1932.
45. Grade and staple length of cotton carried over in the United States as related to the domestic supply, 1928-29 to 1931-32.
46. Car-lot shipments and unloads of important fruits and vegetables for the calendar years 1931 and 1932.
47. Grade, staple length, and tenderability of cotton in the United States, 1928-29 to 1932-33.
48. Cold-storage holdings, year ended December 31, 1933.