# Acreage

National Agricultural Statistics Service

United States Department of Agriculture

Washington, D.C.

Released June 30, 1994, by the Agricultural Statistics Board. Estimates refer to June 1, 1994.

## Corn Acreage Up 7 Percent
## Soybean Acreage Up 4 Percent

**Corn** planted for all purposes is estimated at 78.8 million acres, up 7 percent from last year. Growers expect to harvest 71.8 million acres for grain, up 14 percent from 1993. Excellent weather during planting time allowed producers to complete planting ahead of normal.

**Soybean** growers planted or intend to plant 61.8 million acres in 1994, up 4 percent from 1993. Spring planting got off to a good start with 88 percent of the crop planted by June 5.

**All wheat** seeded is estimated at 70.5 million acres, down 2 percent from last year. Area for grain is expected to total 62.0 million acres, down 1 percent from 1993.

**Winter wheat** planted acres are estimated at 49.5 million acres, down 4 percent from 1993 and 3 percent less than previously published. Grain area is estimated at 41.5 million acres, down 2 percent from the June 1 forecast and off 5 percent from 1993.

**Other spring wheat** plantings are 18.2 million acres, down slightly from 1993. Area for harvest is expected to total 17.8 million acres, up 6 percent from last year's flood and disease reduced level.

Index is located at the end of this report. For information call (202) 720-2127. Office hours are 8:00 a.m. to 4:30 p.m. ET.

Cr Pr 2-2 (6-94)

**All cotton** area planted totals 14.1 million acres, up 5 percent from 1993. Upland cotton acreage accounts for 13.9 million acres of the total, up 5 percent from last year. American-Pima plantings total 178,000 acres, 6 percent below 1993. West Texas planting lagged behind the average pace until late May because of heavy precipitation. Replanting occurred in West Texas, the Delta States, and in the Fresno area of California.

---

**This report was approved on June 30, 1994, by the Secretary of Agriculture and the National Agricultural Statistics Service's Agricultural Statistics Board.**

Secretary of Agriculture
Mike Espy

Agricultural Statistics Board
Chairperson
Rich Allen

Crop Summary: Area Planted and Harvested,
United States, 1993-94
(Domestic Units)

| Crop | Area Planted for All Purposes 1993 | Area Planted for All Purposes 1994 | Area Harvested 1/ 1993 | Area Harvested 1/ 1994 2/ |
|---|---|---|---|---|
| | 1,000 Acres | | | |
| Corn | 73,323.0 | 78,767.0 | 62,991.0 | 71,757.0 |
| Sorghum | 10,492.0 | 10,199.0 | 9,486.0 | 9,297.0 |
| Oats | 7,944.0 | 6,700.0 | 3,793.0 | 4,132.0 |
| Barley | 7,826.0 | 7,307.0 | 6,791.0 | 6,848.0 |
| All Wheat | 72,208.0 | 70,468.0 | 62,647.0 | 61,971.0 |
| Winter | 51,727.0 | 49,547.0 | 43,846.0 | 41,498.0 |
| Durum | 2,191.0 | 2,695.0 | 2,050.0 | 2,639.0 |
| Other Spring | 18,290.0 | 18,226.0 | 16,751.0 | 17,834.0 |
| Rice | 2,920.0 | 3,355.0 | 2,833.0 | 3,300.0 |
| Rye | 1,493.0 | 1,628.0 | 381.0 | 404.0 |
| Soybeans | 59,355.0 | 61,775.0 | 56,447.0 | 60,720.0 |
| Peanuts | 1,733.5 | 1,682.0 | 1,689.8 | 1,657.5 |
| Sunflower | 2,776.0 | 3,430.0 | 2,504.0 | 3,291.0 |
| Canola | 199.0 | 321.0 | 187.0 | 306.0 |
| Mustard Seed | 18.1 | 15.0 | 16.4 | 14.5 |
| Rapeseed | 7.2 | 7.9 | 6.1 | 6.8 |
| Safflower | 404.0 | 255.0 | 292.0 | 239.0 |
| Flaxseed | 206.0 | 189.0 | 191.0 | 182.0 |
| All Cotton | 13,438.3 | 14,066.6 | 12,783.3 | |
| Upland | 13,248.3 | 13,888.6 | 12,594.4 | |
| Amer-Pima | 190.0 | 178.0 | 188.9 | |
| All Hay | | | 60,398.0 | 60,315.0 |
| Alfalfa | | | 24,762.0 | 24,683.0 |
| All Other | | | 35,636.0 | 35,632.0 |
| Dry Edible Beans | 1,849.9 | 2,010.0 | 1,600.0 | 1,939.1 |
| Summer Potatoes | 94.1 | 94.2 | 89.2 | 90.8 |
| Sweetpotatoes | 83.1 | 84.5 | 80.2 | 81.2 |
| All Tobacco | | | 747.5 | 685.2 |
| Sugarbeets | 1,437.7 | 1,469.0 | 1,409.4 | 1,446.9 |
| Sugarcane for Sugar and Seed | | | 948.3 | 952.1 |

1/ Harvested for principal use of each crop, i.e., grain, beans, nuts, etc.
2/ Forecasted.

Crop Summary: Area Planted and Harvested,
United States, 1993-94
(Metric Units)

| Crop | Area Planted for All Purposes | | Area Harvested 1/ | |
|---|---|---|---|---|
| | 1993 | 1994 | 1993 | 1994 2/ |
| | Hectares | | | |
| Corn | 29,673,080 | 31,876,220 | 25,491,830 | 29,039,340 |
| Sorghum | 4,246,010 | 4,127,430 | 3,838,890 | 3,762,400 |
| Oats | 3,214,860 | 2,711,420 | 1,534,990 | 1,672,180 |
| Barley | 3,167,100 | 2,957,070 | 2,748,250 | 2,771,320 |
| All Wheat | 29,221,860 | 28,517,690 | 25,352,610 | 25,079,040 |
| Winter | 20,933,400 | 20,051,180 | 17,744,040 | 16,793,830 |
| Durum | 886,680 | 1,090,640 | 829,610 | 1,067,980 |
| Other Spring | 7,401,780 | 7,375,880 | 6,778,960 | 7,217,240 |
| Rice | 1,181,690 | 1,357,730 | 1,146,490 | 1,335,480 |
| Rye | 604,200 | 658,840 | 154,190 | 163,490 |
| Soybeans | 24,020,370 | 24,999,720 | 22,843,540 | 24,572,780 |
| Peanuts | 701,530 | 680,690 | 683,850 | 670,770 |
| Sunflower | 1,123,420 | 1,388,090 | 1,013,340 | 1,331,830 |
| Canola | 80,530 | 129,910 | 75,680 | 123,840 |
| Mustard Seed | 7,320 | 6,070 | 6,640 | 5,870 |
| Rapeseed | 2,910 | 3,200 | 2,470 | 2,750 |
| Safflower | 163,490 | 103,200 | 118,170 | 96,720 |
| Flaxseed | 83,370 | 76,490 | 77,300 | 73,650 |
| All Cotton | 5,438,350 | 5,692,610 | 5,173,270 | |
| Upland | 5,361,450 | 5,620,580 | 5,096,830 | |
| Amer-Pima | 76,890 | 72,030 | 76,450 | |
| All Hay | | | 24,442,470 | 24,408,880 |
| Alfalfa | | | 10,020,930 | 9,988,960 |
| All Other | | | 14,421,530 | 14,419,910 |
| Dry Edible Beans | 748,640 | 813,430 | 647,500 | 784,730 |
| Summer Potatoes | 38,080 | 38,120 | 36,100 | 36,750 |
| Sweetpotatoes | 33,630 | 34,200 | 32,460 | 32,860 |
| All Tobacco | | | 302,520 | 277,300 |
| Sugarbeets | 581,820 | 594,490 | 570,370 | 585,550 |
| Sugarcane for Sugar and Seed | | | 383,770 | 385,310 |

1/ Harvested for principal use of each crop, i.e., grain, beans, nuts, etc.
2/ Forecasted.

Selected Crops: Historic Area Planted and Harvested, United States, 1985-94 1/

| Year | Corn All Planted | Corn Harvested For Grain | Sorghum All Planted | Sorghum Harvested For Grain | Oats Planted | Oats Harvested |
|---|---|---|---|---|---|---|
| | 1,000 Acres | | | | | |
| 1985 | 83,398 | 75,209 | 18,285 | 16,782 | 13,235 | 8,147 |
| 1986 | 76,580 | 68,907 | 15,339 | 13,862 | 14,671 | 6,840 |
| 1987 | 66,200 | 59,505 | 11,756 | 10,531 | 17,907 | 6,888 |
| 1988 | 67,717 | 58,250 | 10,343 | 9,042 | 13,910 | 5,533 |
| 1989 | 72,221 | 64,703 | 12,642 | 11,103 | 12,085 | 6,882 |
| 1990 | 74,171 | 66,952 | 10,535 | 9,089 | 10,423 | 5,945 |
| 1991 | 75,951 | 68,847 | 11,064 | 9,870 | 8,654 | 4,806 |
| 1992 | 79,340 | 72,162 | 13,277 | 12,150 | 7,961 | 4,492 |
| 1993 | 73,323 | 62,991 | 10,492 | 9,486 | 7,944 | 3,793 |
| 1994 | 78,767 | 71,757 | 10,199 | 9,297 | 6,700 | 4,132 |

| Year | Barley Planted | Barley Harvested | All Wheat Planted | All Wheat Harvested | Winter Wheat Planted | Winter Wheat Harvested |
|---|---|---|---|---|---|---|
| | 1,000 Acres | | | | | |
| 1985 | 13,139 | 11,591 | 75,535 | 64,704 | 57,712 | 47,923 |
| 1986 | 13,024 | 11,974 | 71,998 | 60,688 | 53,895 | 43,170 |
| 1987 | 10,929 | 9,957 | 65,829 | 55,945 | 48,806 | 39,332 |
| 1988 | 9,831 | 7,636 | 65,529 | 53,189 | 48,800 | 39,800 |
| 1989 | 9,125 | 8,313 | 76,615 | 62,189 | 55,091 | 41,509 |
| 1990 | 8,221 | 7,529 | 77,241 | 69,283 | 56,948 | 49,901 |
| 1991 | 8,941 | 8,413 | 69,921 | 57,703 | 51,064 | 39,406 |
| 1992 | 7,809 | 7,325 | 72,264 | 62,411 | 51,057 | 41,893 |
| 1993 | 7,826 | 6,791 | 72,208 | 62,647 | 51,727 | 43,846 |
| 1994 | 7,307 | 6,848 | 70,468 | 61,971 | 49,547 | 41,498 |

| Year | Wheat Durum Planted | Wheat Durum Harvested | Wheat Other Spring Planted | Wheat Other Spring Harvested | Rice Planted | Rice Harvested |
|---|---|---|---|---|---|---|
| | 1,000 Acres | | | | | |
| 1985 | 3,207 | 3,094 | 14,616 | 13,687 | 2,512 | 2,492 |
| 1986 | 2,994 | 2,877 | 15,109 | 14,641 | 2,381 | 2,360 |
| 1987 | 3,341 | 3,279 | 13,682 | 13,334 | 2,356 | 2,333 |
| 1988 | 3,336 | 2,847 | 13,393 | 10,542 | 2,933 | 2,900 |
| 1989 | 3,791 | 3,673 | 17,733 | 17,007 | 2,731 | 2,687 |
| 1990 | 3,570 | 3,507 | 16,723 | 15,875 | 2,897 | 2,823 |
| 1991 | 3,253 | 3,197 | 15,604 | 15,100 | 2,878 | 2,775 |
| 1992 | 2,507 | 2,449 | 18,700 | 18,069 | 3,176 | 3,132 |
| 1993 | 2,191 | 2,050 | 18,290 | 16,751 | 2,920 | 2,833 |
| 1994 | 2,695 | 2,639 | 18,226 | 17,834 | 3,355 | 3,300 |

See footnotes at end of table.

--continued

Selected Crops: Historic Area Planted and Harvested, United States, 1985-94 1/ (continued)

| Year | Rye All Planted | Rye Harvested For Grain | Soybeans All Planted | Soybeans Harvested For Grain | Flaxseed Planted | Flaxseed Harvested |
|---|---|---|---|---|---|---|
| | 1,000 Acres | | | | | |
| 1985 | 2,543 | 708 | 63,145 | 61,599 | 620 | 584 |
| 1986 | 2,334 | 661 | 60,405 | 58,312 | 720 | 683 |
| 1987 | 2,428 | 671 | 58,180 | 57,172 | 470 | 463 |
| 1988 | 2,374 | 595 | 58,840 | 57,373 | 275 | 226 |
| 1989 | 2,014 | 484 | 60,820 | 59,538 | 195 | 163 |
| 1990 | 1,625 | 375 | 57,795 | 56,512 | 260 | 253 |
| 1991 | 1,671 | 396 | 59,180 | 58,011 | 356 | 342 |
| 1992 | 1,582 | 406 | 59,130 | 58,183 | 171 | 165 |
| 1993 | 1,493 | 381 | 59,355 | 56,447 | 206 | 191 |
| 1994 | 1,628 | 404 | 61,775 | 60,720 | 189 | 182 |

| Year | Peanuts Planted | Peanuts Harvested for Nuts | All Sunflower Planted | All Sunflower Harvested |
|---|---|---|---|---|
| | 1,000 Acres | | | |
| 1985 | 1,490.4 | 1,467.4 | 3,055 | 2,844 |
| 1986 | 1,564.7 | 1,535.2 | 2,025 | 1,955 |
| 1987 | 1,567.4 | 1,547.4 | 1,805 | 1,775 |
| 1988 | 1,657.4 | 1,628.4 | 2,038 | 1,921 |
| 1989 | 1,665.2 | 1,644.7 | 1,840 | 1,786 |
| 1990 | 1,840.0 | 1,809.5 | 1,905 | 1,851 |
| 1991 | 2,039.2 | 2,015.7 | 2,746 | 2,673 |
| 1992 | 1,689.6 | 1,672.1 | 2,217 | 2,072 |
| 1993 | 1,733.5 | 1,689.8 | 2,776 | 2,504 |
| 1994 | 1,682.0 | 1,657.5 | 3,430 | 3,291 |

| Year | Cotton Planted | Cotton Harvested | All Hay Harvested | Dry Edible Beans Planted | Dry Edible Beans Harvested |
|---|---|---|---|---|---|
| | 1,000 Acres | | | | |
| 1985 | 10,684.6 | 10,229.0 | 60,461 | 1,569.9 | 1,481.4 |
| 1986 | 10,044.6 | 8,468.4 | 62,334 | 1,653.8 | 1,495.0 |
| 1987 | 10,397.2 | 10,030.3 | 60,133 | 1,782.6 | 1,665.4 |
| 1988 | 12,514.8 | 11,948.2 | 65,055 | 1,485.4 | 1,353.0 |
| 1989 | 10,586.6 | 9,537.7 | 63,300 | 1,824.6 | 1,650.9 |
| 1990 | 12,348.1 | 11,731.6 | 61,407 | 2,177.6 | 2,084.4 |
| 1991 | 14,052.1 | 12,959.5 | 62,475 | 1,964.1 | 1,913.7 |
| 1992 | 13,240.0 | 11,143.3 | 59,597 | 1,640.6 | 1,529.9 |
| 1993 | 13,438.3 | 12,783.3 | 60,398 | 1,849.9 | 1,600.0 |
| 1994 | 14,066.6 | | 60,315 | 2,010.0 | 1,939.1 |

See footnotes at end of table.

--continued

Selected Crops: Historic Area Planted and Harvested, United States, 1985-94 1/ (continued)

| Year | Potatoes Planted | Potatoes Harvested | Sweetpotatoes Planted | Sweetpotatoes Harvested | Tobacco Harvested |
|---|---|---|---|---|---|
| | 1,000 Acres | | | | |
| 1985 | 1,406.7 | 1,358.7 | 108.2 | 103.3 | 688.0 |
| 1986 | 1,256.6 | 1,220.2 | 94.5 | 90.8 | 580.6 |
| 1987 | 1,316.6 | 1,293.4 | 92.3 | 88.9 | 586.3 |
| 1988 | 1,284.7 | 1,259.3 | 89.1 | 85.5 | 634.0 |
| 1989 | 1,305.0 | 1,281.5 | 89.5 | 86.0 | 678.2 |
| 1990 | 1,399.7 | 1,370.6 | 93.9 | 89.5 | 733.3 |
| 1991 | 1,407.5 | 1,374.4 | 81.2 | 77.8 | 763.7 |
| 1992 | 1,339.3 | 1,315.0 | 85.9 | 82.4 | 784.4 |
| 1993 | 1,386.0 | 1,317.8 | 83.1 | 80.2 | 747.5 |
| 1994 | 1,388.9 | 1,324.7 | 84.5 | 81.2 | 685.2 |

| Year | Sugarbeets Planted | Sugarbeets Harvested | Sugarcane for Sugar & Seed Harvested | Principal Crops Planted 2/ | Principal Crops Harvested 3/ |
|---|---|---|---|---|---|
| | 1,000 Acres | | | | |
| 1985 | 1,124.5 | 1,102.5 | 770.0 | 353,042 | 330,255 |
| 1986 | 1,232.5 | 1,192.2 | 796.2 | 338,220 | 310,098 |
| 1987 | 1,266.7 | 1,252.4 | 823.6 | 315,263 | 288,532 |
| 1988 | 1,327.2 | 1,300.7 | 845.3 | 318,319 | 289,275 |
| 1989 | 1,324.4 | 1,294.5 | 851.9 | 331,629 | 305,071 |
| 1990 | 1,400.4 | 1,377.2 | 794.2 | 326,913 | 308,318 |
| 1991 | 1,427.4 | 1,386.7 | 896.9 | 326,032 | 303,864 |
| 1992 | 1,436.7 | 1,411.5 | 925.2 | 327,409 | 307,171 |
| 1993 | 1,437.7 | 1,409.4 | 948.3 | 320,276 | 295,964 |
| 1994 | 1,469.0 | 1,446.9 | 952.1 | 326,198 | 303,183 |

1/ Area harvested forecasted for 1994.

2/ Crops included in area planted are corn, sorghum, oats, barley, winter wheat, rye, durum wheat, other spring wheat, rice, soybeans, peanuts, sunflower, cotton, dry edible beans, potatoes, and sugarbeets. The harvested acreage is used for all hay, tobacco, and sugarcane in computing total area planted. Includes double cropped acres and unharvested small grains planted as cover crops. Fall potatoes carried forward from the previous year for current year totals.

3/ Crops included in area harvested are listed in footnote 2.

Principal Crops: Area Planted, by State and United States, 1992-94 1/ 2/

| State | 1992 | 1993 | 1994 |
|---|---|---|---|
| | 1,000 Acres | | |
| AL | 2,246 | 2,256 | 2,324 |
| AZ | 744 | 710 | 752 |
| AR | 8,310 | 8,425 | 8,150 |
| CA | 4,916 | 4,789 | 5,196 |
| CO | 5,868 | 6,014 | 6,036 |
| CT | 134 | 131 | 128 |
| DE | 531 | 512 | 510 |
| FL | 1,130 | 1,103 | 1,087 |
| GA | 4,039 | 4,065 | 4,294 |
| HI | 68 | 70 | 68 |
| ID | 4,177 | 4,506 | 4,382 |
| IL | 23,940 | 23,325 | 23,800 |
| IN | 12,219 | 12,038 | 12,487 |
| IA | 24,222 | 23,562 | 24,007 |
| KS | 21,916 | 21,869 | 22,407 |
| KY | 5,647 | 5,659 | 5,615 |
| LA | 4,150 | 3,865 | 3,945 |
| ME | 386 | 390 | 384 |
| MD | 1,679 | 1,627 | 1,569 |
| MA | 140 | 138 | 143 |
| MI | 7,059 | 6,922 | 7,115 |
| MN | 19,905 | 19,277 | 20,063 |
| MS | 4,990 | 4,840 | 4,935 |
| MO | 13,201 | 12,826 | 12,836 |
| MT | 9,320 | 9,466 | 9,412 |
| NE | 19,276 | 18,732 | 19,015 |
| NV | 407 | 530 | 517 |
| NH | 105 | 108 | 100 |
| NJ | 446 | 455 | 458 |
| NM | 1,301 | 1,254 | 1,241 |
| NY | 3,325 | 3,236 | 3,264 |
| NC | 4,757 | 4,432 | 4,692 |
| ND | 21,732 | 21,932 | 21,902 |

See footnotes at end of table. --continued

Principal Crops: Area Planted, by State and United States, 1992-94 1/ 2/ (continued)

| State | 1992 | 1993 | 1994 |
|---|---|---|---|
| | 1,000 Acres | | |
| OH | 10,429 | 10,279 | 10,505 |
| OK | 11,112 | 10,777 | 10,869 |
| OR | 2,236 | 2,337 | 2,411 |
| PA | 4,176 | 4,111 | 4,161 |
| RI | 11 | 12 | 11 |
| SC | 1,988 | 1,837 | 2,105 |
| SD | 17,047 | 15,381 | 16,306 |
| TN | 4,522 | 4,640 | 4,592 |
| TX | 23,820 | 22,512 | 22,848 |
| UT | 1,050 | 1,083 | 1,113 |
| VT | 473 | 443 | 431 |
| VA | 2,858 | 2,833 | 2,916 |
| WA | 4,233 | 4,378 | 4,056 |
| WV | 652 | 630 | 647 |
| WI | 8,668 | 8,016 | 8,422 |
| WY | 1,789 | 1,888 | 1,792 |
| US 2/ | 327,409 | 320,276 | 326,198 |

1/ Crops included in area planted are corn, sorghum, oats, barley, winter wheat, rye, durum wheat, other spring wheat, rice, soybeans, peanuts, sunflower, cotton, dry edible beans, potatoes, and sugarbeets. The harvested acreage is used for all hay, tobacco, and sugarcane in computing total area planted. Includes double cropped acres and unharvested small grains planted as cover crops. Fall potatoes are carried forward from the previous year for current year totals.

2/ States do not add to U.S. due to sunflower and sugarbeet unallocated acreage.

Corn: Area Planted and Harvested for Grain by State and United States, 1993-94

| State | Area Planted 1993 | Area Planted 1994 | Area Harvested for Grain 1993 | Area Harvested for Grain 1994 1/ |
|---|---|---|---|---|
| | 1,000 Acres | | | |
| AL | 300 | 300 | 250 | 260 |
| AZ | 19 | 23 | 10 | 13 |
| AR | 100 | 100 | 90 | 90 |
| CA | 400 | 400 | 170 | 180 |
| CO | 940 | 1,000 | 825 | 900 |
| CT 2/ | 45 | 46 | | |
| DE | 165 | 155 | 160 | 152 |
| FL | 140 | 120 | 100 | 80 |
| GA | 650 | 600 | 560 | 540 |
| ID | 125 | 100 | 45 | 30 |
| IL | 10,500 | 11,700 | 10,000 | 11,550 |
| IN | 5,550 | 6,300 | 5,400 | 6,150 |
| IA | 12,000 | 12,700 | 11,000 | 12,400 |
| KS | 2,000 | 2,150 | 1,800 | 2,000 |
| KY | 1,370 | 1,350 | 1,220 | 1,220 |
| LA | 230 | 300 | 210 | 280 |
| ME 2/ | 41 | 38 | | |
| MD | 500 | 460 | 420 | 380 |
| MA 2/ | 32 | 31 | | |
| MI | 2,500 | 2,500 | 2,150 | 2,150 |
| MN | 6,300 | 7,000 | 4,600 | 6,300 |
| MS | 220 | 330 | 190 | 280 |
| MO | 2,200 | 2,400 | 1,850 | 2,300 |
| MT | 65 | 70 | 8 | 25 |
| NE | 8,000 | 8,400 | 7,550 | 8,000 |
| NH 2/ | 18 | 19 | | |
| NJ | 100 | 100 | 80 | 83 |
| NM | 118 | 125 | 85 | 85 |
| NY | 1,150 | 1,110 | 580 | 540 |
| NC | 1,000 | 950 | 850 | 850 |
| ND | 780 | 850 | 365 | 500 |
| OH | 3,500 | 3,750 | 3,280 | 3,500 |
| OK | 170 | 200 | 140 | 180 |
| OR | 42 | 50 | 19 | 25 |
| PA | 1,370 | 1,400 | 970 | 1,000 |
| RI 2/ | 2 | 2 | | |
| SC | 330 | 370 | 240 | 345 |
| SD | 3,350 | 3,800 | 2,550 | 3,300 |
| TN | 660 | 670 | 550 | 570 |
| TX | 2,000 | 2,100 | 1,850 | 1,950 |
| UT | 68 | 67 | 22 | 24 |
| VT 2/ | 93 | 91 | | |
| VA | 490 | 500 | 285 | 340 |
| WA | 120 | 140 | 80 | 95 |
| WV | 75 | 70 | 43 | 40 |
| WI | 3,400 | 3,750 | 2,350 | 3,000 |
| WY | 95 | 80 | 44 | 50 |
| US | 73,323 | 78,767 | 62,991 | 71,757 |

1/ Forecasted.
2/ Area harvested for grain not estimated.

Sorghum: Area Planted and Harvested for Grain
by State and United States, 1993-94

| State | Area Planted | | Area Harvested for Grain | |
|---|---|---|---|---|
| | 1993 | 1994 | 1993 | 1994 1/ |
| | 1,000 Acres | | | |
| AL | 28 | 27 | 19 | 20 |
| AR | 240 | 250 | 215 | 230 |
| CO | 230 | 200 | 190 | 170 |
| GA | 80 | 65 | 40 | 35 |
| IL | 230 | 190 | 210 | 180 |
| KS | 3,000 | 3,000 | 2,800 | 2,800 |
| KY | 14 | 23 | 10 | 20 |
| LA | 130 | 120 | 120 | 110 |
| MS | 70 | 75 | 65 | 70 |
| MO | 720 | 650 | 640 | 620 |
| NE | 1,400 | 1,350 | 1,250 | 1,200 |
| NM | 190 | 190 | 165 | 160 |
| NC | 25 | 35 | 15 | 20 |
| OK | 320 | 320 | 290 | 290 |
| SC | 20 | 24 | 8 | 15 |
| SD | 350 | 250 | 210 | 130 |
| TN | 45 | 30 | 39 | 27 |
| TX | 3,400 | 3,400 | 3,200 | 3,200 |
| US | 10,492 | 10,199 | 9,486 | 9,297 |

1/ Forecasted.

Oats: Area Planted and Harvested by State and United States, 1993-94

| State | Area Planted 1/ | | Area Harvested | |
|---|---|---|---|---|
| | 1993 | 1994 | 1993 | 1994 2/ |
| | 1,000 Acres | | | |
| AL | 60 | 70 | 30 | 38 |
| AR | 35 | 25 | 20 | 20 |
| CA | 300 | 310 | 30 | 35 |
| CO | 87 | 85 | 32 | 30 |
| GA | 75 | 80 | 50 | 50 |
| ID | 70 | 70 | 15 | 25 |
| IL | 550 | 110 | 90 | 90 |
| IN | 130 | 75 | 40 | 40 |
| IA | 950 | 600 | 225 | 450 |
| KS | 70 | 160 | 30 | 120 |
| ME | 33 | 27 | 26 | 21 |
| MD | 10 | 8 | 8 | 6 |
| MI | 150 | 140 | 130 | 110 |
| MN | 850 | 575 | 475 | 475 |
| MO | 35 | 55 | 14 | 30 |
| MT | 140 | 140 | 70 | 80 |
| NE | 270 | 230 | 160 | 160 |
| NY | 135 | 130 | 105 | 100 |
| NC | 60 | 85 | 30 | 40 |
| ND | 800 | 860 | 530 | 520 |
| OH | 200 | 150 | 150 | 120 |
| OK | 80 | 80 | 25 | 30 |
| OR | 65 | 80 | 30 | 55 |
| PA | 220 | 190 | 200 | 170 |
| SC | 50 | 65 | 30 | 40 |
| SD | 750 | 750 | 510 | 560 |
| TX | 800 | 700 | 140 | 150 |
| UT | 50 | 40 | 13 | 10 |
| WA | 65 | 45 | 30 | 20 |
| WV | 9 | 10 | 5 | 5 |
| WI | 790 | 700 | 525 | 500 |
| WY | 55 | 55 | 25 | 32 |
| US | 7,944 | 6,700 | 3,793 | 4,132 |

1/ Includes are planted in preceding fall.
2/ Forecasted.

Barley: Area Planted and Harvested by State and United States, 1993-94

| State | Area Planted 1/ | | Area Harvested | |
|---|---|---|---|---|
| | 1993 | 1994 | 1993 | 1994 2/ |
| | 1,000 Acres | | | |
| AZ | 32 | 35 | 29 | 33 |
| CA | 250 | 290 | 200 | 220 |
| CO | 100 | 90 | 90 | 83 |
| DE | 40 | 35 | 35 | 30 |
| ID | 770 | 740 | 750 | 720 |
| KS | 18 | 15 | 15 | 13 |
| KY | 18 | 16 | 16 | 14 |
| MD | 75 | 65 | 69 | 60 |
| MI | 30 | 35 | 28 | 33 |
| MN | 725 | 650 | 650 | 625 |
| MT | 1,300 | 1,300 | 1,100 | 1,150 |
| NE | 35 | 25 | 32 | 21 |
| NV | 6 | 7 | 5 | 4 |
| NJ | 7 | 7 | 5 | 5 |
| NC | 25 | 30 | 20 | 25 |
| ND | 2,900 | 2,600 | 2,400 | 2,550 |
| OK | 12 | 9 | 8 | 6 |
| OR | 165 | 140 | 150 | 130 |
| PA | 100 | 85 | 95 | 80 |
| SC | 8 | 8 | 7 | 7 |
| SD | 400 | 380 | 360 | 355 |
| TX | 20 | 15 | 7 | 6 |
| UT | 115 | 115 | 110 | 110 |
| VA | 105 | 105 | 85 | 85 |
| WA | 350 | 310 | 345 | 305 |
| WI | 100 | 90 | 70 | 75 |
| WY | 120 | 110 | 110 | 103 |
| US | 7,826 | 7,307 | 6,791 | 6,848 |

1/ Includes are planted in preceding fall.
2/ Forecasted.

All Wheat: Area Planted and Harvested by State
and United States, 1993-94

| State | Area Planted 1/ | | Area Harvested | |
|---|---|---|---|---|
| | 1993 | 1994 | 1993 | 1994 2/ |
| | 1,000 Acres | | | |
| AL | 145 | 140 | 95 | 95 |
| AZ | 95 | 125 | 85 | 123 |
| AR | 1,100 | 980 | 1,000 | 900 |
| CA | 591 | 650 | 540 | 569 |
| CO | 2,835 | 2,945 | 2,583 | 2,642 |
| DE | 65 | 75 | 63 | 70 |
| FL | 40 | 25 | 25 | 20 |
| GA | 400 | 440 | 360 | 400 |
| ID | 1,490 | 1,490 | 1,390 | 1,430 |
| IL | 1,650 | 1,150 | 1,550 | 900 |
| IN | 720 | 680 | 670 | 630 |
| IA | 60 | 55 | 40 | 50 |
| KS | 12,100 | 12,100 | 11,100 | 11,400 |
| KY | 680 | 640 | 470 | 450 |
| LA | 130 | 100 | 95 | 80 |
| MD | 210 | 230 | 200 | 220 |
| MI | 580 | 600 | 540 | 580 |
| MN | 2,755 | 2,680 | 2,298 | 2,627 |
| MS | 250 | 180 | 210 | 160 |
| MO | 1,650 | 1,250 | 1,400 | 1,100 |
| MT | 5,515 | 5,530 | 5,214 | 5,328 |
| NE | 2,350 | 2,200 | 2,100 | 2,000 |
| NV | 11 | 12 | 9 | 9 |
| NJ | 43 | 45 | 33 | 35 |
| NM | 510 | 470 | 270 | 240 |
| NY | 95 | 120 | 85 | 110 |
| NC | 610 | 670 | 560 | 620 |
| ND | 11,700 | 11,440 | 10,800 | 11,285 |
| OH | 1,050 | 1,230 | 1,010 | 1,200 |
| OK | 7,200 | 7,100 | 5,400 | 5,300 |
| OR | 950 | 965 | 925 | 932 |
| PA | 170 | 170 | 165 | 165 |
| SC | 280 | 370 | 260 | 360 |
| SD | 3,820 | 3,620 | 3,488 | 3,319 |
| TN | 490 | 470 | 340 | 290 |
| TX | 6,100 | 6,000 | 3,700 | 3,000 |
| UT | 187 | 194 | 180 | 182 |
| VA | 280 | 280 | 255 | 255 |
| WA | 2,900 | 2,650 | 2,790 | 2,545 |
| WV | 14 | 15 | 11 | 11 |
| WI | 149 | 157 | 125 | 141 |
| WY | 238 | 225 | 213 | 198 |
| US | 72,208 | 70,468 | 62,647 | 61,971 |

1/ Includes area planted in preceding fall.
2/ Forecasted.

Winter Wheat: Area Planted and Harvested by State
and United States, 1993-94

| State | Area Planted 1/ | | Area Harvested | |
|---|---|---|---|---|
| | 1993 | 1994 | 1993 | 1994 2/ |
| | 1,000 Acres | | | |
| AL | 145 | 140 | 95 | 95 |
| AZ | 40 | 30 | 35 | 29 |
| AR | 1,100 | 980 | 1,000 | 900 |
| CA | 550 | 590 | 500 | 510 |
| CO | 2,800 | 2,900 | 2,550 | 2,600 |
| DE | 65 | 75 | 63 | 70 |
| FL | 40 | 25 | 25 | 20 |
| GA | 400 | 440 | 360 | 400 |
| ID | 920 | 840 | 850 | 800 |
| IL | 1,650 | 1,150 | 1,550 | 900 |
| IN | 720 | 680 | 670 | 630 |
| IA | 60 | 55 | 40 | 50 |
| KS | 12,100 | 12,100 | 11,100 | 11,400 |
| KY | 680 | 640 | 470 | 450 |
| LA | 130 | 100 | 95 | 80 |
| MD | 210 | 230 | 200 | 220 |
| MI | 580 | 600 | 540 | 580 |
| MN | 45 | 40 | 40 | 38 |
| MS | 250 | 180 | 210 | 160 |
| MO | 1,650 | 1,250 | 1,400 | 1,100 |
| MT | 2,650 | 1,950 | 2,450 | 1,850 |
| NE | 2,350 | 2,200 | 2,100 | 2,000 |
| NV | 5 | 7 | 4 | 5 |
| NJ | 43 | 45 | 33 | 35 |
| NM | 510 | 470 | 270 | 240 |
| NY | 95 | 120 | 85 | 110 |
| NC | 610 | 670 | 560 | 620 |
| ND | 150 | 40 | 130 | 35 |
| OH | 1,050 | 1,230 | 1,010 | 1,200 |
| OK | 7,200 | 7,100 | 5,400 | 5,300 |
| OR | 880 | 900 | 860 | 870 |
| PA | 170 | 170 | 165 | 165 |
| SC | 280 | 370 | 260 | 360 |
| SD | 1,600 | 1,550 | 1,450 | 1,350 |
| TN | 490 | 470 | 340 | 290 |
| TX | 6,100 | 6,000 | 3,700 | 3,000 |
| UT | 160 | 170 | 155 | 160 |
| VA | 280 | 280 | 255 | 255 |
| WA | 2,600 | 2,400 | 2,500 | 2,300 |
| WV | 14 | 15 | 11 | 11 |
| WI | 135 | 145 | 115 | 130 |
| WY | 220 | 200 | 200 | 180 |
| US | 51,727 | 49,547 | 43,846 | 41,498 |

1/ Includes area planted in preceding fall.
2/ Forecasted.

Durum Wheat: Area Planted and Harvested by State
and United States, 1993-94

| State | Area Planted | | Area Harvested | |
|---|---|---|---|---|
| | 1993 | 1994 | 1993 | 1994 1/ |
| | 1,000 Acres | | | |
| AZ | 55 | 95 | 50 | 94 |
| CA | 41 | 60 | 40 | 59 |
| MN | 10 | 40 | 8 | 39 |
| MT | 115 | 180 | 114 | 178 |
| ND | 1,950 | 2,300 | 1,820 | 2,250 |
| SD | 20 | 20 | 18 | 19 |
| US | 2,191 | 2,695 | 2,050 | 2,639 |

1/ Forecasted.

Other Spring Wheat: Area Planted and Harvested by State
and United States, 1993-94

| State | Area Planted | | Area Harvested | |
|---|---|---|---|---|
| | 1993 | 1994 | 1993 | 1994 1/ |
| | 1,000 Acres | | | |
| CO | 35 | 45 | 33 | 42 |
| ID | 570 | 650 | 540 | 630 |
| MN | 2,700 | 2,600 | 2,250 | 2,550 |
| MT | 2,750 | 3,400 | 2,650 | 3,300 |
| NV | 6 | 5 | 5 | 4 |
| ND | 9,600 | 9,100 | 8,850 | 9,000 |
| OR | 70 | 65 | 65 | 62 |
| SD | 2,200 | 2,050 | 2,020 | 1,950 |
| UT | 27 | 24 | 25 | 22 |
| WA | 300 | 250 | 290 | 245 |
| WI | 14 | 12 | 10 | 11 |
| WY | 18 | 25 | 13 | 18 |
| US | 18,290 | 18,226 | 16,751 | 17,834 |

1/ Forecasted.

Rye: Area Planted and Harvested by State
and United States, 1993-94

| State | Area Planted 1/ | | Area Harvested | |
|---|---|---|---|---|
| | 1993 | 1994 | 1993 | 1994 2/ |
| | 1,000 Acres | | | |
| CO | 11 | 25 | 1 | 2 |
| GA | 300 | 370 | 60 | 80 |
| IL | 40 | 45 | 7 | 7 |
| IN | 25 | 20 | 5 | 5 |
| KS | 70 | 90 | 21 | 15 |
| MD | 30 | 35 | 5 | 4 |
| MI | 80 | 90 | 15 | 17 |
| MN | 30 | 40 | 23 | 30 |
| NE | 100 | 70 | 25 | 20 |
| NJ | 32 | 33 | 7 | 6 |
| NY | 40 | 30 | 8 | 8 |
| NC | 110 | 100 | 30 | 25 |
| ND | 35 | 25 | 30 | 20 |
| OH | 45 | 45 | 5 | 5 |
| OK | 110 | 150 | 30 | 40 |
| PA | 40 | 45 | 10 | 10 |
| SC | 50 | 75 | 20 | 25 |
| SD | 55 | 50 | 50 | 45 |
| TX | 130 | 120 | 11 | 10 |
| VA | 80 | 90 | 5 | 5 |
| WI | 80 | 80 | 13 | 25 |
| US | 1,493 | 1,628 | 381 | 404 |

1/ Includes area planted in preceding fall.
2/ Forecasted.

Rice: Area Planted and Harvested by Class, State, and United States, 1993-94

| Class and State | Area Planted 1993 | Area Planted 1994 | Area Harvested 1993 | Area Harvested 1994 1/ |
|---|---|---|---|---|
| | 1,000 Acres | | | |
| Long Grain | | | | |
| AR | 1,115.0 | 1,199.0 | 1,070.0 | 1,182.0 |
| CA | 14.0 | 10.0 | 14.0 | 10.0 |
| LA | 325.0 | 440.0 | 315.0 | 430.0 |
| MS | 250.0 | 300.0 | 245.0 | 295.0 |
| MO | 105.0 | 120.0 | 93.0 | 112.0 |
| TX | 293.0 | 330.0 | 291.0 | 328.0 |
| US | 2,102.0 | 2,399.0 | 2,028.0 | 2,357.0 |
| Medium Grain | | | | |
| AR | 162.0 | 220.0 | 157.0 | 217.0 |
| CA | 413.0 | 503.0 | 410.0 | 498.0 |
| LA | 220.0 | 210.0 | 215.0 | 205.0 |
| MS 2/ | | | | |
| MO | | | | |
| TX | 7.0 | 10.0 | 7.0 | 10.0 |
| US | 802.0 | 943.0 | 789.0 | 930.0 |
| Short Grain | | | | |
| AR | 3.0 | 1.0 | 3.0 | 1.0 |
| CA | 13.0 | 12.0 | 13.0 | 12.0 |
| US | 16.0 | 13.0 | 16.0 | 13.0 |
| All | | | | |
| AR | 1,280.0 | 1,420.0 | 1,230.0 | 1,400.0 |
| CA | 440.0 | 525.0 | 437.0 | 520.0 |
| LA | 545.0 | 650.0 | 530.0 | 635.0 |
| MS | 250.0 | 300.0 | 245.0 | 295.0 |
| MO | 105.0 | 120.0 | 93.0 | 112.0 |
| TX | 300.0 | 340.0 | 298.0 | 338.0 |
| US | 2,920.0 | 3,355.0 | 2,833.0 | 3,300.0 |

1/ Forecasted.
2/ No medium grain estimated.

Soybeans: Area Planted and Harvested by State and United States, 1993-94

| State | Area Planted 1993 | Area Planted 1994 | Area Harvested 1993 | Area Harvested 1994 1/ |
|---|---|---|---|---|
| | 1,000 Acres | | | |
| AL | 310 | 320 | 295 | 300 |
| AR | 3,500 | 3,300 | 3,450 | 3,250 |
| DE | 220 | 225 | 215 | 220 |
| FL | 55 | 50 | 50 | 48 |
| GA | 600 | 520 | 480 | 510 |
| IL | 9,100 | 9,500 | 8,700 | 9,400 |
| IN | 4,900 | 4,700 | 4,850 | 4,680 |
| IA | 8,500 | 8,900 | 8,200 | 8,850 |
| KS | 1,950 | 2,100 | 1,850 | 2,050 |
| KY | 1,170 | 1,150 | 1,150 | 1,120 |
| LA | 1,300 | 1,200 | 1,250 | 1,150 |
| MD | 580 | 560 | 570 | 550 |
| MI | 1,450 | 1,550 | 1,440 | 1,540 |
| MN | 5,400 | 5,700 | 5,000 | 5,600 |
| MS | 2,000 | 2,000 | 1,950 | 1,900 |
| MO | 4,200 | 4,600 | 3,550 | 4,540 |
| NE | 2,600 | 2,900 | 2,500 | 2,850 |
| NJ | 150 | 150 | 147 | 147 |
| NC | 1,300 | 1,400 | 1,200 | 1,350 |
| ND | 600 | 620 | 540 | 610 |
| OH | 4,100 | 3,900 | 4,040 | 3,850 |
| OK | 280 | 300 | 260 | 280 |
| PA | 300 | 320 | 295 | 315 |
| SC | 600 | 640 | 520 | 620 |
| SD | 1,800 | 2,450 | 1,700 | 2,400 |
| TN | 1,050 | 1,100 | 990 | 1,050 |
| TX | 230 | 220 | 205 | 200 |
| VA | 500 | 540 | 470 | 520 |
| WI | 610 | 860 | 580 | 820 |
| US | 59,355 | 61,775 | 56,447 | 60,720 |

1/ Forecasted.

Soybeans: Percent of Acreage Planted Following Another Crop, Selected States and United States, 1990-94 1/

| State | 1990 | 1991 | 1992 | 1993 | 1994 | State | 1990 | 1991 | 1992 | 1993 | 1994 |
|---|---|---|---|---|---|---|---|---|---|---|---|
| AL | 31 | 17 | 27 | 38 | 29 | MS | 20 | 13 | 8 | 9 | 4 |
| AR | 39 | 28 | 27 | 30 | 26 | MO | 15 | 11 | 11 | 13 | 10 |
| DE | 26 | 51 | 53 | 48 | 42 | NJ | 22 | 22 | 21 | 16 | 18 |
| FL | 44 | 24 | 27 | 29 | 21 | NC | 29 | 27 | 34 | 32 | 33 |
| GA | 46 | 49 | 44 | 49 | 55 | OH | 1 | 1 | 2 | 1 | 1 |
| IL | 4 | 4 | 4 | 3 | 2 | OK | 29 | 18 | 23 | 16 | 27 |
| IN | 4 | 4 | 4 | 3 | 4 | PA | 8 | 19 | 17 | 17 | 19 |
| KS | 6 | 14 | 15 | 12 | 10 | SC | 44 | 39 | 40 | 39 | 47 |
| KY | 36 | 39 | 39 | 33 | 33 | TN | 36 | 29 | 31 | 32 | 22 |
| LA | 15 | 12 | 9 | 2 | 2 | TX | 0 | 0 | 7 | 0 | 0 |
| MD | 33 | 36 | 41 | 40 | 37 | VA | 49 | 59 | 55 | 57 | 54 |
| | | | | | | US | 11 | 9 | 9 | 8 | 8 |

1/ Data as obtained from area frame samples. These data do not represent official estimates of the Agricultural Statistics Board but provide raw data as obtained from survey respondents. The purpose of these data is to portray trends in soybean production practices.

Peanuts for Nuts: Area Planted and Harvested by State and United States, 1993-94

| State | Area Planted | | Area Harvested | |
|---|---|---|---|---|
| | 1993 | 1994 | 1993 | 1994 1/ |
| | 1,000 Acres | | | |
| AL | 240.0 | 237.0 | 239.0 | 236.0 |
| FL | 98.0 | 83.0 | 84.0 | 75.0 |
| GA | 702.0 | 680.0 | 697.0 | 675.0 |
| NM | 22.0 | 23.0 | 21.8 | 23.0 |
| NC | 152.0 | 151.0 | 143.0 | 150.0 |
| OK | 105.0 | 110.0 | 102.0 | 107.0 |
| SC | 14.5 | 15.0 | 14.0 | 14.5 |
| TX | 305.0 | 290.0 | 295.0 | 285.0 |
| VA | 95.0 | 93.0 | 94.0 | 92.0 |
| US | 1,733.5 | 1,682.0 | 1,689.8 | 1,657.5 |

1/ Forecasted.

Sunflower: Area Planted and Harvested by Type, State, and United States, 1993-94

| Varietal Type and State | Area Planted 1993 | Area Planted 1994 | Area Harvested 1993 | Area Harvested 1994 1/ |
|---|---|---|---|---|
| | 1,000 Acres | | | |
| Oil | | | | |
| CO | 60 | 72 | 54 | 69 |
| KS | 140 | 170 | 133 | 160 |
| MN | 300 | 400 | 260 | 390 |
| NE | 39 | 40 | 35 | 39 |
| ND | 1,100 | 1,500 | 970 | 1,430 |
| SD | 630 | 670 | 601 | 651 |
| TX | 13 | 10 | 12 | 10 |
| Oth Sts | 35 | 41 | 28 | 37 |
| US | 2,317 | 2,903 | 2,093 | 2,786 |
| Non-Oil | | | | |
| CO | 25 | 33 | 23 | 31 |
| KS | 40 | 90 | 37 | 85 |
| MN | 90 | 100 | 80 | 95 |
| NE | 23 | 15 | 22 | 15 |
| ND | 220 | 220 | 195 | 215 |
| SD | 20 | 30 | 19 | 29 |
| TX | 20 | 15 | 17 | 14 |
| Oth Sts | 21 | 24 | 18 | 21 |
| US | 459 | 527 | 411 | 505 |
| All | | | | |
| CO | 85 | 105 | 77 | 100 |
| KS | 180 | 260 | 170 | 245 |
| MN | 390 | 500 | 340 | 485 |
| NE | 62 | 55 | 57 | 54 |
| ND | 1,320 | 1,720 | 1,165 | 1,645 |
| SD | 650 | 700 | 620 | 680 |
| TX | 33 | 25 | 29 | 24 |
| Oth Sts | 56 | 65 | 46 | 58 |
| US | 2,776 | 3,430 | 2,504 | 3,291 |

1/ Forecasted.

Flaxseed: Area Planted and Harvested by State and United States, 1993-94

| State | Area Planted 1993 | Area Planted 1994 | Area Harvested 1993 | Area Harvested 1994 1/ |
|---|---|---|---|---|
| | 1,000 Acres | | | |
| MN | 15 | 10 | 10 | 10 |
| ND | 165 | 150 | 156 | 145 |
| SD | 20 | 20 | 19 | 18 |
| Other States | 6 | 9 | 6 | 9 |
| US | 206 | 189 | 191 | 182 |

1/ Forecasted.

Special Oilseeds: Area Planted and Harvested, United States, 1993-94

| Crop | Area Planted 1993 | Area Planted 1994 | Area Harvested 1993 | Area Harvested 1994 1/ |
|---|---|---|---|---|
| | 1,000 Acres | | | |
| Canola | 199.0 | 321.0 | 187.0 | 306.0 |
| Rapeseed | 7.2 | 7.9 | 6.1 | 6.8 |
| Safflower | 404.0 | 255.0 | 292.0 | 239.0 |
| Mustard Seed | 18.1 | 15.0 | 16.4 | 14.5 |

1/ Forecasted.

Cotton: Area Planted and Harvested by Type, State, and United States, 1993-94

| Type and State | Area Planted 1993 | Area Planted 1994 | Area Harvested 1993 | Area Harvested 1994 1/ |
|---|---|---|---|---|
| | 1,000 Acres | | | |
| Upland | | | | |
| AL | 443.0 | 470.0 | 430.0 | |
| AZ | 316.0 | 315.0 | 315.0 | |
| AR | 990.0 | 950.0 | 970.0 | |
| CA | 1,050.0 | 1,090.0 | 1,045.0 | |
| FL | 54.0 | 70.0 | 53.5 | |
| GA | 615.0 | 850.0 | 600.0 | |
| KS | 1.6 | 1.6 | 1.4 | |
| LA | 890.0 | 900.0 | 875.0 | |
| MS | 1,330.0 | 1,300.0 | 1,300.0 | |
| MO | 345.0 | 380.0 | 335.0 | |
| NM | 53.5 | 60.0 | 48.7 | |
| NC | 390.0 | 490.0 | 385.0 | |
| OK | 370.0 | 400.0 | 350.0 | |
| SC | 202.0 | 210.0 | 198.0 | |
| TN | 625.0 | 560.0 | 615.0 | |
| TX | 5,550.0 | 5,800.0 | 5,050.0 | |
| VA | 23.2 | 42.0 | 22.8 | |
| US | 13,248.3 | 13,888.6 | 12,594.4 | |
| Amer-Pima | | | | |
| AZ | 57.0 | 48.0 | 56.9 | |
| CA | 91.0 | 80.0 | 91.0 | |
| NM | 11.0 | 15.0 | 11.0 | |
| TX | 31.0 | 35.0 | 30.0 | |
| US | 190.0 | 178.0 | 188.9 | |
| All | | | | |
| AL | 443.0 | 470.0 | 430.0 | |
| AZ | 373.0 | 363.0 | 371.9 | |
| AR | 990.0 | 950.0 | 970.0 | |
| CA | 1,141.0 | 1,170.0 | 1,136.0 | |
| FL | 54.0 | 70.0 | 53.5 | |
| GA | 615.0 | 850.0 | 600.0 | |
| KS | 1.6 | 1.6 | 1.4 | |
| LA | 890.0 | 900.0 | 875.0 | |
| MS | 1,330.0 | 1,300.0 | 1,300.0 | |
| MO | 345.0 | 380.0 | 335.0 | |
| NM | 64.5 | 75.0 | 59.7 | |
| NC | 390.0 | 490.0 | 385.0 | |
| OK | 370.0 | 400.0 | 350.0 | |
| SC | 202.0 | 210.0 | 198.0 | |
| TN | 625.0 | 560.0 | 615.0 | |
| TX | 5,581.0 | 5,835.0 | 5,080.0 | |
| VA | 23.2 | 42.0 | 22.8 | |
| US | 13,438.3 | 14,066.6 | 12,783.3 | |

1/ Estimates to be released August 11, 1994.

Hay: Area Harvested by Type, State, and United States
*1993 and Forecasted 1994*

| State | All Hay 1993 | All Hay 1994 | Alfalfa and Alfalfa Mixtures 1993 | Alfalfa and Alfalfa Mixtures 1994 | All Other 1993 | All Other 1994 |
|---|---|---|---|---|---|---|
| | 1,000 Acres | | | | | |
| AL 1/ | 720 | 750 | | | 720 | 750 |
| AZ | 185 | 200 | 150 | 160 | 35 | 40 |
| AR | 1,180 | 1,125 | 30 | 25 | 1,150 | 1,100 |
| CA | 1,380 | 1,510 | 920 | 950 | 460 | 560 |
| CO | 1,400 | 1,350 | 850 | 860 | 550 | 490 |
| CT | 84 | 80 | 20 | 19 | 64 | 61 |
| DE | 17 | 15 | 5 | 4 | 12 | 11 |
| FL 1/ | 220 | 240 | | | 220 | 240 |
| GA 1/ | 600 | 650 | | | 600 | 650 |
| ID | 1,330 | 1,250 | 1,050 | 1,020 | 280 | 230 |
| IL | 1,250 | 1,100 | 790 | 700 | 460 | 400 |
| IN | 700 | 700 | 330 | 350 | 370 | 350 |
| IA | 2,050 | 1,750 | 1,550 | 1,250 | 500 | 500 |
| KS | 2,450 | 2,500 | 850 | 850 | 1,600 | 1,650 |
| KY | 2,200 | 2,250 | 280 | 300 | 1,920 | 1,950 |
| LA 1/ | 250 | 280 | | | 250 | 280 |
| ME | 235 | 238 | 19 | 19 | 216 | 219 |
| MD | 210 | 200 | 65 | 60 | 145 | 140 |
| MA | 103 | 108 | 31 | 33 | 72 | 75 |
| MI | 1,500 | 1,550 | 1,200 | 1,200 | 300 | 350 |
| MN | 2,250 | 2,300 | 1,600 | 1,600 | 650 | 700 |
| MS 1/ | 720 | 750 | | | 720 | 750 |
| MO | 3,560 | 3,370 | 460 | 370 | 3,100 | 3,000 |
| MT | 2,370 | 2,300 | 1,500 | 1,550 | 870 | 750 |
| NE | 3,650 | 3,500 | 1,400 | 1,500 | 2,250 | 2,000 |
| NV | 505 | 490 | 235 | 240 | 270 | 250 |
| NH | 90 | 81 | 15 | 16 | 75 | 65 |
| NJ | 120 | 120 | 30 | 30 | 90 | 90 |
| NM | 325 | 335 | 255 | 265 | 70 | 70 |
| NY | 1,750 | 1,800 | 700 | 680 | 1,050 | 1,120 |
| NC | 470 | 510 | 20 | 20 | 450 | 490 |
| ND | 2,950 | 2,900 | 1,700 | 1,550 | 1,250 | 1,350 |
| OH | 1,350 | 1,400 | 650 | 680 | 700 | 720 |
| OK | 2,130 | 2,200 | 330 | 320 | 1,800 | 1,880 |
| OR | 1,040 | 1,100 | 420 | 420 | 620 | 680 |
| PA | 1,880 | 1,920 | 780 | 800 | 1,100 | 1,120 |
| RI | 9 | 8 | 2 | 2 | 7 | 6 |
| SC 1/ | 230 | 280 | | | 230 | 280 |
| SD | 4,200 | 4,300 | 2,300 | 2,500 | 1,900 | 1,800 |
| TN | 1,700 | 1,700 | 60 | 50 | 1,640 | 1,650 |
| TX | 3,485 | 3,690 | 85 | 90 | 3,400 | 3,600 |
| UT | 650 | 685 | 500 | 525 | 150 | 160 |
| VT | 350 | 340 | 90 | 95 | 260 | 245 |
| VA | 1,200 | 1,210 | 130 | 150 | 1,070 | 1,060 |
| WA | 750 | 720 | 480 | 480 | 270 | 240 |
| WV | 530 | 550 | 40 | 40 | 490 | 510 |
| WI | 2,800 | 2,700 | 2,200 | 2,250 | 600 | 450 |
| WY | 1,270 | 1,210 | 640 | 660 | 630 | 550 |
| US | 60,398 | 60,315 | 24,762 | 24,683 | 35,636 | 35,632 |

1/ Alfalfa and alfalfa mixtures included in all other hay.

Dry Edible Beans: Area Planted and Harvested by State and United States, 1993-94 1/

| State | Area Planted | | Area Harvested | |
|---|---|---|---|---|
| | 1993 | 1994 | 1993 | 1994 2/ |
| | 1,000 Acres | | | |
| CA | 107.0 | 160.0 | 104.0 | 155.0 |
| CO | 205.0 | 215.0 | 185.0 | 205.0 |
| ID | 125.0 | 140.0 | 123.0 | 138.0 |
| KS | 29.0 | 30.0 | 27.0 | 28.0 |
| MI | 390.0 | 400.0 | 380.0 | 390.0 |
| MN | 110.0 | 125.0 | 88.0 | 120.0 |
| MT | 12.0 | 8.5 | 11.0 | 8.3 |
| NE | 170.0 | 190.0 | 130.0 | 180.0 |
| NM | 14.0 | 12.5 | 14.0 | 12.5 |
| NY | 37.0 | 45.0 | 34.0 | 43.0 |
| ND | 510.0 | 540.0 | 380.0 | 520.0 |
| OR | 9.0 | 10.0 | 8.9 | 9.9 |
| TX | 30.0 | 30.0 | 27.0 | 27.0 |
| UT | 6.4 | 6.0 | 6.1 | 5.6 |
| WA | 43.0 | 41.0 | 42.0 | 41.0 |
| WI | 10.5 | 10.0 | 10.0 | 9.8 |
| WY | 42.0 | 47.0 | 30.0 | 46.0 |
| US | 1,849.9 | 2,010.0 | 1,600.0 | 1,939.1 |

1/ Excludes beans grown for garden seed.
2/ Forecasted.

Sweetpotatoes: Area Planted and Harvested by State and United States, 1993-94

| State | Area Planted 1993 | Area Planted 1994 | Area Harvested 1993 | Area Harvested 1994 1/ |
|---|---|---|---|---|
| | 1,000 Acres | | | |
| AL | 4.5 | 4.3 | 4.4 | 4.2 |
| CA | 8.3 | 8.4 | 8.3 | 8.4 |
| GA | 3.2 | 2.8 | 3.0 | 2.6 |
| LA | 17.0 | 19.0 | 16.5 | 18.0 |
| MD | 0.3 | 0.3 | 0.3 | 0.3 |
| MS | 6.0 | 6.0 | 5.5 | 5.5 |
| NJ | 1.5 | 1.5 | 1.4 | 1.4 |
| NC | 33.0 | 34.0 | 32.0 | 33.0 |
| SC | 2.4 | 2.0 | 2.2 | 1.9 |
| TX | 6.3 | 5.7 | 6.0 | 5.4 |
| VA | 0.6 | 0.5 | 0.6 | 0.5 |
| US | 83.1 | 84.5 | 80.2 | 81.2 |

1/ Forecasted.

Summer Potatoes: Area Planted and Harvested by State and United States, 1993-94

| State | Area Planted 1993 | Area Planted 1994 | Area Harvested 1993 | Area Harvested 1994 1/ |
|---|---|---|---|---|
| | 1,000 Acres | | | |
| AL | 7.3 | 7.2 | 7.1 | 7.0 |
| CA | 4.8 | 5.2 | 4.8 | 5.2 |
| CO | 8.3 | 9.2 | 8.2 | 9.0 |
| DE | 5.0 | 4.6 | 5.0 | 4.6 |
| IL | 5.1 | 5.0 | 4.5 | 4.6 |
| IA | 1.5 | 1.7 | 0.7 | 1.7 |
| MD | 2.5 | 2.2 | 2.3 | 2.0 |
| MI | 13.0 | 14.0 | 12.5 | 13.5 |
| MN | 7.2 | 7.7 | 7.1 | 7.6 |
| MO | 7.7 | 7.2 | 6.7 | 6.7 |
| NE | 3.1 | 3.5 | 3.0 | 3.4 |
| NJ | 3.4 | 2.9 | 3.3 | 2.8 |
| NM | 4.9 | 4.5 | 4.3 | 4.5 |
| NC | 1.3 | 1.5 | 1.2 | 1.4 |
| TX | 8.0 | 7.8 | 7.5 | 7.3 |
| VA | 11.0 | 10.0 | 11.0 | 9.5 |
| US | 94.1 | 94.2 | 89.2 | 90.8 |

1/ Forecasted.

Alaska: Area Planted by Crop, 1992-94 1/

| Crop | Area Planted | | |
|---|---|---|---|
| | 1992 | 1993 | 1994 |
| | Acres | | |
| All Oats | 2,000 | 2,000 | 2,400 |
| All Barley | 3,500 | 4,700 | 6,900 |
| Grain Hay and Silage 2/ 3/ | 600 | 800 | 1,300 |
| Grass Hay and Silage 3/ | 16,400 | 20,200 | 22,500 |
| Potatoes | 830 | 850 | 820 |

1/ Estimates are provided to meet special needs of users for crops and livestock production statistics. Estimates are excluded from commodity data tables.
2/ Included in the above grain crop estimates.
3/ Area harvested.

Tobacco: Area Harvested by State and United States, 1992-93 and Forecasted 1994

| State | Area Harvested | | |
|---|---|---|---|
| | 1992 | 1993 | 1994 |
| | Acres | | |
| CT | 1,450 | 1,500 | 1,610 |
| FL | 7,500 | 7,100 | 7,000 |
| GA | 44,000 | 43,000 | 39,000 |
| IN | 9,000 | 8,100 | 7,100 |
| KY | 234,700 | 207,300 | 186,400 |
| MD | 9,700 | 9,500 | 8,500 |
| MA | 370 | 400 | 460 |
| MO | 1,700 | 2,800 | 3,300 |
| NC | 270,200 | 271,000 | 252,500 |
| OH | 10,400 | 9,000 | 8,500 |
| PA | 10,800 | 10,000 | 9,600 |
| SC | 52,000 | 52,000 | 48,000 |
| TN | 72,200 | 69,940 | 62,440 |
| VA | 51,100 | 49,100 | 45,500 |
| WV | 2,100 | 2,000 | 2,000 |
| WI | 7,200 | 4,800 | 3,300 |
| US | 784,420 | 747,540 | 685,210 |

Tobacco: Area Harvested by Class, Type, State, and United States, 1993 and Forecasted 1994

| Class and Type | Area Harvested 1993 | Area Harvested 1994 |
|---|---|---|
| | Acres | |
| Class 1, Flue-Cured | | |
| Type 11, Old and Middle Belts | | |
| NC | 99,000 | 93,000 |
| VA | 36,000 | 33,000 |
| US | 135,000 | 126,000 |
| Type 12, Eastern NC Belt | | |
| NC | 129,000 | 120,000 |
| Type 13, NC Border & SC Belt | | |
| NC | 34,000 | 31,000 |
| SC | 52,000 | 48,000 |
| US | 86,000 | 79,000 |
| Type 14, GA-FL Belt | | |
| FL | 7,100 | 7,000 |
| GA | 43,000 | 39,000 |
| US | 50,100 | 46,000 |
| Total 11-14 | 400,100 | 371,000 |
| Class 2, Fire-Cured | | |
| Type 21, VA Belt | | |
| VA | 1,200 | 1,400 |
| Type 22, Eastern District | | |
| KY | 3,500 | 3,800 |
| TN | 7,700 | 8,200 |
| US | 11,200 | 12,000 |
| Type 23, Western District | | |
| KY | 3,650 | 3,800 |
| TN | 600 | 640 |
| US | 4,250 | 4,440 |
| Total 21-23 | 16,650 | 17,840 |
| Class 3, Air-Cured | | |
| Class 3A, Light Air-Cured | | |
| Type 31, Burley | | |
| IN | 8,100 | 7,100 |
| KY | 196,000 | 175,000 |
| MO | 2,800 | 3,300 |
| NC | 9,000 | 8,500 |
| OH | 9,000 | 8,500 |
| TN | 61,000 | 53,000 |
| VA | 11,800 | 11,000 |
| WV | 2,000 | 2,000 |
| US | 299,700 | 268,400 |
| Type 32, Southern MD Belt | | |
| MD | 9,500 | 8,500 |
| PA | 3,500 | 3,600 |
| US | 13,000 | 12,100 |
| Total 31-32 | 312,700 | 280,500 |
| Class 3, Air-Cured | | |
| Class 3B, Dark Air-Cured | | |
| Type 35, One Sucker Belt | | |
| KY | 2,700 | 2,500 |
| TN | 640 | 600 |
| US | 3,340 | 3,100 |
| Type 36, Green River Belt | | |
| KY | 1,450 | 1,300 |
| Type 37, VA Sun-Cured Belt | | |
| VA | 100 | 100 |
| Total 35-37 | 4,890 | 4,500 |
| Class 4, Cigar Filler | | |
| Type 41, PA Seedleaf | | |
| PA | 6,500 | 6,000 |
| Class 5, Cigar Binder | | |
| Class 5A, CT Valley Binder | | |
| Type 51, CT Valley Broadleaf | | |
| CT | 650 | 700 |
| MA | 170 | 180 |
| US | 820 | 880 |
| Class 5B, WI Binder | | |
| Type 54, Southern WI | | |
| WI | 2,900 | 2,200 |
| Type 55, Northern WI | | |
| WI | 1,900 | 1,100 |
| Total 54-55 | 4,800 | 3,300 |
| Total 51-55 | 5,620 | 4,180 |
| Class 6, Cigar Wrapper | | |
| Type 61, CT Valley Shade-grown | | |
| CT | 850 | 910 |
| MA | 230 | 280 |
| US | 1,080 | 1,190 |
| All Cigar Types | | |
| Total 41-61 | 13,200 | 11,370 |
| All Tobacco | 747,540 | 685,210 |

Sugarbeets: Area Planted and Harvested by State
and United States, 1993-94 1/

| State | Area Planted | | Area Harvested | |
|---|---|---|---|---|
| | 1993 | 1994 | 1993 | 1994 2/ |
| | 1,000 Acres | | | |
| CA | 138.0 | 139.0 | 136.0 | 137.0 |
| CO | 40.3 | 44.0 | 40.0 | 43.3 |
| ID | 206.0 | 202.0 | 204.0 | 200.0 |
| MI | 189.0 | 196.0 | 187.0 | 193.0 |
| MN | 390.0 | 415.0 | 379.0 | 410.0 |
| MT | 54.4 | 54.4 | 54.1 | 54.4 |
| NE | 82.3 | 82.3 | 79.6 | 76.0 |
| ND | 193.8 | 204.0 | 190.9 | 203.0 |
| OH | 19.1 | 16.1 | 17.5 | 16.0 |
| OR | 16.0 | 16.0 | 15.2 | 15.5 |
| TX | 40.3 | 25.5 | 39.2 | 25.0 |
| WY | 66.0 | 63.0 | 64.4 | 62.0 |
| Oth Sts 3/ | 2.5 | 11.7 | 2.5 | 11.7 |
| US | 1,437.7 | 1,469.0 | 1,409.4 | 1,446.9 |

1/ Relates to year of intended harvest except for overwintered spring planted beets in CA.
2/ Forecasted.
3/ Includes NM and WA.

Sugarcane for Sugar and Seed: Area Harvested by State
and United States, 1993 and Forecasted 1994

| State | Area Harvested | |
|---|---|---|
| | 1993 | 1994 |
| | 1,000 Acres | |
| FL | 444.0 | 445.0 |
| HI | 69.9 | 67.7 |
| LA | 390.0 | 395.0 |
| TX | 44.4 | 44.4 |
| US | 948.3 | 952.1 |

## Spring Weather Review

### March-May 1994

Spring began and ended dry across the central Plains and the Corn Belt, broken by a wet April (except in eastern Nebraska and Iowa). In the southern Plains, several months of dry weather yielded abruptly to early-May downpours. Farther east, the Southeastern States turned dry during the spring for the second consecutive year. Areas from the middle Mississippi and Ohio River Valleys into the Northeast weathered a minor bout with spring flooding as conditions gradually turned drier. The Pacific Northwest closed out a drier-than-normal wet season with sub-par spring rains.

All 3 months featured above-normal temperatures in the Northwestern States, resulting in spring departures of +4 degrees F or more from southern Idaho to western Nebraska. Normal to below-normal spring temperatures were confined to parts of a strip from Texas to New England.

**March:** Overall atmospheric features from the 1993-94 winter months (west coast ridge and Great Lakes trough) remained intact. As a result, wet weather persisted from the Tennessee Valley to the Northeast, while dryness prevailed in the West. California had its 14th driest March since 1895, marking an early end to a disappointing wet season (about 65 percent of normal precipitation). It was the driest March on record for Idaho, Montana, Wyoming, and Nebraska. Farther east, monthly precipitation exceeded twice normal and totaled 8 inches or more in many locations from eastern Tennessee to New Jersey. One storm, between March 1 and 4, left a fresh 1- to 2-foot blanket of snow in the central and northern Appalachians. A late-month storm pushed seasonal snowfall at Scranton, PA, and Elkins, WV, to record magnitudes, but will be remembered for its tornado outbreak that left more than 40 people dead across the Southeast on March 27.

Monthly temperatures averaged above normal except in parts of the Northeast. Near-record warmth, with departures commonly +4 to +6 degrees F, covered areas from the northern Plains southward into the Southwest.

**April:** Stormy weather encompassed a large part of the Nation, but temperatures remained mostly above normal. Acute wetness shifted a bit westward, striking the middle Mississippi and lower Ohio Valleys. Rainfall, in combination with above-normal Ohio Basin runoff, brought short-lived minor to moderate flooding to the middle Mississippi Valley and points downstream. Several late-season snowstorms plagued the central third of the Nation, including one that left a depth of 9 inches in Sioux Falls, SD, on April 29. In contrast, dryness developed in the southern Atlantic Coast States.

Although temperatures averaged only 1 to 2 degrees F below normal in the Plains, occasional wintry blasts persisted. Even late in the month (April 27), Amarillo, TX, reported 30 degrees F and Tribune, KS, measured 21 degrees F. Monthly temperatures averaged 4 degrees F or more above normal in the interior Northwest and in many areas south of the Ohio River.

**May:** Long-term moisture deficits were vanquished in the southern High Plains, but dry weather returned to the central Plains and the Corn Belt. Dryness continued in the Southeast, where a single, early-month storm broke an otherwise dry pattern. The month's most vigorous and slow-moving storm spent 2 weeks spinning its way across the Southwestern and South Central States. The system unleashed excessive rainfall on the southern High Plains between May 9 and 12.

A trough became re-established over the Nation's northeastern quadrant, permitting cool air to slide southeastward across the Midwest and the East. Monthly temperatures averaged 2 to 4 degrees F below normal in a broad strip from New York to Texas. An offshore storm roiled Atlantic coastal waters and brought exceptionally cool weather to the Eastern States for several days beginning at mid-month. Farther west, over the High Plains and the Northwest, a building ridge resulted in several hot days. Monthly temperature departures topped +5 degrees F from southern Idaho to western Nebraska.

**Spring Fieldwork:** Planting progress for Spring 1994 was one of the fastest on record. The spring began with snowstorms and melting snow hindering field preparation but by mid-March warm, dry weather allowed producers to progress rapidly. Spring storms in early April left fields too wet to prepare for planting and triggered concern of a repeat of last year's problems. By the end of April warm, windy weather dried fields and allowed farmers to begin planting. Early May storm systems saturated some fields and hindered planting but did not slow planting. Poor germination from cool, wet weather in mid-May and then later in the month from hot, dry weather had little effect on planting progress. All the right conditions for crop development existed until later in the spring when dry soil conditions began to increase concern for field crops.

In early March, snow hit the south-central and Atlantic Seaboard States depositing significant amounts of snow which restricted field work. The storm systems left fields too wet to allow planting to begin, which left field preparation behind schedule. A warming trend, starting in the Pacific Coast and moving eastward early in the month, melted the snow pack for winter wheat. The warm weather triggered wheat growth in the upper Great Plains, but the rapid snow melt produced some flooding and soil erosion. In mid-month, the Southeastern half of the Nation experienced warm, windy weather, which helped dry fields and allowed field preparation and planting to start. The small grains responded to the mild weather as winter wheat broke out of dormancy in the lower High Plains. For most of the month, the Central Plains remained below the desired soil-moisture level for wheat growers. In Kansas, some slight insect and disease problems were reported. By the end of the month, drier Spring weather let farmers plant and prepare fields in the Ohio Valley and Mid-western regions. Early planting and field preparation proceeded rapidly in the latter half of the month.

In early April, spring storm systems hit the central Plains and lessened dry soil moisture conditions. In the Tennessee and Ohio Valleys, saturated fields hampered spring planting. Producers in the Corn Belt planted wherever and whenever possible despite rain and wet fields. Corn planting began with progress equal to the five-year average, while all other field crops started ahead of the five-year average. Small grains in the Great Plains were rescued by mid-month rains, while dry conditions in the Pacific Coast continued with some fields of small grains abandoned. A late month snow storm slowed fieldwork

and left the soil temperatures below the desired level for corn planting. Despite the cool weather in the Great Plains that limited crop development, planting proceeded. In the middle Mississippi Valley, late month thunderstorms restrained corn planting progress to average levels. By month's end, corn planting was ahead of the average prompted in part by the threat of a cool, wet spring that spurred producers to plant early and avoid a repeat of last year's problems.

In early May, many producers were waiting for fields to dry so spring planting could resume. Rain and cool weather in the Mississippi and Ohio Valleys interrupted planting progress and slowed crop development and stressed some early emerged plants. The Nation's row crops began the month of May ahead the five-year average and the dry weather allowed producers to maintain their lead. The small grains seeding started the month slightly behind the average and remained there. The summer-like weather stimulated wheat development in the middle Great Plains causing winter wheat to turn color early. Winter wheat harvest started in the Southernmost States slightly ahead of average. Cool weather lowered some germination success in the Corn Belt. By mid-month, a warm weather system brought summer-like temperatures that dried fields and boosted crop development. Continued hot, dry weather later in the month inhibited some germination of row crops in the Midwest. The Central Plains and Southeastern States reported short soil moisture levels for the month's end. By month's end corn, rice, and spring wheat plantings were completed. With corn planted, producers turned their attention to planting soybeans. Continued dry weather in the Corn Belt allowed farmers to plant soybeans at the brisk pace set by corn producers. Producers planted soybeans early despite cool weather early in the month that threatened germination. The month began with cool, damp conditions that slowed cotton germination and development in Texas. Cotton planting doubled during the month starting at 42 percent and ending at 84 percent completed.

**Corn:** Corn planted for all purposes is estimated at 78.8 million acres, up 7 percent from 1993. Growers expect to harvest 71.8 million acres for grain, up 14 percent from the flood reduced acreage of 1993. Most of the increase in corn planted acreage can be attributed to the decrease from 10 percent to zero percent of the Acreage Reduction Program of the 1994 Feed Grain Program. Excellent weather during planting time allowed farmers to get the crop in ahead of normal and match earlier expectations of acreage.

As of June 5, corn planting was virtually complete, compared with a 93 percent average. Corn condition was rated mostly fair to good, with 73 percent rated good to excellent. Last year 51 percent of the crop was rated good to excellent at the same time.

**Sorghum:** Acreage planted for all purposes in 1994 is expected to total 10.2 million acres, down 3 percent from 1993. This will be the smallest area devoted to sorghum since 1930. Area for grain is placed at 9.30 million acres, down 2 percent from a year ago.

Planting progress for this year's crop had reached 95 percent complete in the 12 major producing States as of June 19. The four largest producing States (Kansas, Missouri, Nebraska, and Texas) total 8.40 million acres planted and 7.82 million acres for grain. Both levels are down 1 percent from last season.

Late seeded fields need moisture in Kansas. Favorable weather allowed Nebraska producers to complete planting about 10 days ahead of average.

**Oats:** Oats planted last fall and this spring totaled 6.70 million acres, down 16 percent from 1993. Acreage seeded in Illinois, Iowa, and Minnesota dropped sharply as corn growers participating in the Feed Grain Program will have a zero acreage reduction requirement for corn (ARP) and will not be seeding oats as a cover crop.

Growers intend to harvest 4.13 million acres for grain in 1994, up 9 percent from the 3.79 million acres harvested in 1993.

**Barley:** Seedings last fall and this spring totaled 7.31 million acres, down 7 percent from last year. North Dakota continues to lead all States with 2.60 million acres seeded, down 10 percent from a year earlier. The area to be harvested for grain is expected to total 6.85 million acres, up 1 percent from last year.

**All Wheat:** Seeded area for 1994 is estimated at 70.5 million acres, down 2 percent from last year. Area for grain is expected to total 62.0 million acres, down 1 percent from 1993.

**Winter Wheat:** Seeded area is 49.5 million acres, down 4 percent from 1993 and off 3 percent from previously published 1994 levels. Grain area is placed at 41.5 million acres, 2 percent less than the June 1 forecast and down 5 percent from last year.

**Durum Wheat:** Planted area for 1994 is estimated at 2.70 million acres; 2.64 million acres are expected to be harvested for grain. The planted level is up 23 percent from 1993 and the grain area is up 29 percent. North Dakota's plantings finished in early June. Crop development trails average progress. Condition ratings are ahead of last year. Arizona's durum harvest reached 86 percent by mid-June; California's Imperial Valley harvest is nearly complete.

**Other Spring Wheat:** Seedings this spring total 18.2 million acres, down slightly from 1993. Growers intend to harvest 17.8 million acres for grain, up 6 percent from a year ago. Minnesota and North Dakota's harvested areas are up from their flood and disease reduced 1993 crops. Montana's increase is largely due to an even more reduced winter seeded area than previously thought.

**Rye:** Planted area is estimated at 1.63 million acres, up 9 percent from 1993. Area for grain is estimated at 404,000 acres, 6 percent more than last year. Harvested acreage increased in Georgia and Minnesota but decreased in Nebraska, North Dakota, and South Dakota.

**Rice:** Area planted to rice is estimated at 3.36 million acres in the six major producing States, 15 percent above 1993. Acreage increased in all six States. Expected area for harvest is 3.30 million acres, up 16 percent from last year.

Area planted to long grain rice, representing 72 percent of the total, rose 14 percent. Medium grain acreage increased 18 percent and accounted for 28 percent of the total. The acreage planted to short grain varieties dropped 19 percent and continued to make up a small amount of the all rice total.

Rice planting got off to a good start in the six States and by May 29 most States had completed planting. Rice condition in late June was rated as mostly good to fair.

**Soybeans:** Growers planted or intend to plant 61.8 million acres in 1994, up 4 percent from 1993. Area to be harvested is estimated at 60.7 million acres this year.

Hot, dry weather allowed farmers to plant more soybeans than expected. South Dakota farmers planted 650,000 acres more than last year. Illinois, Iowa, and Missouri acreages were up 400,000 acres each from a year ago. Both Minnesota and Nebraska increased 300,000 acres from 1993. Overall, growers in 18 States are planting more acreage this year than last, 9 States are planting less, and 2 States are estimating no change from 1993.

Planting progress by May 29 in the 19 major States was 78 percent complete, 26 points ahead of the five-year average. By June 12, 91 percent of the soybean acreage was planted, 12 points ahead of the average. Soybean condition in late June was rated as mostly good to fair.

**Peanuts:** Producers planted 1.68 million acres of peanuts this year, down 3 percent from the 1993 planted area of 1.73 million acres and slightly below the 1992 acreage. Acres declined in all States except South Carolina, New Mexico, and Oklahoma.

Area for harvest is estimated at 1.66 million acres, down 2 percent from the 1993 level of 1.69 million acres. If realized, North and South Carolina, New Mexico, and Oklahoma will exceed their harvested acreage of 1993.

Southeast growers (Alabama, Florida, Georgia, and South Carolina) planted 1.02 million acres, 4 percent less than last year but slightly above 1992. Planting ended slightly ahead of normal. Generally favorable conditions prevailed placing most of the crop in good condition. Frequent showers in early June improved crop development after a brief dry period in late May. As of June 19, development in Georgia was ahead of last year and normal with 73 percent blooming compared with 55 percent last year and 59 percent normally. Over three-fourths of the acreage was rated in good condition. Alabama producers rated most of the crop in good condition. The Florida and South Carolina crops are mostly in good condition with near normal development.

In the Virginia-North Carolina region, producers planted 244,000 acres, 1 percent less than the last two years. Planting finished slightly behind normal but nearly all the crop was rated in good condition as of mid-June. Some replanting was necessary in Virginia.

Acreage planted in the Southwest (New Mexico, Oklahoma, and Texas) totaled 423,000 acres, 2 percent below 1993 and 1 percent below 1992. The Oklahoma crop rated 57 percent good and 43 percent fair by early June. Planting was

90 percent complete by June 19, well ahead of the 74 percent average for the date. Planting in New Mexico ended under typical dry, windy conditions but with no damaging weather. In Texas, early planted peanuts emerged in good condition. Excess moisture slowed planting in South Central Texas, while in the Cross Timbers, area planting was delayed due to dry conditions.

**Sunflower:** Planted area is estimated at 3.43 million acres in 1994, 24 percent above 1993 plantings. Oil type varieties total 2.90 million acres this year, 25 percent above 1993. Non-oil type varieties were planted on 527,000 acres, up 15 percent from the previous year.

Planting progress has been ahead of the average most of the season for North Dakota, South Dakota, and Minnesota. North Dakota continues to lead all States with 1.72 million acres seeded. Crop condition in North Dakota was mostly good by mid-June.

**Flaxseed:** Acreage seeded for 1994 is estimated at 189,000 acres, down 8 percent from 1993. Area harvested is estimated at 182,000 acres, down 5 percent from last year.

In North Dakota, planting started slowly but by June 5 was equal to or ahead of the average. As of June 19, the majority of the crop was emerged and in mostly good condition in North Dakota.

**Special Oilseeds:** Planted area of canola is estimated at 321,000 acres, up 61 percent from 1993. Farmers intend to harvest 306,000 acres, up 64 percent from last year. Planted acreage of rapeseed is estimated at 7,900 acres, up 10 percent from 1993. Growers expect to harvest 6,800 acres, up 11 percent from a year ago. Area planted for safflower is estimated at 255,000 acres, down 37 percent from 1993. Area for harvest is expected to total 239,000 acres, down 18 percent from last year. Mustard seed growers planted 15,000 acres this year, down 17 percent from 1993. Harvested acres are estimated at 14,500, down 12 percent from 1993.

**Cotton:** Area planted to all cotton for 1994 totaled 14.1 million acres, 5 percent above 1993 plantings and 6 percent above the 1992 area. Upland cotton accounts for 13.9 million acres, up 5 percent from last year. Growers decreased plantings of American-Pima cotton to 178,000 acres, 6 percent below last year's amount.

Upland growers in the Delta States (Arkansas, Louisiana, Mississippi, Missouri, and Tennessee) planted 4.09 million acres. This total declined 2 percent from 1993 and is 3 percent below two years ago. Early precipitation caused Missouri and Tennessee producers to trail their average planting pace. By mid-May, fields had dried and progress equalled or exceeded normal. Some acreage in Arkansas suffered the effects of cool, damp weather during late May and was replanted. Although 22 percent of the Arkansas crop and 11 percent of the Missouri crop were rated poor in early June, all of the region rated fair to good by late June, as beneficial weather improved conditions.

Texas and Oklahoma producers planted 6.20 million acres, a 5 percent increase from last year and 6 percent greater than two years ago. Early in the season, South Texas plantings exceeded the average pace. Because of heavy rains during

early May. West Texas planting lagged until late May. Some replanting was necessary in the Plains due to heavy rains. In late June, 80 percent of the Texas crop was rated fair to good, and 14 percent was rated in excellent condition. About half of the Oklahoma acreage was rated fair and half was in good condition.

In the Southeast (Alabama, Georgia, North Carolina, and South Carolina), producers planted 2.02 million acres, up 22 percent from 1993 and 39 percent greater than 1992. Alabama planting progress was behind the normal rate throughout the season due to cool temperatures and dry soils. In early June, 17 percent of the North Carolina crop and 30 percent of the South Carolina crop were in poor condition. Condition improved late in the month, as 98 percent of North Carolina's acreage and 99 percent of South Carolina's crop were in fair to good condition.

Upland acreage in the Western States (Arizona, California, and New Mexico) is estimated at 1.47 million acres, a 3 percent increase from last year and up 6 percent from 1992. Seeding rates in Arizona and California equalled or exceeded the average pace. In late April, heavy rains in the Fresno area caused replanting, and this activity was hampered to some extent because of cool temperatures and wet soil. In early June, the desert crop was blooming but germination was slow in the San Joaquin Valley. In late June, crop development in that valley remained one week behind normal. Ninety-five percent of the State's crop was in fair to good condition. Arizona conditions improved slightly during the month. By late June, 89 percent of the crop was in good to excellent condition.

American-Pima plantings decreased from 1993 in Arizona and California by 16 and 12 percent, respectively. These two States account for 72 percent of the acreage. Acreage in New Mexico and Texas increased from 1993 by 36 percent and 13 percent, respectively.

**Hay:** Growers expect to harvest 60.3 million acres of hay in 1994, down slightly from 1993. The area harvested of alfalfa and alfalfa mixtures cut for hay is estimated at 24.7 million acres, virtually unchanged from 1993. Acreage of all other hay harvested is estimated at 35.6 million acres, virtually unchanged from last year.

Harvesting of first cutting hay had a good start with most States near completion and some beginning their second cutting. Hay conditions for the Nation have been mostly fair to good. Hot, dry weather has forced some producers to feed hay to livestock.

**Dry Beans:** Planted acreage of dry beans is estimated at 2.01 million acres this year, up 9 percent from last year and 23 percent above two years ago. Area for harvest is forecast at 1.94 million acres, up 21 percent from a year ago when considerable acreage was rained and flooded out in North Central States. Dry bean acreage increased from last year in nearly all major producing States.

California's planted acreage jumped 50 percent from last year on the prospects of better prices. Oregon increased 11 percent. Idaho and Wyoming gained 12 percent each. Colorado's acres are up 5 percent. Most beans in the Intermountain West are as much as 20 percent ahead of their normal growing schedule.

Washington is the only major dry bean State with fewer acres than last year. Montana, Utah, and New Mexico are also down. In the North Central States, acreage is up substantially except in Wisconsin and growth is ahead of normal. Minnesota's acreage is up 14 percent. Nebraska jumped 12 percent with some beans going in on abandoned corn and sugarbeet land. North Dakota's plantings increased 6 percent over last year and growing progress is good. Farther east, Michigan growers increased plantings by 3 percent. Progress in Michigan is 20 percent ahead of average. New York's acreage jumped 22 percent.

**Sweetpotatoes:** Planted area totaled 84,500 acres this year, up 2 percent from last year but was 2 percent below 1992. Acreage increased in the three leading States: North Carolina, Louisiana, and California. Acreage declined in Alabama, Georgia, South Carolina, Texas, and Virginia. Area for harvest is forecast at 81,200 acres, a gain of 1 percent from 1993 but 1 percent below 1992.

Planting is generally finished except for small areas across the Southeast. North Carolina's planting was 88 percent complete by June 20. Crop condition is good. Planting in South Carolina and Alabama started early but slowed in June because of hot, dry weather. Planting in Louisiana was ahead of schedule with mostly favorable conditions throughout the spring. By June 20, planting was 71 percent complete. Planting in nearby Mississippi was slow up to June 1. The California crop started slow because of cool, wet conditions and early weed problems but picked up in the last couple of weeks.

**Summer Potatoes:** Growers planted 94,200 acres this year, up slightly from last year and 6 percent above two years ago. Acreage for harvest is forecast at 90,800 acres, 2 percent above last year and 6 percent above 1992. Summer potato acreage is lower in the Atlantic Coast, Mississippi Valley, and Southwest but higher in the Midwest and West.

Growers along the Atlantic Coast reduced acreage from 15 percent in New Jersey to 8 percent in Delaware. North Carolina is up 200 acres on their relatively small acreage. Crop prospects look good on the Delmarva Peninsula as temperatures and precipitation were good from March through June. Harvest began on the Virginia Peninsula in the week ending June 26 with good size and quality. Alabama growers have a good crop enhanced by favorable weather. Michigan potato acreage is up 8 percent from last year as chip deals are gaining on fresh market. Minnesota acreage is up 7 percent. Iowa and Nebraska jumped 13 percent each. Planted acreage in Illinois is down but growers expect to harvest more than last year. Missouri has a good crop. Colorado summer acreage jumped again this year as a new sorting shed was built in Yuma County. New Mexico and Texas acreages are off but good crops are expected. California summer potato acreage is 8 percent above last year. Harvest should start in early July.

**Tobacco:** The Nation's total area of tobacco for harvest in 1994 is estimated at 685,210 acres, 8 percent below last year and 13 percent below 1992.

Flue cured acreage, at 371,000 acres, is down 7 percent from last year. Planting started on schedule in Virginia. With rain received in mid-June, the crop condition was rated as fair to good. Florida and Georgia reported dry weather early in the season and that harvest was underway by mid-June.

Burley acreage, at 268,400 acres, dropped 10 percent from a year ago. Dry soil conditions resulted in higher than normal loss this year for late set plants, requiring some fields to be reset in Kentucky.

Dark fire-cured types are estimated at 17,840 acres, up 7 percent from a year ago. Acreage for cigar types is estimated at 11,370 acres, down 14 percent from 1993. Tobacco condition was rated as fair to good.

**Sugarbeets:** Growers planted an estimated 1.47 million acres of sugarbeets for 1994, up 2 percent from last year. Planted acres increased 6 percent in Minnesota, 5 percent in North Dakota, and 4 percent in Michigan. Growers in Idaho reported a decrease in planted acres of 2 percent. Ohio and Texas reported decreases of 16 and 37 percent, respectively. Dry conditions in California prevailed over most of the season resulting in new crop sugarbeets being irrigated. Planting in Colorado got off to a good start but a late April freeze caused nearly half of the acreage to be replanted. Replanting in Nebraska, due to hailstorms, resulted in smaller plants that were unable to survive the hot, dry, windy weather later in the spring. In the upper Great Plains, dry conditions allowed planting to begin ahead of last year but wet weather later in the spring slowed planting progress to finish the same as last year.

**Sugarcane for Sugar and Seed:** Growers intend to harvest 952,100 acres of sugarcane in 1994, up slightly from last year and if realized will be a record high. The increase from last year resulted from Louisiana's continued expansion in sugarcane production. Florida growers expect to harvest 445,000 acres, compared to 444,000 acres harvested in 1993. Sugarcane acres for harvest in Hawaii were down 3 percent from 1993, reflecting the withdrawal from sugarcane production by some growers. Overall, condition of the crop is rated as fair to good.

## Reliability of Acreage Data in This Report

**Survey Procedures:** The estimates of planted and harvested acreages in this report are based primarily on surveys conducted the first 2 weeks of June. These surveys are based on a probability area frame survey with a sample of over 15,500 segments or parcels of land (average approximately 1 square mile) and a probability list sample of over 70,000 farm operators. Enumerators conducting the area survey contact all farmers having operations within the sampled segments of land and account for their operations. From these data, estimates can be calculated. The list survey sample is contacted by mail, telephone, or personal interviews to obtain information on these operations. Responses from the list sample plus data from the area operations that were not on the list to be sampled are combined to provide another estimate of planted and harvested acreages.

**Estimating Procedures:** National, Regional, State, and grower reported data were reviewed for reasonableness and consistency with general cultural practices, farm legislation, and historical estimates. The survey estimates were also reviewed considering weather patterns and planting progress. Each State Statistical Office submits their analysis of the current situation to the Agricultural Statistics Board (ASB). Planted acreage estimates were based on survey estimates and the historical relationship of official estimates to survey estimates. Harvested acreage estimates were based on survey estimates and the historical relationship between planted and harvested acres.

**Revision Policy:** Planted acreage estimates are subject to revision August 1 if actual plantings are significantly different than those reported in early June. Also, planted acreage estimates can be reviewed at the end of the season and again the following year, if new information is available that would justify a change. Harvested acres can be adjusted anytime a change is made in planted acres. In addition, harvested acres are subject to change anytime a production forecast is made. Estimates will also be reviewed after data for the 5-year Census of Agriculture, conducted by the Department of Commerce, are available. No revisions will be made after this date.

**Reliability:** The surveys used to make acreage estimates are subject to sampling and non-sampling type errors that are common to all surveys. Sampling errors for major crops generally are between 1 and 5 percent. Sampling errors represent the variability between estimates that would result if many different samples were surveyed at the same time. Sampling errors cannot be applied directly to the acreage published in this report to determine confidence intervals since the official estimates represent a composite of information from more than a single source. The sampling errors from the 1994 area frame survey for U.S. planted acres were: barley 4.4 percent, corn 1.0 percent, upland cotton 3.3 percent, sorghum 5.6 percent, soybeans 1.1 percent, winter wheat 1.7 percent, and other spring wheat 3.1 percent.

Non-sampling errors cannot be measured directly but may occur due to planting intentions, incorrect reporting, and/or recording, data omissions or duplications and errors in processing. To minimize non-sampling errors, vigorous quality controls are used in the data collection process and all data are carefully reviewed for consistency and reasonableness.

A method of evaluating the reliability of production forecasts in this report is the **"Root Mean Square Error"**, a statistical measure based on past performances shown below for selected crops. This is computed by expressing the deviations between the mid-year acreage estimates and the final estimates as a percent of the final estimates and averaging the squared percentage deviations for the 1974-1993 20-year period; the square root of this average becomes statistically the "Root Mean Square Error." Probability statements can be made concerning expected differences in the current estimates relative to the final estimates assuming that factors affecting this year's estimate are not different than those influencing the past 20 years.

For example, the "Root Mean Square Error" for the corn planted estimate is 1.2 percent. This means that chances are 2 out of 3 that the current acreage estimate of 78.8 million acres will not be above or below the final estimate by *more than 1.2 percent or approximately 945,204 acres. Chances are 9 out of 10* (90 percent confidence level) that difference will not exceed 2.1 percent or approximately 1.65 million acres.

Also shown in the table is a 10-year record for selected crops of the difference between the mid-year planted acres estimate and the final estimates. Using corn *again as an example, changes between the mid-year* estimates and the final estimates during the past 10 years have averaged 340,000 acres ranging from 10,000 acres to 1,179,000 acres. The mid-year planted acres have been below the final estimate 6 times and above 4 times. This does not imply that the mid-year planted estimate this year is likely to understate or overstate the final estimate.

Reliability of Mid-Year Planted Acreage Estimates

| Crop | Root Mean Square Error | | | 10 Year Record of Differences Between Mid-Year and Final Estimates | | | | |
|---|---|---|---|---|---|---|---|---|
| | | 90% Percent Confidence Level | | Thousand Acres | | | Number of Years | |
| | Percent | Percent | Thousand Acres | Average | Small | Large | Below Final | Above Final |
| Corn | 1.2 | 2.1 | 1,654 | 340 | 10 | 1,179 | 6 | 4 |
| Sorghum | 3.4 | 5.9 | 602 | 424 | 10 | 1,060 | 6 | 4 |
| Oats | 2.8 | 4.8 | 322 | 77 | 3 | 185 | 4 | 6 |
| Barley | 2.6 | 4.5 | 329 | 169 | 10 | 907 | 5 | 5 |
| Winter Wheat | .60 | 1.0 | 495 | 174 | 15 | 410 | 2 | 8 |
| Spring Wheat | 1.1 | 1.9 | 346 | 88 | 0 | 250 | 4 | 5 |
| Soybeans | 1.4 | 2.4 | 1,483 | 634 | 150 | 2,220 | 3 | 7 |
| Upland Cotton | 2.8 | 4.9 | 681 | 179 | 35 | 369 | 3 | 7 |

## Index


| | Page | |
|---|---|---|
| | Table | Narrative |
| Area Planted and Harvested, U.S. 1985-93 | A- 5 | |
| Alaska | A-27 | |
| Area Planted, by States | A- 8 | |
| Barley | A-13 | B-4 |
| Beans, Dry Edible | A-25 | B-7 |
| Canola | A-22 | B-6 |
| Corn | A-10 | B-3 |
| Cotton | A-23 | B-6 |
| Crop Summary | A- 3 | |
| Flaxseed | A-22 | B-6 |
| Hay | A-24 | B-7 |
| Mustard Seed | A-22 | B-6 |
| Oats | A-12 | B-4 |
| Peanuts | A-20 | B-5 |
| Potatoes, Summer | A-26 | B-8 |
| Rapeseed | A-22 | B-6 |
| Rice | A-18 | B-4 |
| Rye | A-17 | B-4 |
| Safflower | A-22 | B-6 |
| Sorghum | A-11 | B-3 |
| Soybeans | A-19 | B-5 |
| Special Oilseeds | | B-6 |
| Sugarbeets | A-29 | B-9 |
| Sugarcane for Sugar and Seed | A-29 | B-9 |
| Sunflower | A-21 | B-6 |
| Sweetpotatoes | A-26 | B-8 |
| Tobacco, by Class and Type | A-28 | B-8 |
| Tobacco, by States | A-27 | B-8 |
| Wheat, All | A-14 | B-4 |
| Wheat, Durum | A-16 | B-4 |
| Wheat, Other Spring | A-16 | B-4 |
| Wheat, Winter | A-15 | B-4 |

## Report Features

*The next* **"Acreage"** *report will be released in June 1995.*

Listed below are the commodity specialists in the Crops Branch of the National Agricultural Statistics Service to contact for additional information.

| | | |
|---|---|---|
| John D. Witzig, Chief | (202) | 720-2127 |
| Field Crops Section | | |
| Bill Dowdy, Head | (202) | 720-3843 |
| Dan Kerestes - Soybeans, Minor Oilseeds, Rice | (202) | 720-9526 |
| Greg Preston - Sugar, Tobacco, Hay, Weekly Crop Weather | (202) | 720-7621 |
| Vaughn Siegenthaler - Rye, Sorghum, Wheat | (202) | 720-8068 |
| Charles Van Lahr - Barley, Corn, Oats, Pasture Condition | (202) | 720-7369 |
| Fruit, Vegetable & Special Crops Section | | |
| Stephen Ropel, Head | (202) | 720-3843 |
| Jim Brewster - Fruits, Fresh Vegetables | (202) | 720-7688 |
| *Arvin Budge - Potatoes, Dry Beans, Onions* | (202) | 720-4285 |
| Kirby Cavett - Peanuts | (202) | 720-8843 |
| Roger Latham - Cotton | (202) | 720-5944 |
| David Mueller - Fresh and Processing Vegetables | (202) | 720-6054 |
| Blair Smith - Fruits, Citrus, Nuts | (202) | 720-5412 |

The next "Acreage" report will be released at 3 p.m. ET in June 1995.

**********

The United States Department of Agriculture (USDA) prohibits discrimination in its programs on the basis of race, color, national origin, sex, religion, age, disability, political beliefs, and marital or familial status. Persons with disabilities who require alternative means for communication of program information (braille, large print, audiotape, etc.) should contact the USDA Office of Communications at (202) 720-5881 (voice) or (202) 720-7808 (TDD).

To file a complaint, write the Secretary of Agriculture, USDA, Washington, D.C., 20250, or call (202) 720-7327 (voice) or (202) 720-1127 (TDD). USDA is an equal employment opportunity employer.

## ☑ Want to subscribe? ☑ Time to renew?

Subscribe to *Crop Production* **today!** If you already subscribe to this timely periodical, note that expiration information about your subscription appears on the top line of your mailing label. **Expiration information will appear in one of two formats:** 1-PCP-2 (this means you have TWO issues remaining in your subscription) or APR95 (expiration date is April 1995). Disregard this notice if no renewal information appears.

**Need it fast?** Subscribe to *Crop Production* via CALL-ERS/NASS, an electronic Bulletin Board Service that supports 1200/2400/9600 baud communications. Reports may be accessed by 9 a.m. ET the day following release. An electronic subscription costs the same as paper!

**Call toll free, 1-800-999-6779,** and tell us whether you want to subscribe or renew, or return this form to: **ERS-NASS, 341 Victory Drive, Herndon, VA 22070.**

### Crop Production (PCP)

| | 1 Year | 2 Years | 3 Years |
|---|---|---|---|
| Domestic | ☐ $49.00 | ☐ $96.00 | ☐ $144.00 |
| Foreign | ☐ $61.25 | ☐ $120.00 | ☐ $180.00 |

☐ Yes! I want to **start** my subscription.
☐ Yes! I want to **renew** my subscription.

**Check one:** ☐ Paper
☐ Electronic (1-year subscriptions only)
☐ BOTH (rate is DOUBLE the listed price)

**New subscribers:**

Name: _____

Address: _____

City, State, Zip: _____

Daytime phone: (_____) _____

**Renewals:**

ATTACH MAILING LABEL HERE

**Payment method:**

☐ Enclosed is $_____ Use purchase orders, checks drawn on U.S. banks, cashier's checks, or international money orders. **Make payable to ERS-NASS.**

**Credit card orders:** ☐ MasterCard ☐ Visa Total charges $_____.

Credit card number: ☐☐☐☐☐☐☐☐☐☐☐☐☐☐☐☐ Card expiration date: ☐☐ Month/Year

**For *fastest* service, call our toll-free order desk 1-800-999-6779, in the U.S. and Canada; other areas please call 703-834-0125, or FAX this page to 703-834-0110.**